

上部消化管検査判定結果

判定	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
	異常なし	男	14,305 (86.6%)	10	603	2944	4539	4166	1861
	女	8,333 (88.5%)	2	215	1634	2799	2592	985	106
	計	22,638 (87.3%)	12	818	4578	7338	6758	2846	288
放置可	男	871 (5.3%)	0	2	94	247	351	157	20
	女	394 (4.2%)	0	1	50	140	149	48	6
	計	1,265 (4.9%)	0	3	144	387	500	205	26
要経過観察	男	607 (3.7%)	0	4	67	145	261	116	14
	女	340 (3.6%)	1	4	36	119	120	49	11
	計	947 (3.7%)	1	8	103	264	381	165	25
要精密検査	男	662 (4.0%)	0	3	68	170	246	155	20
	女	325 (3.5%)	0	6	45	118	105	44	7
	計	987 (3.8%)	0	9	113	288	351	199	27
要治療	男	22 (0.1%)	0	0	3	4	10	4	1
	女	5 (0.1%)	0	0	0	1	4	0	0
	計	27 (0.1%)	0	0	3	5	14	4	1
治療中	男	56 (0.3%)	0	1	2	7	19	23	4
	女	15 (0.2%)	0	0	0	2	10	1	2
	計	71 (0.3%)	0	1	2	9	29	24	6
合計	男	16,523	10	613	3178	5112	5053	2316	241
	女	9,412	3	226	1765	3179	2980	1127	132
	計	25,935	13	839	4943	8291	8033	3443	373

胃内視鏡検査所見

所見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
	食道裂孔ヘルニア	男	182 (21.1%)	0	0	15	60	75	27
	女	21 (5.8%)	0	0	5	7	6	3	0
	計	203 (16.6%)	0	0	20	67	81	30	5
逆流性食道炎	男	147 (17.0%)	0	0	17	44	70	14	2
	女	18 (5.0%)	0	0	1	6	8	2	1
	計	165 (13.5%)	0	0	18	50	78	16	3
粘膜下腫瘍	男	2 (0.2%)	0	0	0	1	1	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	2 (0.2%)	0	0	0	1	1	0	0
ポリープ	男	6 (0.7%)	0	0	1	2	3	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	6 (0.5%)	0	0	1	2	3	0	0
その他の所見	男	39 (4.5%)	0	0	2	6	24	6	1
	女	18 (5.0%)	0	0	2	9	7	0	0
	計	57 (4.7%)	0	0	4	15	31	6	1
慢性胃炎	男	485 (56.2%)	0	1	47	118	240	72	7
	女	188 (52.2%)	0	0	14	63	84	25	2
	計	673 (55.0%)	0	1	61	181	324	97	9
萎縮性胃炎	男	50 (5.8%)	0	0	2	9	27	8	4
	女	16 (4.4%)	0	0	0	5	7	1	3
	計	66 (5.4%)	0	0	2	14	34	9	7
びらん性胃炎	男	32 (3.7%)	0	0	4	4	16	7	1
	女	11 (3.1%)	0	0	2	1	8	0	0
	計	43 (3.5%)	0	0	6	5	24	7	1
タコイボびらん	男	5 (0.6%)	0	0	0	3	1	1	0
	女	3 (0.8%)	0	0	0	1	1	1	0
	計	8 (0.7%)	0	0	0	4	2	2	0
胃潰瘍	男	9 (1.0%)	0	0	0	1	6	2	0
	女	1 (0.3%)	0	0	0	1	0	0	0
	計	10 (0.8%)	0	0	0	2	6	2	0
胃潰瘍瘢痕	男	16 (1.9%)	0	0	0	2	11	1	2
	女	2 (0.6%)	0	0	0	1	1	0	0
	計	18 (1.5%)	0	0	0	3	12	1	2
ポリープ	男	128 (14.8%)	0	0	13	44	52	17	2
	女	88 (24.4%)	0	0	13	37	32	6	0
	計	216 (17.7%)	0	0	26	81	84	23	2
粘膜下腫瘍	男	11 (1.3%)	0	0	0	1	8	2	0
	女	6 (1.7%)	0	0	2	1	2	1	0
	計	17 (1.4%)	0	0	2	2	10	3	0
その他の所見	男	108 (12.5%)	0	1	14	27	50	13	3
	女	30 (8.3%)	0	0	2	6	14	8	0
	計	138 (11.3%)	0	1	16	33	64	21	3
十二指腸潰瘍	男	8 (0.9%)	0	0	0	0	5	3	0
	女	2 (0.6%)	0	0	0	0	1	1	0
	計	10 (0.8%)	0	0	0	0	6	4	0
十二指腸潰瘍瘢痕	男	50 (5.8%)	0	0	7	7	21	13	2
	女	11 (3.1%)	0	0	2	2	5	2	0
	計	61 (5.0%)	0	0	9	9	26	15	2
慢性十二指腸炎	男	31 (3.6%)	0	0	6	8	14	3	0
	女	1 (0.3%)	0	0	0	0	1	0	0
	計	32 (2.6%)	0	0	6	8	15	3	0
ポリープ	男	18 (2.1%)	0	0	0	4	12	2	0
	女	1 (0.3%)	0	0	0	0	1	0	0
	計	19 (1.6%)	0	0	0	4	13	2	0
憩室	男	1 (0.1%)	0	0	0	0	0	1	0
	女	1 (0.3%)	0	0	0	0	0	1	0
	計	2 (0.2%)	0	0	0	0	0	2	0
その他の所見	男	60 (7.0%)	0	0	4	19	29	6	2
	女	14 (3.9%)	0	0	3	3	5	3	0
	計	74 (6.1%)	0	0	7	22	34	9	2
検診者総数	男	863	0	2	109	247	386	105	14
	女	360	0	1	55	126	138	35	5
	計	1,223	0	3	164	373	524	140	19