

胃部×線検査所見

所見		総数								所見		総数									
		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上				
食道部										胃部											
拡張	男	1	(0.0%)	0	0	0	0	0	1	0	胃下垂	男	0	(0.0%)	0	0	0	0	0	0	
	女	0	(0.0%)	0	0	0	0	0	0	0		女	9	(0.1%)	0	0	3	4	0	1	1
	計	1	(0.0%)	0	0	0	0	0	1	0		計	9	(0.0%)	0	0	3	4	0	1	1
狭窄	男	1	(0.0%)	0	0	0	0	1	0	0	胃炎の疑い	男	106	(0.8%)	0	1	25	31	38	10	1
	女	1	(0.0%)	0	0	0	0	1	0	0		女	40	(0.5%)	1	4	9	8	13	5	0
	計	2	(0.0%)	0	0	0	0	2	0	0		計	146	(0.7%)	1	5	34	39	51	15	1
辺縁不整	男	2	(0.0%)	0	0	0	1	1	0	0	胃潰瘍の疑い	男	74	(0.6%)	0	0	11	14	25	19	5
	女	0	(0.0%)	0	0	0	0	0	0	0		女	22	(0.3%)	0	0	4	8	7	3	0
	計	2	(0.0%)	0	0	0	1	1	0	0		計	96	(0.5%)	0	0	15	22	32	22	5
粘膜不整	男	1	(0.0%)	0	0	0	0	0	1	0	巨大皺壁の疑い	男	7	(0.1%)	0	0	0	0	3	4	0
	女	0	(0.0%)	0	0	0	0	0	0	0		女	1	(0.0%)	0	0	0	0	1	0	0
	計	1	(0.0%)	0	0	0	0	0	1	0		計	8	(0.0%)	0	0	0	0	4	4	0
食道裂孔ヘルニア	男	9	(0.1%)	0	0	0	1	4	2	2	隆起性病変の疑い	男	211	(1.7%)	0	2	35	59	66	43	6
	女	7	(0.1%)	0	0	0	0	1	5	1		女	388	(4.8%)	0	4	57	175	110	35	7
	計	16	(0.1%)	0	0	0	1	5	7	3		計	599	(2.9%)	0	6	92	234	176	78	13
食道炎の疑い	男	6	(0.0%)	0	0	0	2	2	1	1	陥凹性病変の疑い	男	4	(0.0%)	0	0	2	1	1	0	0
	女	2	(0.0%)	0	0	0	1	0	1	0		女	0	(0.0%)	0	0	0	0	0	0	0
	計	8	(0.0%)	0	0	0	3	2	2	1		計	4	(0.0%)	0	0	2	1	1	0	0
隆起性病変の疑い	男	13	(0.1%)	0	0	1	3	3	5	1	憩室の疑い	男	25	(0.2%)	0	0	4	9	9	2	1
	女	7	(0.1%)	0	0	1	2	3	1	0		女	18	(0.2%)	0	1	3	7	6	0	1
	計	20	(0.1%)	0	0	2	5	6	6	1		計	43	(0.2%)	0	1	7	16	15	2	2
憩室の疑い	男	28	(0.2%)	0	0	0	1	12	15	0	術後胃	男	68	(0.5%)	0	0	1	8	21	34	4
	女	12	(0.1%)	0	0	0	3	3	3	3		女	8	(0.1%)	0	0	0	1	3	3	1
	計	40	(0.2%)	0	0	0	4	15	18	3		計	76	(0.4%)	0	0	1	9	24	37	5
その他の所見	男	34	(0.3%)	0	0	2	4	15	9	4	その他の所見	男	33	(0.3%)	0	0	5	7	11	7	3
	女	10	(0.1%)	0	0	0	2	4	2	2		女	23	(0.3%)	0	0	1	7	7	7	1
	計	44	(0.2%)	0	0	2	6	19	11	6		計	56	(0.3%)	0	0	6	14	18	14	4
胃部										十二指腸部											
伸展不良	男	37	(0.3%)	0	0	1	13	12	10	1	十二指腸球部変形	男	153	(1.2%)	0	2	19	35	68	26	3
	女	6	(0.1%)	0	0	0	2	1	1	2		女	40	(0.5%)	0	0	3	8	20	8	1
	計	43	(0.2%)	0	0	1	15	13	11	3		計	193	(0.9%)	0	2	22	43	88	34	4
辺縁不整	男	8	(0.1%)	0	0	1	1	2	4	0	十二指腸潰瘍の疑い	男	29	(0.2%)	0	0	6	6	11	6	0
	女	3	(0.0%)	0	1	0	0	0	2	0		女	11	(0.1%)	0	0	0	4	5	2	0
	計	11	(0.1%)	0	1	1	1	2	6	0		計	40	(0.2%)	0	0	6	10	16	8	0
変形	男	277	(2.2%)	0	0	21	80	93	72	11	十二指腸潰瘍瘢痕の疑い	男	15	(0.1%)	0	0	1	5	4	3	2
	女	28	(0.3%)	0	0	3	5	15	5	0		女	0	(0.0%)	0	0	0	0	0	0	0
	計	305	(1.5%)	0	0	24	85	108	77	11		計	15	(0.1%)	0	0	1	5	4	3	2
粘膜不整	男	2	(0.0%)	0	0	0	0	2	0	0	十二指腸炎の疑い	男	0	(0.0%)	0	0	0	0	0	0	0
	女	7	(0.1%)	0	0	1	3	2	0	1		女	1	(0.0%)	0	0	1	0	0	0	0
	計	9	(0.0%)	0	0	1	3	4	0	1		計	1	(0.0%)	0	0	1	0	0	0	0
ニッシュ	男	3	(0.0%)	0	0	0	0	2	1	0	十二指腸憩室の疑い	男	7	(0.1%)	0	0	0	2	2	3	0
	女	1	(0.0%)	0	0	0	1	0	0	0		女	1	(0.0%)	0	0	0	1	0	0	0
	計	4	(0.0%)	0	0	0	1	2	1	0		計	8	(0.0%)	0	0	0	3	2	3	0
レリーフ集中係	男	17	(0.1%)	0	0	2	1	7	7	0	造影不良	男	12	(0.1%)	0	0	4	6	2	0	0
	女	5	(0.1%)	0	0	0	3	1	0	1		女	2	(0.0%)	0	0	1	1	0	0	0
	計	22	(0.1%)	0	0	2	4	8	7	1		計	14	(0.1%)	0	0	5	7	2	0	0
レリーフ異常	男	1	(0.0%)	0	0	0	0	0	1	0	十二指腸拡張不良	男	4	(0.0%)	0	0	0	2	1	1	0
	女	0	(0.0%)	0	0	0	0	0	0	0		女	0	(0.0%)	0	0	0	0	0	0	0
	計	1	(0.0%)	0	0	0	0	0	1	0		計	4	(0.0%)	0	0	0	2	1	1	0
バリウム斑	男	2	(0.0%)	0	0	0	0	2	0	0	検診者総数	男	12,513		10	347	2,499	3,761	3,725	1,956	215
	女	0	(0.0%)	0	0	0	0	0	0	0		女	8,151		3	214	1,484	2,684	2,601	1,042	123
	計	2	(0.0%)	0	0	0	0	2	0	0		計	20,664		13	561	3,983	6,445	6,326	2,998	338