

動脈硬化検査判定結果

判定	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
	男	女							
異常なし	男	40 (17.6%)	0	0	3	12	24	1	0
	女	30 (29.1%)	0	0	4	11	12	3	0
	計	70 (21.2%)	0	0	7	23	36	4	0
放置可	男	4 (1.8%)	0	0	0	2	2	0	0
	女	6 (5.8%)	0	0	0	1	2	3	0
	計	10 (3.0%)	0	0	0	3	4	3	0
要経過観察	男	159 (70.0%)	0	0	4	18	104	29	4
	女	58 (56.3%)	0	0	1	5	27	19	6
	計	217 (65.8%)	0	0	5	23	131	48	10
要精密検査	男	22 (9.7%)	0	0	0	0	4	10	8
	女	9 (8.7%)	0	0	0	0	1	5	3
	計	31 (9.4%)	0	0	0	0	5	15	11
要治療	男	0 (0.0%)	0	0	0	0	0	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	0 (0.0%)	0	0	0	0	0	0	0
治療中	男	2 (0.9%)	0	0	0	0	0	1	1
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	2 (0.6%)	0	0	0	0	0	1	1
合計	男	227	0	0	7	32	134	41	13
	女	103	0	0	5	17	42	30	9
	計	330	0	0	12	49	176	71	22

血圧脈波検査の性年齢別平均値

検査項目	総数		29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		
	男	女	男	女	男	女	男	女	男	女	男	女	男	女	
ABI検査・右	男	227	1.136	0	0	7	1.094	32	1.143	134	1.137	41	1.155	13	1.071
	女	102	1.104	0	0	5	1.108	17	1.076	41	1.1	30	1.122	9	1.107
	計	329	1.126	0	0	12	1.1	49	1.119	175	1.129	71	1.141	22	1.085
ABI検査・左	男	227	1.111	0	0	7	1.069	32	1.118	134	1.114	41	1.124	13	1.039
	女	102	1.082	0	0	5	1.07	17	1.065	41	1.074	30	1.095	9	1.111
	計	329	1.102	0	0	12	1.069	49	1.1	175	1.104	71	1.112	22	1.069
PWV検査・右 cm/s	男	227	1397.2	0	0	7	1325	32	1351.4	134	1348.2	41	1513.1	13	1688.8
	女	102	1385.3	0	0	5	1296	17	1200.6	41	1321.5	30	1488.1	9	1731.3
	計	329	1393.5	0	0	12	1312.9	49	1299.1	175	1341.9	71	1502.6	22	1706.2
PWV検査・左 cm/s	男	226	1381.3	0	0	7	1259.1	32	1321.8	134	1338.5	41	1519.8	12	1616.3
	女	102	1384	0	0	5	1277	17	1189.6	41	1317.2	30	1498.8	9	1731.9
	計	328	1382.1	0	0	12	1266.6	49	1275.9	175	1333.5	71	1510.9	21	1665.9

頭部動脈超音波検査所見

所見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
	男	女							
プラーク個数	男	226 (99.6%)	0	0	7	32	133	41	13
	女	103 (100.0%)	0	0	5	17	42	30	9
	計	329 (99.7%)	0	0	12	49	175	71	22
不安定プラーク (III a・III b・IV)	男	4 (1.8%)	0	0	0	0	2	1	1
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	4 (1.2%)	0	0	0	0	2	1	1
頭動脈の内腔狭窄 (50-75%)	男	2 (0.9%)	0	0	0	0	0	0	2
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	2 (0.6%)	0	0	0	0	0	0	2
頭動脈の高度狭窄 (75%以上)	男	1 (0.4%)	0	0	0	0	0	0	1
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	1 (0.3%)	0	0	0	0	0	0	1
頭動脈蛇行	男	1 (0.4%)	0	0	0	0	0	0	1
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	1 (%)	0	0	0	0	0	0	1
頭動脈描出不良	男	15 (6.6%)	0	0	1	5	2	5	2
	女	3 (2.9%)	0	0	0	0	2	1	0
	計	18 (5.5%)	0	0	1	5	4	6	2
不安定プラーク (III a)	男	0 (0.0%)	0	0	0	0	0	0	0
	女	3 (2.9%)	0	0	0	0	0	2	1
	計	3 (0.9%)	0	0	0	0	0	2	1
不安定プラーク (III b)	男	1 (0.4%)	0	0	0	0	0	1	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	1 (0.3%)	0	0	0	0	0	1	0
総頭動脈壁肥厚	男	7 (3.1%)	0	0	0	0	2	4	1
	女	5 (4.9%)	0	0	0	0	1	2	2
	計	12 (3.6%)	0	0	0	0	3	6	3
その他の所見	男	8 (3.5%)	0	0	0	0	3	5	0
	女	3 (2.9%)	0	0	0	0	1	2	0
	計	11 (3.3%)	0	0	0	0	4	7	0
検診者総数	男	227	0	0	7	32	134	41	13
	女	103	0	0	5	17	42	30	9
	計	330	0	0	12	49	176	71	22

頭部動脈超音波検査の計測値の性年齢別平均値

検査項目	総数		29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		
	男	女	男	女	男	女	男	女	男	女	男	女	男	女	
平均IMT右 mm	男	227	0.707	0	0	7	0.58	32	0.632	134	0.683	41	0.82	13	0.855
	女	103	0.724	0	0	5	0.556	17	0.62	42	0.716	30	0.781	9	0.857
	計	330	0.712	0	0	12	0.57	49	0.628	176	0.691	71	0.804	22	0.856
平均IMT左 mm	男	227	0.764	0	0	7	0.59	32	0.69	134	0.724	41	0.908	13	0.992
	女	103	0.755	0	0	5	0.632	17	0.619	42	0.717	30	0.838	9	0.977
	計	330	0.761	0	0	12	0.608	49	0.665	176	0.722	71	0.878	22	0.986
最大IMT右 mm	男	227	1.298	0	0	7	0.856	32	0.999	134	1.263	41	1.561	13	1.805
	女	103	1.17	0	0	5	0.932	17	1.052	42	1.161	30	1.257	9	1.272
	計	330	1.258	0	0	12	0.888	49	1.018	176	1.239	71	1.433	22	1.587
最大IMT左 mm	男	227	1.295	0	0	7	0.853	32	1.027	134	1.254	41	1.601	13	1.655
	女	103	1.219	0	0	5	0.918	17	0.985	42	1.104	30	1.409	9	1.727
	計	330	1.271	0	0	12	0.88	49	1.012	176	1.218	71	1.52	22	1.685