

前立腺検査の判定結果

| 判定 | 総数 | | 19歳以下 | 20-29歳 | 30-39歳 | 40-49歳 | 50-59歳 | 60-69歳 | 70歳以上 |
|-------|-------|---------|-------|--------|--------|--------|--------|--------|-------|
| 異常なし | 4,102 | (95.3%) | 0 | 5 | 75 | 324 | 2,658 | 904 | 136 |
| 放置可 | 0 | (0.0%) | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 要経過観察 | 5 | (0.1%) | 0 | 0 | 0 | 0 | 2 | 2 | 1 |
| 要精密検査 | 111 | (2.6%) | 0 | 0 | 3 | 1 | 50 | 52 | 5 |
| 要治療 | 0 | (0.0%) | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 治療中 | 87 | (2.0%) | 0 | 0 | 2 | 2 | 31 | 34 | 18 |
| 合計 | 4,305 | 0 | 0 | 5 | 80 | 327 | 2,741 | 992 | 160 |

P S A の年齢別平均値

| 検査項目 | 総数 | | 29歳以下 | | 30-39歳 | | 40-49歳 | | 50-59歳 | | 60-69歳 | | 70歳以上 | |
|--------------|-------|------|-------|------|--------|------|--------|------|--------|------|--------|------|-------|------|
| PSA ng/mL | 4,306 | 1.29 | 5 | 0.78 | 80 | 1.23 | 327 | 0.91 | 2,741 | 1.16 | 992 | 1.66 | 161 | 1.98 |

骨密度検査判定結果

| 判定 | 総数 | | 19歳以下 | 20-29歳 | 30-39歳 | 40-49歳 | 50-59歳 | 60-69歳 | 70歳以上 | |
|-------|----|-------|---------|--------|--------|--------|--------|--------|-------|----|
| 異常なし | 男 | 140 | (65.4%) | 0 | 0 | 12 | 20 | 58 | 38 | 12 |
| | 女 | 830 | (58.0%) | 91 | 17 | 125 | 254 | 280 | 57 | 6 |
| | 計 | 970 | (59.0%) | 91 | 17 | 137 | 274 | 338 | 95 | 18 |
| 放置可 | 男 | 2 | (0.9%) | 0 | 0 | 2 | 0 | 0 | 0 | 0 |
| | 女 | 54 | (3.8%) | 0 | 4 | 27 | 22 | 1 | 0 | 0 |
| | 計 | 56 | (3.4%) | 0 | 4 | 29 | 22 | 1 | 0 | 0 |
| 要経過観察 | 男 | 54 | (25.2%) | 0 | 0 | 0 | 3 | 18 | 18 | 15 |
| | 女 | 428 | (29.9%) | 0 | 5 | 33 | 63 | 188 | 116 | 23 |
| | 計 | 482 | (29.3%) | 0 | 5 | 33 | 66 | 206 | 134 | 38 |
| 要精密検査 | 男 | 13 | (6.1%) | 0 | 0 | 0 | 0 | 8 | 2 | 3 |
| | 女 | 80 | (5.6%) | 0 | 0 | 0 | 5 | 31 | 33 | 11 |
| | 計 | 93 | (5.7%) | 0 | 0 | 0 | 5 | 39 | 35 | 14 |
| 要治療 | 男 | 0 | (0.0%) | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | (0.0%) | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 0 | (0.0%) | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 治療中 | 男 | 5 | (2.3%) | 0 | 0 | 0 | 0 | 1 | 2 | 2 |
| | 女 | 39 | (2.7%) | 0 | 0 | 0 | 3 | 7 | 20 | 9 |
| | 計 | 44 | (2.7%) | 0 | 0 | 0 | 3 | 8 | 22 | 11 |
| 合計 | 男 | 214 | | 0 | 0 | 14 | 23 | 85 | 60 | 32 |
| | 女 | 1,431 | | 91 | 26 | 185 | 347 | 507 | 226 | 49 |
| | 計 | 1,645 | | 91 | 26 | 199 | 370 | 592 | 286 | 81 |

超音波骨密度検査の性年齢別平均値

| 検査項目 | 総数 | | 29歳以下 | | 30-39歳 | | 40-49歳 | | 50-59歳 | | 60-69歳 | | 70歳以上 | | |
|---------------|----|-------|-------|-----|--------|-----|--------|-----|--------|-----|--------|-----|-------|----|------|
| 骨密度 スティフネス | 男 | 214 | 98.7 | 0 | — | 14 | 106.6 | 23 | 111.8 | 85 | 99.0 | 60 | 97.8 | 32 | 87.2 |
| | 女 | 1,431 | 91.0 | 117 | 121.3 | 185 | 96.7 | 347 | 96.8 | 507 | 86.4 | 226 | 76.3 | 49 | 72.5 |
| | 計 | 1,645 | 92.0 | 117 | 121.3 | 199 | 97.4 | 370 | 97.8 | 592 | 88.2 | 286 | 80.8 | 81 | 78.3 |