

動脈硬化検査判定結果

判定	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
	男	女							
異常なし	男	118 (18.1%)	0	2	10	54	49	3	0
	女	148 (29.7%)	0	0	12	70	60	5	1
	計	266 (23.1%)	0	2	22	124	109	8	1
放置可	男	46 (7.1%)	0	0	2	13	29	2	0
	女	32 (6.4%)	0	0	0	4	18	9	1
	計	78 (6.8%)	0	0	2	17	47	11	1
要経過観察	男	419 (64.3%)	0	0	15	83	185	111	25
	女	291 (58.3%)	0	0	1	38	122	107	23
	計	710 (61.7%)	0	0	16	121	307	218	48
要精密検査	男	63 (9.7%)	0	0	0	5	24	21	13
	女	28 (5.6%)	0	0	1	0	3	15	9
	計	91 (7.9%)	0	0	1	5	27	36	22
要治療	男	1 (0.2%)	0	0	0	0	1	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	1 (0.1%)	0	0	0	0	1	0	0
治療中	男	5 (0.8%)	0	0	0	0	0	2	3
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	5 (0.4%)	0	0	0	0	0	2	3
合計	男	652	0	2	27	155	288	139	41
	女	499	0	0	14	112	203	136	34
	計	1,151	0	2	41	267	491	275	75

頸部動脈超音波検査所見

所見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
	男	女							
頸動脈の内腔狭窄 (50-75%)	男	12 (1.8%)	0	0	0	1	5	3	3
	女	4 (0.8%)	0	0	0	1	1	0	2
	計	16 (1.4%)	0	0	0	2	6	3	5
頸動脈の高度狭窄 (75%以上)	男	1 (0.2%)	0	0	0	0	0	0	1
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	1 (0.1%)	0	0	0	0	0	0	1
総頸動脈内径拡大	男	1 (0.2%)	0	0	0	0	0	1	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	1 (0.1%)	0	0	0	0	0	1	0
頸動脈描出不良	男	29 (4.4%)	0	0	3	4	11	7	4
	女	16 (3.2%)	0	0	0	3	7	4	2
	計	45 (3.9%)	0	0	3	7	18	11	6
不安定プラーク (Ⅲa)	男	2 (0.3%)	0	0	0	0	1	1	0
	女	1 (0.2%)	0	0	1	0	0	0	0
	計	3 (0.3%)	0	0	1	0	1	1	0
不安定プラーク (Ⅲb)	男	4 (0.6%)	0	0	0	1	3	0	0
	女	6 (1.2%)	0	0	0	1	1	3	1
	計	10 (0.9%)	0	0	0	2	4	3	1
不安定プラーク (Ⅳ)	男	1 (0.2%)	0	0	0	0	1	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	1 (0.1%)	0	0	0	0	1	0	0
総頸動脈壁肥厚	男	48 (7.4%)	0	0	0	4	12	20	12
	女	20 (4.0%)	0	0	0	0	2	10	8
	計	68 (5.9%)	0	0	0	4	14	30	20
その他の所見	男	5 (0.8%)	0	0	0	1	3	1	0
	女	1 (0.2%)	0	0	0	0	0	0	1
	計	6 (0.5%)	0	0	0	1	3	1	1
検診者総数	男	652	0	2	27	155	288	139	41
	女	500	0	0	14	113	203	136	34
	計	1,152	0	2	41	268	491	275	75

頸部動脈超音波検査の計測値の性年齢別平均値

検査項目	総数		29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		
	男	女	男	女	男	女	男	女	男	女	男	女	男	女	
平均IMT 右mm	男	652	0.720	2	0.620	27	0.560	155	0.630	288	0.720	139	0.810	41	0.870
	女	500	0.690	0	0.000	14	0.510	113	0.580	203	0.670	136	0.780	34	0.930
	計	1,152	0.710	2	0.620	41	0.540	268	0.610	491	0.700	275	0.800	75	0.900
平均IMT 左mm	男	652	0.780	2	0.600	27	0.600	155	0.690	288	0.770	139	0.890	41	1.020
	女	500	0.720	0	0.000	14	0.510	113	0.620	203	0.700	136	0.790	34	0.950
	計	1,152	0.760	2	0.600	41	0.570	268	0.660	491	0.740	275	0.840	75	0.990
最大IMT 右mm	男	652	1.340	2	0.820	27	0.920	155	1.130	288	1.350	139	1.510	41	1.770
	女	500	1.210	0	0.000	14	0.890	113	0.990	203	1.200	136	1.330	34	1.600
	計	1,152	1.280	2	0.820	41	0.910	268	1.070	491	1.290	275	1.420	75	1.690
最大IMT 左mm	男	652	1.380	2	0.770	27	0.970	155	1.140	288	1.360	139	1.650	41	1.770
	女	500	1.220	0	0.000	14	0.810	113	1.020	203	1.210	136	1.380	34	1.490
	計	1,152	1.310	2	0.770	41	0.910	268	1.090	491	1.300	275	1.520	75	1.650

血圧脈波検査の性年齢別平均値

検査項目	総数		29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		
	男	女	男	女	男	女	男	女	男	女	男	女	男	女	
ABI検査 右	男	652	1.150	2	1.100	27	1.120	155	1.150	288	1.160	139	1.160	41	1.150
	女	496	1.130	0	0.000	14	1.110	112	1.120	201	1.130	136	1.140	33	1.140
	計	1,148	1.140	2	1.100	41	1.120	267	1.140	489	1.150	275	1.150	74	1.150
ABI検査 左	男	652	1.130	2	1.110	27	1.100	155	1.130	288	1.130	139	1.150	41	1.120
	女	496	1.100	0	0.000	14	1.080	112	1.100	201	1.100	136	1.120	33	1.120
	計	1,148	1.120	2	1.110	41	1.090	267	1.120	489	1.120	275	1.130	74	1.120
PWV検査 右 cm/s	男	651	1424.4	2	1222.5	27	1338.3	154	1310.2	288	1391.9	139	1537.6	41	1765.0
	女	496	1367.4	0	0.0	14	1151.0	112	1190.2	201	1320.1	136	1516.5	33	1734.0
	計	1,147	1399.8	2	1222.5	41	1274.3	266	1259.7	489	1362.4	275	1527.2	74	1751.2
PWV検査 左 cm/s	男	651	1407.6	2	1238.0	27	1322.4	154	1291.1	288	1374.4	139	1545.3	41	1676.4
	女	496	1368.5	0	0.0	14	1133.4	112	1189.0	201	1319.2	136	1524.0	33	1736.5
	計	1,147	1390.7	2	1238.0	41	1257.9	266	1248.1	489	1351.7	275	1534.8	74	1703.2