

動脈硬化検査判定結果

判定	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
	男	女							
異常なし	男	109 (16.9%)	0	0	11	57	35	6	0
	女	130 (25.9%)	0	0	11	64	45	10	0
	計	239 (20.9%)	0	0	22	121	80	16	0
放置可	男	61 (9.5%)	0	0	3	25	27	6	0
	女	59 (11.8%)	0	0	2	13	35	8	1
	計	120 (10.5%)	0	0	5	38	62	14	1
要経過観察	男	400 (62.1%)	0	0	4	85	172	105	34
	女	280 (55.8%)	0	0	2	33	124	106	15
	計	680 (59.3%)	0	0	6	118	296	211	49
要精密検査	男	66 (10.2%)	0	0	0	5	17	38	6
	女	33 (6.6%)	0	0	0	2	4	16	11
	計	99 (8.6%)	0	0	0	7	21	54	17
要治療	男	0 (0.0%)	0	0	0	0	0	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	0 (0.0%)	0	0	0	0	0	0	0
治療中	男	8 (1.2%)	0	0	0	0	1	5	2
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	8 (0.7%)	0	0	0	0	1	5	2
合計	男	644	0	0	18	172	252	160	42
	女	502	0	0	15	112	208	140	27
	計	1,146	0	0	33	284	460	300	69

頸部動脈超音波検査所見

所見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
	男	女							
頸動脈の内腔狭窄 (50-75%)	男	17 (2.6%)	0	0	0	0	2	12	3
	女	3 (0.6%)	0	0	0	0	1	0	2
	計	20 (1.7%)	0	0	0	0	3	12	5
頸動脈の高度狭窄 (75%以上)	男	2 (0.3%)	0	0	0	0	0	1	1
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	2 (0.2%)	0	0	0	0	0	1	1
総頸動脈内径拡大	男	4 (0.6%)	0	0	0	0	1	3	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	4 (0.3%)	0	0	0	0	1	3	0
頸動脈描出不良	男	33 (5.1%)	0	0	1	9	10	9	4
	女	16 (3.2%)	0	0	0	3	3	8	2
	計	49 (4.3%)	0	0	1	12	13	17	6
不安定プラーク (III b)	男	3 (0.5%)	0	0	0	0	0	3	0
	女	3 (0.6%)	0	0	0	1	0	1	1
	計	6 (0.5%)	0	0	0	1	0	4	1
不安定プラーク (IV)	男	1 (0.2%)	0	0	0	0	0	1	0
	女	1 (0.2%)	0	0	0	0	0	1	0
	計	2 (0.2%)	0	0	0	0	0	2	0
総頸動脈壁肥厚	男	59 (9.2%)	0	0	0	4	14	29	12
	女	19 (3.8%)	0	0	0	0	3	11	5
	計	78 (6.8%)	0	0	0	4	17	40	17
その他の所見	男	4 (0.6%)	0	0	0	1	1	2	0
	女	1 (0.2%)	0	0	0	0	0	1	0
	計	5 (0.4%)	0	0	0	1	1	3	0
検診者総数	男	644	0	0	18	172	252	160	42
	女	502	0	0	15	112	208	140	27
	計	1,146	0	0	33	284	460	300	69

頸部動脈超音波検査の計測値の性年齢別平均値

検査項目	総数		29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		
	男	女	男	女	男	女	男	女	男	女	男	女	男	女	
平均IMT 右mm	男	644	0.740	0	—	18	0.590	172	0.620	252	0.720	160	0.850	42	0.940
	女	502	0.690	0	—	15	0.510	112	0.590	208	0.670	140	0.760	27	0.890
	計	1,146	0.720	0	—	33	0.560	284	0.610	460	0.700	300	0.810	69	0.920
平均IMT 左mm	男	644	0.790	0	—	18	0.620	172	0.670	252	0.790	160	0.880	42	1.050
	女	502	0.720	0	—	15	0.530	112	0.610	208	0.710	140	0.790	27	0.950
	計	1,146	0.760	0	—	33	0.580	284	0.640	460	0.750	300	0.840	69	1.010
最大IMT 右mm	男	644	1.360	0	—	18	0.960	172	1.130	252	1.300	160	1.630	42	1.830
	女	502	1.230	0	—	15	1.010	112	1.060	208	1.230	140	1.330	27	1.470
	計	1,146	1.300	0	—	33	0.980	284	1.100	460	1.270	300	1.490	69	1.690
最大IMT 左mm	男	644	1.400	0	—	18	0.980	172	1.140	252	1.360	160	1.690	42	1.770
	女	502	1.250	0	—	15	0.910	112	1.050	208	1.260	140	1.370	27	1.530
	計	1,146	1.330	0	—	33	0.950	284	1.100	460	1.310	300	1.540	69	1.670

血圧脈波検査の性年齢別平均値

検査項目	総数		29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		
	男	女	男	女	男	女	男	女	男	女	男	女	男	女	
ABI検査 右	男	641	1.150	0	—	18	1.130	172	1.140	251	1.150	159	1.160	41	1.150
	女	498	1.120	0	—	15	1.090	111	1.120	207	1.120	139	1.130	26	1.130
	計	1,139	1.140	0	—	33	1.120	283	1.130	458	1.140	298	1.150	67	1.140
ABI検査 左	男	641	1.140	0	—	18	1.110	172	1.120	251	1.140	159	1.150	41	1.140
	女	498	1.110	0	—	15	1.090	111	1.110	207	1.110	139	1.120	26	1.120
	計	1,139	1.130	0	—	33	1.100	283	1.120	458	1.130	298	1.140	67	1.130
PWV検査 右 cm/s	男	641	1434.6	0	—	18	1213.3	172	1313.2	251	1390.1	159	1583.1	41	1737.6
	女	498	1348.9	0	—	15	1139.1	111	1156.3	207	1303.8	139	1516.3	26	1756.7
	計	1,139	1397.1	0	—	33	1179.6	283	1251.7	458	1351.1	298	1551.9	67	1745.0
PWV検査 左 cm/s	男	641	1425.0	0	—	18	1199.4	172	1303.2	251	1385.2	159	1577.7	41	1686.0
	女	498	1360.0	0	—	15	1134.3	111	1160.4	207	1317.2	139	1531.6	26	1764.9
	計	1,139	1396.6	0	—	33	1169.9	283	1247.2	458	1354.5	298	1556.2	67	1716.6