

総合健診結果（コース別健診、市民健診）

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70-79歳	80歳以上
異常なし	男	3,130	(7.2%)	102	817	870	479	400	200	99	7
	女	5,447	(12.0%)	164	973	1,281	1,180	621	640	243	17
	計	8,577	(9.7%)	236	1,790	2,151	1,659	1,021	840	342	24
放置可	男	3,570	(8.3%)	79	1,036	1,034	705	395	285	227	36
	女	6,479	(14.3%)	111	1,574	1,322	1,519	983	945	423	44
	計	10,049	(11.3%)	190	2,610	2,356	2,224	1,378	1,230	650	80
要経過観察	男	16,535	(38.2%)	181	1,775	3,172	3,886	3,122	2,458	1,649	394
	女	16,232	(37.2%)	114	1,134	2,207	3,393	3,305	3,701	2,068	437
	計	32,767	(35.7%)	295	2,909	5,379	7,279	6,427	6,159	3,717	831
要精密検査	男	6,983	(16.1%)	16	239	1,365	1,895	1,521	983	463	129
	女	6,006	(13.2%)	8	224	926	1,505	1,287	1,071	506	157
	計	12,989	(14.6%)	24	463	2,291	3,400	2,808	2,054	969	286
要治療	男	1,270	(2.9%)	1	34	159	322	298	282	128	40
	女	871	(1.9%)	1	7	46	128	208	292	159	27
	計	2,141	(2.4%)	2	41	205	450	506	574	287	67
治療中	男	11,780	(27.2%)	2	34	338	1,340	2,909	3,467	2,849	837
	女	10,381	(22.9%)	1	30	119	572	1,661	3,579	3,327	1,086
	計	22,161	(25.0%)	3	64	457	1,912	4,570	7,046	6,176	1,923
合計	男	43,268		351	3,935	6,938	8,627	8,645	7,675	5,415	1,443
	女	45,416		399	3,942	5,901	8,297	8,065	10,228	6,726	1,768
	計	88,684		750	7,877	12,839	16,924	16,710	17,903	12,141	3,211

胸部検査判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	22,804	(90.1%)	217	2,749	4,867	6,073	5,574	2,999	325
	女	19,349	(93.6%)	177	3,062	4,069	5,217	4,663	1,956	205
	計	42,153	(91.7%)	394	5,811	8,936	11,290	10,237	4,955	530
放置可	男	1,750	(6.9%)	4	45	140	381	622	488	70
	女	913	(4.4%)	3	49	67	210	320	221	43
	計	2,663	(5.8%)	7	94	207	591	942	709	113
要経過観察	男	340	(1.3%)	0	4	21	66	119	112	18
	女	117	(0.6%)	0	2	10	21	35	40	9
	計	457	(1.0%)	0	6	31	87	154	152	27
要精密検査	男	401	(1.6%)	0	12	48	82	131	105	23
	女	274	(1.3%)	0	8	30	64	101	60	11
	計	675	(1.5%)	0	20	78	146	232	165	34
要治療	男	1	(0.0%)	0	0	0	0	0	1	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	1	(0.0%)	0	0	0	0	0	1	0
治療中	男	22	(0.1%)	0	0	3	0	7	9	3
	女	13	(0.1%)	0	1	1	6	1	3	1
	計	35	(0.1%)	0	1	4	6	8	12	4
合 計	男	25,318		221	2,810	5,079	6,602	6,453	3,714	439
	女	20,666		180	3,122	4,177	5,518	5,120	2,280	269
	計	45,984		401	5,932	9,256	12,120	11,573	5,994	708

喀痰細胞診所見

所 見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
B 1 (Class I (異常なし))	男	12 (21.1%)	0	0	2	3	4	3	0
	女	3 (21.4%)	0	0	0	2	1	0	0
	計	15 (21.1%)	0	0	2	5	5	3	0
B 2 (Class II (悪性所見なし))	男	45 (78.9%)	2	4	3	10	16	9	1
	女	11 (78.6%)	1	0	0	5	3	2	0
	計	56 (78.9%)	3	4	3	15	19	11	1
検診者総数	男	57	2	4	5	13	20	12	1
	女	14	1	0	0	7	4	2	0
	計	71	3	4	5	20	24	14	1

胸部ヘリカルＣＴ検査所見

所 見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
線状陰影	男	37 (3.1%)	0	0	0	6	16	12	3
	女	10 (4.1%)	0	0	0	1	2	4	3
	計	47 (3.3%)	0	0	0	7	18	16	6
帯状陰影	男	21 (1.8%)	0	0	1	4	11	5	0
	女	7 (2.9%)	0	0	0	0	3	3	1
	計	28 (1.9%)	0	0	1	4	14	8	1
微細斑点状陰影	男	1 (0.1%)	0	0	0	0	0	1	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	1 (0.1%)	0	0	0	0	0	1	0
結節陰影	男	193 (16.1%)	0	0	9	40	94	42	8
	女	54 (22.0%)	0	1	2	17	15	14	5
	計	247 (17.1%)	0	1	11	57	109	56	13
浸潤陰影	男	23 (1.9%)	0	0	1	3	14	4	1
	女	8 (3.3%)	0	0	0	0	3	4	1
	計	31 (2.1%)	0	0	1	3	17	8	2
網目状陰影	男	4 (0.3%)	0	0	0	1	1	2	0
	女	3 (1.2%)	0	0	0	0	1	0	2
	計	7 (0.5%)	0	0	0	1	2	2	2
石灰化陰影	男	10 (0.8%)	0	0	2	2	2	2	2
	女	6 (2.4%)	0	0	0	0	2	2	2
	計	16 (1.1%)	0	0	2	2	4	4	4
囊胞陰影	男	156 (13.0%)	0	0	7	44	74	27	4
	女	11 (4.5%)	0	0	0	3	4	3	1
	計	167 (11.6%)	0	0	7	47	78	30	5
気腫性変化	男	39 (3.3%)	0	0	1	8	22	8	0
	女	1 (0.4%)	0	0	0	0	1	0	0
	計	40 (2.8%)	0	0	1	8	23	8	0
すりガラス状陰影	男	38 (3.2%)	0	0	1	9	16	10	2
	女	8 (3.3%)	0	0	0	1	5	1	1
	計	46 (3.2%)	0	0	1	10	21	11	3
粒状陰影	男	3 (0.3%)	0	0	0	1	2	0	0
	女	3 (1.2%)	0	0	1	0	0	1	1
	計	6 (0.4%)	0	0	1	1	2	1	1
胸膜肥厚	男	31 (2.6%)	0	0	1	6	14	10	0
	女	12 (4.9%)	0	0	0	3	5	2	2
	計	43 (3.0%)	0	0	1	9	19	12	2
胸膜癒着	男	11 (0.9%)	0	0	0	1	3	7	0
	女	4 (1.6%)	0	0	0	0	2	2	0
	計	15 (1.0%)	0	0	0	1	5	9	0
肺気腫	男	39 (3.3%)	0	0	0	2	24	9	4
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	39 (2.7%)	0	0	0	2	24	9	4
縦隔腫瘍の疑い	男	0 (0.0%)	0	0	0	0	0	0	0
	女	1 (0.4%)	0	0	1	0	0	0	0
	計	1 (0.1%)	0	0	1	0	0	0	0
その他の異常	男	1 (0.1%)	0	0	0	1	0	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	1 (0.1%)	0	0	0	1	0	0	0
その他の所見	男	122 (10.2%)	0	0	7	21	55	36	3
	女	19 (7.8%)	0	0	0	5	8	5	1
	計	141 (9.8%)	0	0	7	26	63	41	4
検診者総数	男	1,197	0	3	83	310	556	212	33
	女	245	0	2	19	63	88	57	16
	計	1,442	0	5	102	373	644	269	49

胸部×線単純検査所見

所 見		総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上	所 見		総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
心陰影拡大	男	58	(0.2%)	0	1	10	9	16	16	6	気腫様陰影	男	1	(0.0%)	0	0	0	0	1	0	0
	女	18	(0.1%)	0	1	0	2	4	11	0		女	0	(0.0%)	0	0	0	0	0	0	0
	計	76	(0.2%)	0	2	10	11	20	27	6		計	1	(0.0%)	0	0	0	0	1	0	0
肺門陰影拡大	男	17	(0.1%)	0	0	1	4	3	8	1	胸膜肥厚	男	326	(1.3%)	0	2	24	73	108	110	9
	女	14	(0.1%)	0	0	2	2	5	3	2		女	242	(1.2%)	0	2	9	53	105	68	5
	計	31	(0.1%)	0	0	3	6	8	11	3		計	568	(1.2%)	0	4	33	126	213	178	14
右胸心	男	2	(0.0%)	0	0	0	2	0	0	0	胸膜癒着	男	214	(0.8%)	0	5	15	44	61	76	13
	女	2	(0.0%)	0	0	0	2	0	0	0		女	121	(0.6%)	0	1	7	27	45	29	12
	計	4	(0.0%)	0	0	0	4	0	0	0		計	335	(0.7%)	0	6	22	71	106	105	25
肺紋理増強	男	1	(0.0%)	0	0	0	0	1	0	0	胸水陰影	男	2	(0.0%)	0	0	0	0	2	0	0
	女	1	(0.0%)	0	0	0	1	0	0	0		女	0	(0.0%)	0	0	0	0	0	0	0
	計	2	(0.0%)	0	0	0	1	1	0	0		計	2	(0.0%)	0	0	0	0	2	0	0
右大動脈弓	男	20	(0.1%)	0	0	4	5	8	3	0	上葉縮小	男	5	(0.0%)	0	0	0	2	3	0	0
	女	13	(0.1%)	0	4	0	2	3	4	0		女	5	(0.0%)	0	0	1	1	1	2	0
	計	33	(0.1%)	0	4	4	7	11	7	0		計	10	(0.0%)	0	0	1	3	4	2	0
大動脈拡張・縮小	男	11	(0.0%)	0	0	0	0	3	7	1	横隔膜高位	男	72	(0.3%)	0	3	9	18	29	11	2
	女	1	(0.0%)	0	0	0	0	0	0	1		女	18	(0.1%)	0	1	2	1	6	6	2
	計	12	(0.0%)	0	0	0	0	3	7	2		計	90	(0.2%)	0	4	11	19	35	17	4
線状陰影	男	134	(0.5%)	0	4	6	30	50	38	6	横隔膜低位	男	1	(0.0%)	0	0	0	1	0	0	0
	女	66	(0.3%)	0	2	6	16	21	15	6		女	0	(0.0%)	0	0	0	0	0	0	0
	計	200	(0.4%)	0	6	12	46	71	53	12		計	1	(0.0%)	0	0	0	1	0	0	0
帯状陰影	男	64	(0.3%)	0	0	7	19	16	22	0	肋骨異常	男	25	(0.1%)	0	0	4	6	8	6	1
	女	28	(0.1%)	1	0	2	6	9	9	1		女	8	(0.0%)	0	3	1	2	2	0	0
	計	92	(0.2%)	1	0	9	25	25	31	1		計	33	(0.1%)	0	3	5	8	10	6	1
石灰沈着	男	1	(0.0%)	0	0	0	0	0	1	0	胸椎異常	男	97	(0.4%)	2	17	6	29	25	16	2
	女	1	(0.0%)	0	0	0	0	0	1	0		女	167	(0.8%)	2	29	32	51	33	17	3
	計	2	(0.0%)	0	0	0	0	0	2	0		計	264	(0.6%)	4	46	38	80	58	33	5
索状陰影	男	5	(0.0%)	0	0	0	1	1	2	1	鎖骨異常	男	1	(0.0%)	0	0	1	0	0	0	0
	女	4	(0.0%)	0	0	0	0	4	0	0		女	1	(0.0%)	0	0	0	0	0	1	0
	計	9	(0.0%)	0	0	0	1	5	2	1		計	2	(0.0%)	0	0	1	0	0	1	0
大動脈蛇行	男	26	(0.1%)	0	0	0	2	11	10	3	漏斗胸	男	3	(0.0%)	0	1	0	0	2	0	0
	女	6	(0.0%)	0	0	0	0	1	2	3		女	10	(0.0%)	0	1	4	3	2	0	0
	計	32	(0.1%)	0	0	0	2	12	12	6		計	13	(0.0%)	0	2	4	3	4	0	0
微細斑点状陰影	男	15	(0.1%)	0	0	1	1	3	9	1	手術後	男	90	(0.4%)	2	8	18	16	24	20	2
	女	7	(0.0%)	0	0	0	3	1	3	0		女	25	(0.1%)	0	3	1	5	8	8	0
	計	22	(0.0%)	0	0	1	4	4	12	1		計	115	(0.2%)	2	11	19	21	32	28	2
粒状陰影	男	3	(0.0%)	0	0	1	0	0	1	1	胸郭形成	男	0	(0.0%)	0	0	0	0	0	0	0
	女	1	(0.0%)	0	0	1	0	0	0	0		女	1	(0.0%)	0	0	0	0	1	0	0
	計	4	(0.0%)	0	0	2	0	0	1	1		計	1	(0.0%)	0	0	0	0	1	0	0
輪状陰影	男	0	(0.0%)	0	0	0	0	0	0	0	陈旧性肺結核の疑い	男	2	(0.0%)	0	0	0	0	0	0	2
	女	1	(0.0%)	0	0	0	0	0	0	1		女	1	(0.0%)	0	0	0	1	0	0	0
	計	1	(0.0%)	0	0	0	0	0	0	1		計	3	(0.0%)	0	0	0	1	0	0	2
結節陰影	男	329	(1.3%)	0	9	35	61	115	96	13	肺気腫の疑い	男	34	(0.1%)	0	0	1	7	11	12	3
	女	283	(1.4%)	0	8	26	73	106	60	10		女	4	(0.0%)	0	0	0	1	0	2	1
	計	612	(1.3%)	0	17	61	134	221	156	23		計	38	(0.1%)	0	0	1	8	11	14	4
円形陰影	男	4	(0.0%)	0	0	1	0	1	2	0	慢性気管支炎の疑い	男	13	(0.1%)	0	0	1	3	3	6	0
	女	1	(0.0%)	0	0	0	0	0	0	1		女	4	(0.0%)	0	0	0	0	2	2	0
	計	5	(0.0%)	0	0	1	0	1	2	1		計	17	(0.0%)	0	0	1	3	5	8	0
塊状陰影	男	25	(0.1%)	0	2	3	8	8	3	1	気管支拡張症の疑い	男	9	(0.0%)	0	0	2	2	0	4	1
	女	20	(0.1%)	0	0	3	4	9	4	0		女	3	(0.0%)	0	0	0	1	0	1	1
	計	45	(0.1%)	0	2	6	12	17	7	1		計	12	(0.0%)	0	0	2	3	0	5	2
浸潤陰影	男	111	(0.4%)	0	4	13	20	28	39	7	間質性肺炎の疑い	男	39	(0.2%)	0	0	0	2	9	21	7
	女	79	(0.4%)	0	3	4	13	26	27	6		女	7	(0.0%)	0	0	0	0	3	3	1
	計	190	(0.4%)	0	7	17	33	54	66	13		計	46	(0.1%)	0	0	0	2	12	24	8
網目状陰影	男	1	(0.0%)	0	0	0	0	0	1	0	塵肺の疑い	男	1	(0.0%)	0	0	0	0	0	1	0
	女	2	(0.0%)	0	0	0	0	1	0	1		女	0	(0.0%)	0	0	0	0	0	0	0
	計	3	(0.0%)	0	0	0	0	1	1	1		計	1	(0.0%)	0	0	0	0	0	1	0
石灰化陰影	男	122	(0.5%)	0	0	6	11	47	46	12	気胸の疑い	男	2	(0.0%)	0	1	0	1	0	0	0
	女	70	(0.3%)	0	1	4	13	25	21	6		女	0	(0.0%)	0	0	0	0	0	0	0
	計	192	(0.4%)	0	1	10	24	72	67	18		計	2	(0.0%)	0	1	0	1	0	0	0
空洞陰影	男	1	(0.0%)	0	0	1	0	0	0	0	その他の所見(心所見)	男	3	(0.0%)	0	0	0	1	1	1	0
	女	0	(0.0%)	0	0	0	0	0	0	0		女	0	(0.0%)	0	0	0	0	0	0	0
	計	1	(0.0%)	0	0	1	0	0	0	0		計	3	(0.0%)	0	0	0	1	1	1	0
嚢胞陰影	男	149	(0.6%)	0	4	17	39	56	30	3	その他の所見(肺所見)	男	134	(0.5%)	0	5	8	19	37	55	10
	女	8	(0.0%)	0	0	0	2	5	1	0		女	42	(0.2%)	0	3	3	11	11	12	2
	計	157	(0.3%)	0	4	17	41	61	31	3		計	176	(0.4%)	0	8	11	30	48	67	12
無気肺陰影	男	8	(0.0%)	0	0	0	2	4	2	0	検診者総数	男	25,339		221	2,820	5,086	6,605	6,453	3,715	439
	女	3	(0.0%)	0	0	0	0	2	1	0		女	20,664		180	3,122	4,177	5,516	5,120	2,280	269
	計	11	(0.0%)	0	0	0	2	6	3	0		計	46,003		401	5,942	9,263	12,121	11,573	5,995	708

上部消化管検査判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	15,653	(86.1%)	17	685	3,046	4,969	4,521	2,110	305
	女	8,445	(82.2%)	4	240	1,501	2,789	2,514	1,196	201
	計	24,098	(84.7%)	21	925	4,547	7,758	7,035	3,306	506
放置可	男	911	(5.0%)	0	3	95	259	357	165	32
	女	522	(5.1%)	0	0	67	188	197	56	14
	計	1,433	(5.0%)	0	3	162	447	554	221	46
要経過観察	男	936	(5.2%)	0	1	98	266	363	181	27
	女	894	(8.7%)	0	3	129	297	312	133	20
	計	1,830	(6.4%)	0	4	227	563	675	314	47
要精密検査	男	588	(3.2%)	0	3	48	150	193	165	29
	女	391	(3.8%)	0	4	43	126	127	69	22
	計	979	(3.4%)	0	7	91	276	320	234	51
要治療	男	32	(0.2%)	0	0	1	10	16	4	1
	女	8	(0.1%)	0	0	1	1	5	0	1
	計	40	(0.1%)	0	0	2	11	21	4	2
治療中	男	54	(0.3%)	0	0	0	8	24	18	4
	女	11	(0.1%)	0	0	1	2	3	4	1
	計	65	(0.2%)	0	0	1	10	27	22	5
合計	男	18,174		17	692	3,288	5,662	5,474	2,643	398
	女	10,271		4	247	1,742	3,403	3,158	1,458	259
	計	28,445		21	939	5,030	9,065	8,632	4,101	657

胃内視鏡検査所見

所 見	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
十二指腸潰瘍	男	7	(0.5%)	0	0	2	4	1	0	0
	女	3	(0.4%)	0	0	0	1	2	0	0
	計	10	(0.5%)	0	0	2	5	3	0	0
慢性胃炎	男	446	(32.4%)	0	0	34	113	231	61	7
	女	162	(23.5%)	0	0	23	51	62	23	3
	計	608	(29.5%)	0	0	57	164	293	84	10
食道裂孔ヘルニア	男	242	(17.6%)	0	0	36	68	112	22	4
	女	50	(7.3%)	0	0	9	19	16	5	1
	計	292	(14.1%)	0	0	45	87	128	27	5
十二指腸潰瘍瘢痕	男	65	(4.7%)	0	0	5	23	21	15	1
	女	21	(3.1%)	0	0	1	7	8	4	1
	計	86	(4.2%)	0	0	6	30	29	19	2
萎縮性胃炎	男	42	(3.1%)	0	0	3	6	20	12	1
	女	23	(3.3%)	0	0	1	5	9	7	1
	計	65	(3.1%)	0	0	4	11	29	19	2
逆流性食道炎	男	193	(14.0%)	0	0	16	59	94	21	3
	女	20	(2.9%)	0	0	2	7	4	4	3
	計	213	(10.3%)	0	0	18	66	98	25	6
出血性胃炎	男	1	(0.1%)	0	0	0	1	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	1	(0.0%)	0	0	0	1	0	0	0
びらん性胃炎	男	84	(6.1%)	0	0	4	33	30	17	0
	女	25	(3.6%)	0	0	3	4	15	3	0
	計	109	(5.3%)	0	0	7	37	45	20	0
胃憩室	男	0	(0.0%)	0	0	0	0	0	0	0
	女	2	(0.3%)	0	0	1	0	0	1	0
	計	2	(0.1%)	0	0	1	0	0	1	0
胃潰瘍	男	2	(0.1%)	0	0	0	0	0	2	0
	女	2	(0.3%)	0	0	0	1	1	0	0
	計	4	(0.2%)	0	0	0	1	1	2	0
胃潰瘍瘢痕	男	23	(1.7%)	0	0	0	0	12	9	2
	女	11	(1.6%)	0	0	1	4	4	1	1
	計	34	(1.6%)	0	0	1	4	16	10	3
胃ポリープ	男	184	(13.4%)	0	0	21	62	81	18	2
	女	181	(26.3%)	0	0	30	69	71	9	2
	計	365	(17.7%)	0	0	51	131	152	27	4
粘膜下腫瘍	男	19	(1.4%)	0	0	0	5	10	4	0
	女	13	(1.9%)	0	0	1	4	5	2	1
	計	32	(1.6%)	0	0	1	9	15	6	1
その他の所見 (十二指腸所見)	男	67	(4.9%)	0	0	8	14	34	10	1
	女	11	(1.6%)	0	0	1	5	4	1	0
	計	78	(3.8%)	0	0	9	19	38	11	1
その他の所見 (胃所見)	男	137	(10.0%)	0	0	5	50	57	22	3
	女	59	(8.6%)	0	0	12	17	19	11	0
	計	196	(9.5%)	0	0	17	67	76	33	3
その他の所見 (食道所見)	男	49	(3.6%)	0	0	5	14	21	9	0
	女	15	(2.2%)	0	0	6	3	3	3	0
	計	64	(3.1%)	0	0	11	17	24	12	0
検診者総数	男	1,376		0	3	192	457	531	177	16
	女	688		0	6	143	234	234	66	5
	計	2,064		0	9	335	691	765	243	21

胃部×線検査所見

所 見		総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上	所 見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上	
胃 部											食 道 部										
伸展不良	男	23	(0.2%)	0	0	0	3	10	9	1	狭 窄	男	0	(0.0%)	0	0	0	0	0	0	0
	女	5	(0.1%)	0	0	0	0	2	3	0		女	1	(0.0%)	0	0	0	1	0	0	0
	計	28	(0.1%)	0	0	0	3	12	12	1		計	1	(0.0%)	0	0	0	1	0	0	0
辺縁不整	男	7	(0.1%)	0	0	0	1	3	2	1	食道裂孔ヘルニア	男	11	(0.1%)	0	0	0	1	2	3	5
	女	4	(0.0%)	0	0	0	1	2	1	0		女	7	(0.1%)	0	0	0	0	0	2	5
	計	11	(0.1%)	0	0	0	2	5	3	1		計	18	(0.1%)	0	0	0	1	2	5	10
変形	男	170	(1.3%)	0	0	13	44	50	53	10	食道炎の疑い	男	1	(0.0%)	0	0	0	0	0	1	0
	女	18	(0.2%)	0	0	0	5	6	5	2		女	0	(0.0%)	0	0	0	0	0	0	0
	計	188	(0.9%)	0	0	13	49	56	58	12		計	1	(0.0%)	0	0	0	0	0	1	0
粘膜不整	男	22	(0.2%)	0	0	2	9	2	8	1	その他の所見	男	27	(0.2%)	0	0	5	2	7	11	2
	女	13	(0.2%)	0	0	2	5	3	2	1		女	10	(0.1%)	0	0	0	1	3	3	3
	計	35	(0.2%)	0	0	4	14	5	10	2		計	37	(0.2%)	0	0	5	3	10	14	5
レリーフ集中像	男	16	(0.1%)	0	0	3	2	6	5	0	十二指腸部										
	女	6	(0.1%)	0	0	1	1	1	2	1	十二指腸球部変形	男	147	(1.2%)	0	2	12	33	66	26	8
	計	22	(0.1%)	0	0	4	3	7	7	1		女	23	(0.3%)	0	0	0	8	10	5	0
バリウム斑	男	2	(0.0%)	0	0	1	0	0	1	0		十二指腸潰瘍の疑い	計	170	(0.8%)	0	2	12	41	76	31
	女	1	(0.0%)	0	0	0	0	1	0	0	男		32	(0.3%)	0	0	5	9	11	7	0
	計	3	(0.0%)	0	0	1	0	1	1	0	女		6	(0.1%)	0	0	0	4	2	0	0
胃下垂	男	0	(0.0%)	0	0	0	0	0	0	0	十二指腸潰瘍瘢痕の疑い	計	38	(0.2%)	0	0	5	13	13	7	0
	女	1	(0.0%)	0	0	0	0	1	0	0		男	27	(0.2%)	0	0	2	7	13	5	0
	計	1	(0.0%)	0	0	0	0	1	0	0		女	6	(0.1%)	0	0	2	0	4	0	0
胃炎の疑い	男	44	(0.3%)	0	0	2	12	20	9	1	十二指腸憩室の疑い	計	33	(0.2%)	0	0	4	7	17	5	0
	女	31	(0.4%)	0	0	5	5	13	5	3		男	9	(0.1%)	0	0	0	1	4	2	2
	計	75	(0.4%)	0	0	7	17	33	14	4		女	6	(0.1%)	0	0	0	1	2	1	2
胃潰瘍瘢痕の疑い	男	2	(0.0%)	0	0	0	0	0	2	0	造影不良	計	15	(0.1%)	0	0	0	2	6	3	4
	女	0	(0.0%)	0	0	0	0	0	0	0		男	10	(0.1%)	0	0	1	3	3	3	0
	計	2	(0.0%)	0	0	0	0	0	2	0		女	2	(0.0%)	0	0	0	0	1	1	0
胃潰瘍の疑い	男	60	(0.5%)	0	0	5	16	14	20	5	十二指腸拡張不良	計	12	(0.1%)	0	0	1	3	4	4	0
	女	14	(0.2%)	0	0	1	3	4	4	2		男	1	(0.0%)	0	0	0	1	0	0	0
	計	74	(0.4%)	0	0	6	19	18	24	7		女	0	(0.0%)	0	0	0	0	0	0	0
巨大皺壁の疑い	男	8	(0.1%)	0	0	1	2	2	3	0	その他の所見	計	1	(0.0%)	0	0	0	1	0	0	0
	女	2	(0.0%)	0	0	0	0	0	1	1		男	1	(0.0%)	0	0	0	0	0	1	0
	計	10	(0.0%)	0	0	1	2	2	4	1		女	0	(0.0%)	0	0	0	0	0	0	0
隆起性病変の疑い	男	550	(4.3%)	0	3	70	180	155	115	27	検診者総数	計	1	(0.0%)	0	0	0	0	0	1	0
	女	1,022	(12.3%)	0	4	140	371	355	130	22		男	12,747		8	395	2,285	3,949	3,615	2,127	368
	計	1,572	(7.5%)	0	7	210	551	510	245	49		女	8,340		4	211	1,295	2,650	2,625	1,309	246
陥凹性病変の疑い	男	1	(0.0%)	0	0	0	0	0	1	0		計	21,087		12	606	3,580	6,599	6,240	3,436	614
	女	0	(0.0%)	0	0	0	0	0	0	0											
	計	1	(0.0%)	0	0	0	0	0	1	0											
憩室の疑い	男	49	(0.4%)	0	0	3	11	17	13	5											
	女	43	(0.5%)	0	0	2	10	16	10	5											
	計	92	(0.4%)	0	0	5	21	33	23	10											
術後胃	男	56	(0.4%)	0	0	1	5	14	30	6											
	女	6	(0.1%)	0	0	0	0	1	2	3											
	計	62	(0.3%)	0	0	1	5	15	32	9											
その他の所見	男	36	(0.3%)	0	0	2	9	8	13	4											
	女	25	(0.3%)	0	0	2	7	7	7	2											
	計	61	(0.3%)	0	0	4	16	15	20	6											

ペプシノゲン検査の性年齢別陽性率

検査項目	性別	総数			29歳以下			30-39歳			40-49歳			50-59歳			60-69歳			70歳以上		
		受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率
ペプシノゲン	男	4,801	147	3.1	304	2	0.7	885	15	1.7	1,567	26	1.7	1,667	63	3.8	361	37	10.2	17	4	23.5
	女	1,366	51	3.7	21	0	—	322	8	2.5	601	22	3.7	342	11	3.2	73	9	12.3	7	1	14.3
	計	6,167	198	3.2	325	2	0.6	1,207	23	1.9	2,168	48	2.2	2,009	74	3.7	434	46	10.6	24	5	20.8

陽性率%

ヘリコバクター・ピロリ抗体検査の性年齢別陽性率

検査項目	性別	総数			29歳以下			30-39歳			40-49歳			50-59歳			60-69歳			70歳以上		
		受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率
ヘリコバクター・ピロリ抗体	男	865	223	25.8	9	0	—	71	9	12.7	295	56	19.0	341	97	28.4	131	52	39.7	18	9	50.0
	女	634	146	23.0	14	3	21.4	82	10	12.2	206	42	20.4	231	53	22.9	82	28	34.1	19	10	52.6
	計	1,499	369	24.6	23	3	13.0	153	19	12.4	501	98	19.6	572	150	26.2	213	80	37.6	37	19	51.4

陽性率%

心電図検査判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	19,870	(76.2%)	201	1,744	4,177	5,859	5,222	2,410	257
	女	16,893	(85.0%)	197	1,763	3,317	5,086	4,445	1,822	263
	計	36,763	(80.0%)	398	3,507	7,494	10,945	9,667	4,232	520
放置可	男	2,239	(8.6%)	62	299	490	552	501	298	37
	女	1,250	(6.3%)	32	153	243	322	296	165	39
	計	3,489	(7.6%)	94	452	733	874	797	463	76
要経過観察	男	3,166	(12.1%)	12	138	383	758	1023	715	137
	女	1,476	(7.4%)	7	64	157	365	450	340	93
	計	4,642	(10.1%)	19	202	540	1123	1,473	1055	230
要精密検査	男	154	(0.6%)	3	4	9	30	51	52	5
	女	69	(0.3%)	1	6	4	18	16	18	6
	計	223	(0.5%)	4	10	13	48	67	70	11
要治療	男	17	(0.1%)	0	0	0	2	8	7	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	17	(0.0%)	0	0	0	2	8	7	0
治療中	男	622	(2.4%)	0	3	11	70	230	247	61
	女	194	(1.0%)	0	3	8	17	56	60	50
	計	816	(1.8%)	0	6	19	87	286	307	111
合計	男	26,068		278	2,188	5,070	7,271	7,035	3,729	497
	女	19,882		237	1,989	3,729	5,808	5,263	2,405	451
	計	45,950		515	4,177	8,799	13,079	12,298	6,134	948

マスター負荷心電図所見

所 見	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
陽性S T - T の変化 （ダブル）	男	2	(0.7%)	0	0	0	0	2	0	0
	女	4	(5.1%)	0	0	0	2	1	0	1
	計	6	(1.7%)	0	0	0	2	3	0	1
疑陽性S T - T の変化 （ダブル）	男	3	(1.1%)	0	0	0	0	3	0	0
	女	4	(5.1%)	0	0	0	2	1	1	0
	計	7	(2.0%)	0	0	0	2	4	1	0
不整脈の出現 （ダブル）	男	15	(5.4%)	0	1	0	6	3	5	0
	女	4	(5.1%)	0	0	0	2	0	1	1
	計	19	(5.4%)	0	1	0	8	3	6	1
疑陽性S T - T の変化 （シングル）	男	1	(0.4%)	0	0	0	0	0	1	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	1	(0.3%)	0	0	0	0	0	1	0
その他の所見	男	1	(0.4%)	0	0	0	0	1	0	0
	女	1	(1.3%)	0	0	0	0	1	0	0
	計	2	(0.6%)	0	0	0	0	2	0	0
検診者総数	男	276		0	47	8	71	103	38	9
	女	78		0	0	3	22	40	9	4
	計	354		0	47	11	93	143	47	13

自転車エルゴメータ負荷心電図所見

所 見	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
不整脈の出現	男	44	(6.5%)	0	0	6	17	12	9	0
	女	8	(4.3%)	0	0	1	5	1	0	1
	計	52	(6.0%)	0	0	7	22	13	9	1
S T - T の変化	男	20	(2.9%)	0	0	5	4	9	2	0
	女	13	(7.1%)	0	0	1	4	6	2	0
	計	33	(3.8%)	0	0	6	8	15	4	0
血圧の変化	男	244	(35.8%)	0	10	31	100	84	17	2
	女	24	(13.0%)	0	0	1	9	8	5	1
	計	268	(31.0%)	0	10	32	109	92	22	3
検診者総数	男	681		0	47	189	235	169	36	5
	女	184		0	0	51	68	45	16	4
	計	865		0	47	240	303	214	52	9

安静心電図検査所見

所 見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上	所 見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上	所 見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上			
洞性徐脈	男	745	(2.9%)	14	103	148	172	163	123	22	上室性期外収縮	男	166	(0.6%)	1	4	19	37	41	48	16	S T 低下	男	86	(0.3%)	0	1	3	17	30	29	6
	女	285	(1.4%)	8	32	53	64	88	37	3		女	110	(0.6%)	0	6	12	34	20	24	14		女	129	(0.6%)	0	3	12	28	45	31	10
	計	1,030	(2.2%)	22	135	201	236	251	160	25		計	276	(0.6%)	1	10	31	71	61	72	30		計	215	(0.5%)	0	4	15	45	75	60	16
洞性頻脈	男	114	(0.4%)	0	4	23	32	32	20	3	心室性期外収縮	男	299	(1.1%)	1	17	27	61	102	82	9	S T - T 異常	男	120	(0.5%)	0	1	8	21	39	44	7
	女	78	(0.4%)	1	4	14	28	18	9	4		女	195	(1.0%)	0	12	25	63	56	33	6		女	43	(0.2%)	0	0	5	6	11	18	3
	計	192	(0.4%)	1	8	37	60	50	29	7		計	494	(1.1%)	1	29	52	124	158	115	15		計	163	(0.4%)	0	1	13	27	50	62	10
洞性不整脈	男	220	(0.8%)	15	69	56	44	22	12	2	心房細動	男	130	(0.5%)	1	0	2	17	43	55	12	異常 Q 波の疑い	男	211	(0.8%)	0	8	12	54	69	58	10
	女	152	(0.8%)	18	44	36	23	18	8	5		女	15	(0.1%)	0	0	0	1	3	4	7		女	75	(0.4%)	0	3	2	18	28	18	6
	計	372	(0.8%)	33	113	92	67	40	20	7		計	145	(0.3%)	1	0	2	18	46	59	19		計	286	(0.6%)	0	11	14	72	97	76	16
異所性上室調律	男	59	(0.2%)	4	13	13	10	7	9	3	心房粗動	男	1	(0.0%)	0	0	0	0	1	0	0	r 波増高不良	男	7	(0.0%)	0	0	0	2	1	3	1
	女	73	(0.4%)	1	12	21	17	13	8	1		女	0	(0.0%)	0	0	0	0	0	0	0		女	21	(0.1%)	0	4	7	7	3	0	0
	計	132	(0.3%)	5	25	34	27	20	17	4		計	1	(0.0%)	0	0	0	0	1	0	0		計	28	(0.1%)	0	4	7	9	4	3	1
移動性調律	男	5	(0.0%)	0	1	1	1	0	1	1	左房負荷	男	6	(0.0%)	0	1	1	1	1	2	0	陰性 T 波	男	229	(0.9%)	0	4	15	41	92	66	11
	女	3	(0.0%)	0	0	1	2	0	0	0		女	1	(0.0%)	0	0	1	0	0	0	0		女	301	(1.5%)	1	15	32	51	107	70	25
	計	8	(0.0%)	0	1	2	3	0	1	1		計	7	(0.0%)	0	1	2	1	1	2	0		計	530	(1.2%)	1	19	47	92	199	136	36
右軸偏位 (100° 以上)	男	76	(0.3%)	2	18	28	17	7	4	0	左房負荷疑い	男	107	(0.4%)	0	2	12	25	42	23	3	T 波平低化	男	132	(0.5%)	0	2	10	33	57	25	5
	女	85	(0.4%)	0	15	22	36	12	0	0		女	41	(0.2%)	0	5	6	11	12	7	0		女	203	(1.0%)	0	3	18	51	70	46	15
	計	161	(0.4%)	2	33	50	53	19	4	0		計	148	(0.3%)	0	7	18	36	54	30	3		計	335	(0.7%)	0	5	28	84	127	71	20
左軸偏位 (-30° 以下)	男	235	(0.9%)	0	2	14	46	89	70	14	右房負荷	男	1	(0.0%)	0	0	0	1	0	0	0	T 波增高	男	385	(1.5%)	10	67	90	105	72	35	6
	女	145	(0.7%)	1	3	16	36	37	39	13		女	2	(0.0%)	0	0	0	2	0	0	0		女	8	(0.0%)	0	3	1	1	2	1	0
	計	380	(0.8%)	1	5	30	82	126	109	27		計	3	(0.0%)	0	0	0	3	0	0	0		計	393	(0.9%)	10	70	91	106	74	36	6
不定軸	男	81	(0.3%)	1	4	10	24	30	10	2	右房負荷疑い	男	11	(0.0%)	1	0	1	4	4	1	0	S T 上昇	男	311	(1.2%)	9	37	53	83	86	37	6
	女	49	(0.2%)	0	1	14	18	14	2	0		女	15	(0.1%)	0	0	1	2	8	3	1		女	23	(0.1%)	0	0	0	5	13	4	1
	計	130	(0.3%)	1	5	24	42	44	12	2		計	26	(0.1%)	1	0	2	6	12	4	1		計	334	(0.7%)	9	37	53	88	99	41	7
第一度房室ブロック	男	568	(2.2%)	2	17	56	92	167	186	48	左室肥大	男	105	(0.4%)	1	0	3	24	43	30	4	早期再分極 症候群	男	21	(0.1%)	6	4	3	3	3	2	0
	女	146	(0.7%)	0	6	15	25	45	37	18		女	16	(0.1%)	0	0	0	2	2	11	1		女	4	(0.0%)	2	1	0	0	1	0	0
	計	714	(1.6%)	2	23	71	117	212	223	66		計	121	(0.3%)	1	0	3	26	45	41	5		計	25	(0.1%)	8	5	3	3	4	2	0
第二度房室ブロック	男	4	(0.0%)	0	0	2	0	0	2	0	左室肥大疑い	男	813	(3.1%)	5	26	121	187	264	179	31	Q T 延長	男	95	(0.4%)	2	8	3	19	30	26	7
	女	6	(0.0%)	1	3	0	1	1	0	0		女	149	(0.7%)	0	1	7	19	39	63	20		女	107	(0.5%)	5	5	10	45	25	13	4
	計	10	(0.0%)	1	3	2	1	1	2	0		計	962	(2.1%)	5	27	128	206	303	242	51		計	202	(0.4%)	7	13	13	64	55	39	11
W P W 症候群	男	45	(0.2%)	0	1	11	11	19	3	0	右室肥大	男	2	(0.0%)	0	1	1	0	0	0	0	Q T 短縮	男	8	(0.0%)	0	1	1	5	1	0	0
	女	19	(0.1%)	0	1	4	6	4	4	0		女	0	(0.0%)	0	0	0	0	0	0	0		女	8	(0.0%)	0	2	2	1	1	2	0
	計	64	(0.1%)	0	2	15	17	23	7	0		計	2	(0.0%)	0	1	1	0	0	0	0		計	16	(0.0%)	0	3	3	6	2	2	0
P Q 短縮	男	40	(0.2%)	2	4	15	9	8	2	0	右室肥大疑い	男	57	(0.2%)	2	5	16	19	11	4	0	Brugada 症候群	男	5	(0.0%)	0	0	0	2	2	1	0
	女	80	(0.4%)	2	16	22	25	14	1	0		女	39	(0.2%)	0	3	8	14	12	2	0		女	1	(0.0%)	0	0	0	0	0	0	1
	計	120	(0.3%)	4	20	37	34	22	3	0		計	96	(0.2%)	2	8	24	33	23	6	0		計	6	(0.0%)	0	0	0	2	2	1	1
不完全右脚ブロック	男	404	(1.5%)	6	33	104	105	104	44	8	心筋梗塞疑い	男	2	(0.0%)	0	0	0	1	0	1	0	低電位差	男	34	(0.1%)	0	0	1	13	12	4	4
	女	123	(0.6%)	1	7	23	25	31	28	8		女	0	(0.0%)	0	0	0	0	0	0	0		女	73	(0.4%)	0	4	17	26	20	5	1
	計	527	(1.1%)	7	40	127	130	135	72	16		計	2	(0.0%)	0	0	0	1	0	1	0		計	107	(0.2%)	0	4	18	39	32	9	5
完全右脚ブロック	男	492	(1.9%)	1	12	38	99	162	148	32	陈旧性心筋梗塞	男	19	(0.1%)	0	0	1	1	6	9	2	右胸心	男	2	(0.0%)	0	0	0	1	0	1	0
	女	173	(0.9%)	0	6	18	38	50	43	18		女	0	(0.0%)	0	0	0	0	0	0	0		女	3	(0.0%)	0	0	0	3	0	0	0
	計	665	(1.4%)	1	18	56	137	212	191	50		計	19	(0.0%)	0	0	1	1	6	9	2		計	5	(0.0%)	0	0	0	4	0	1	0
ヘミブロック	男	136	(0.5%)	1	5	18	29	38	37	8	陈旧性心筋梗塞疑い	男	28	(0.1%)	0	0	0	4	12	8	4	ペースメーカー調律	男	11	(0.0%)	0	1	0	2	1	4	3
	女	57	(0.3%)	0	7	8	13	11	14	4		女	7	(0.0%)	0	0	0	2	1	3	1		女	7	(0.0%)	0	0	1	0	0	0	6
	計	193	(0.4%)	1	12	26	42	49	51	12		計	35	(0.1%)	0	0	0	6	13	11	5		計	18	(0.0%)	0	1	1	2	1	4	9
完全左脚ブロック	男	11	(0.0%)	0	0	0	1	6	4	0	心筋障害	男	1	(0.0%)	0	0	0	0	1	0	0	その他の所見	男	282	(1.1%)	0	1	4	42	112	105	18
	女	19	(0.1%)	0	0	0	6	2	7	4		女	0	(0.0%)	0	0	0	0	0	0	0		女	109	(0.5%)	0	4	4	13	32	37	19
	計	30	(0.1%)	0	0	0	7	8	11	4		計	1	(0.0%)	0	0	0	0	1	0	0		計	391	(0.9%)	0	5	8	55	144	142	37
右室伝導遅延	男	332	(1.3%)	2	36	62	78	93	55	6	心筋障害疑い	男	6	(0.0%)	0	0	0	1	2	3	0	検診者総数	男	26,066		278	2,188	5,070	7,271	7,033	3,729	497
	女	130	(0.7%)	1	16	8	23	41	32	9		女	2	(0.0%)	0	0	0	0	0	1	1		女	19,882		237	1,989	3,729	5,808	5,263	2,405	451
	計	462	(1.0%)	3	52	70	101	134	87	15		計	8	(0.0%)	0	0	0	1	2	4	1		計	45,948		515	4,177	8,799	13,079	12,296	6,134	948

腹部超音波判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	1,248	(16.5%)	4	206	366	397	221	50	4
	女	1,886	(34.9%)	3	164	482	652	429	138	18
	計	3,134	(24.2%)	7	370	848	1,049	650	188	22
放置可	男	407	(5.4%)	0	32	75	132	111	51	6
	女	405	(7.5%)	0	27	81	108	103	67	19
	計	812	(6.3%)	0	59	156	240	214	118	25
要経過観察	男	5,529	(73.2%)	3	143	704	1,573	1,978	918	210
	女	2,880	(53.3%)	1	43	306	742	963	624	201
	計	8,409	(64.9%)	4	186	1010	2,315	2,941	1,542	411
要精密検査	男	338	(4.5%)	0	5	46	79	113	63	32
	女	228	(4.2%)	0	4	22	62	66	51	23
	計	566	(4.4%)	0	9	68	141	179	114	55
要治療	男	8	(0.1%)	0	0	0	1	5	2	0
	女	3	(0.1%)	0	0	0	1	2	0	0
	計	11	(0.1%)	0	0	0	2	7	2	0
治療中	男	22	(0.3%)	0	0	2	3	12	5	0
	女	6	(0.1%)	0	0	0	0	3	2	1
	計	28	(0.2%)	0	0	2	3	15	7	1
合計	男	7,552		7	386	1,193	2,185	2,440	1,089	252
	女	5,408		4	238	891	1,565	1,566	882	262
	計	12,960		11	624	2,084	3,750	4,006	1,971	514

腹部超音波検査所見

所 見	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上	所 見	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上	所 見	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
脂肪肝の疑い	男	2,494	(33.0%)	1	45	309	774	939	366	60	胆嚢除筋腫症の疑い	男	122	(1.6%)	0	0	4	28	64	22	4	腎占拠性病変の疑い	男	6	(0.1%)	0	0	1	1	2	0	2
	女	769	(14.2%)	0	3	58	175	300	189	44		女	37	(0.7%)	0	0	1	11	15	7	3		女	1	(0.0%)	0	0	0	0	1	0	0
	計	3,263	(25.2%)	1	48	367	949	1,239	555	104		計	159	(1.2%)	0	0	5	39	79	29	7		計	7	(0.1%)	0	0	1	1	3	0	2
肝嚢胞の疑い	男	1,137	(15.1%)	0	5	71	267	482	255	57	胆嚢壁の肥厚	男	89	(1.2%)	0	3	13	19	34	16	4	水腎症の疑い	男	5	(0.1%)	0	0	1	1	1	1	1
	女	980	(18.1%)	0	8	68	224	355	259	66		女	45	(0.8%)	0	0	3	20	13	8	1		女	3	(0.1%)	0	0	0	2	0	0	1
	計	2,117	(16.3%)	0	13	139	491	837	514	123		計	134	(1.0%)	0	3	16	39	47	24	5		計	8	(0.1%)	0	0	1	3	1	1	2
肝血管腫の疑い	男	285	(3.8%)	0	8	57	92	88	28	12	胆嚢壁内結石の疑い	男	226	(3.0%)	0	0	17	59	94	47	9	重複腎孟の疑い	男	36	(0.5%)	0	1	6	9	14	4	2
	女	283	(5.2%)	0	4	65	103	77	29	5		女	58	(1.1%)	0	1	2	10	20	22	3		女	26	(0.5%)	0	0	2	8	10	6	0
	計	568	(4.4%)	0	12	122	195	165	57	17		計	284	(2.2%)	0	1	19	69	114	69	12		計	62	(0.5%)	0	1	8	17	24	10	2
肝占拠性病変の疑い	男	7	(0.1%)	0	0	0	2	2	2	1	胆嚢変形	男	6	(0.1%)	0	0	0	2	1	3	0	腎萎縮の疑い	男	6	(0.1%)	0	0	1	0	2	3	0
	女	5	(0.1%)	0	0	3	0	1	1	0		女	0	(0.0%)	0	0	0	0	0	0	0		女	6	(0.1%)	0	0	0	1	4	1	0
	計	12	(0.1%)	0	0	3	2	3	3	1		計	6	(0.0%)	0	0	0	2	1	3	0		計	12	(0.1%)	0	0	1	1	6	4	0
肝内結石の疑い	男	1	(0.0%)	0	0	0	1	0	0	0	胆嚢腫大	男	5	(0.1%)	0	0	0	0	4	1	0	腎臓描出不良	男	3	(0.0%)	0	0	0	0	2	0	1
	女	1	(0.0%)	0	0	0	0	0	1	0		女	5	(0.1%)	0	0	0	1	0	4	0		女	8	(0.1%)	0	0	1	1	0	3	3
	計	2	(0.0%)	0	0	0	1	0	1	0		計	10	(0.1%)	0	0	0	1	4	5	0		計	11	(0.1%)	0	0	1	1	2	3	4
肝内石灰化の疑い	男	57	(0.8%)	0	0	5	10	26	13	3	胆嚢摘出後	男	191	(2.5%)	0	1	3	30	84	51	22	腎盂・腎杯の拡張	男	34	(0.5%)	0	0	3	5	10	8	8
	女	32	(0.6%)	0	0	1	4	16	8	3		女	150	(2.8%)	0	0	4	21	55	50	20		女	40	(0.7%)	0	1	1	8	10	13	7
	計	89	(0.7%)	0	0	6	14	42	21	6		計	341	(2.6%)	0	1	7	51	139	101	42		計	74	(0.6%)	0	1	4	13	20	21	15
肝臓に高エコーのものあり	男	89	(1.2%)	0	3	2	16	31	29	8	胆嚢伸展不良	男	2	(0.0%)	0	0	0	1	1	0	0	その他の所見(腎臓)	男	58	(0.8%)	0	1	5	16	17	13	6
	女	39	(0.7%)	0	1	4	7	16	7	4		女	1	(0.0%)	0	0	0	0	1	0	0		女	44	(0.8%)	0	0	0	12	15	14	3
	計	128	(1.0%)	0	4	6	23	47	36	12		計	3	(0.0%)	0	0	0	1	2	0	0		計	102	(0.8%)	0	1	5	28	32	27	9
肝左葉辺縁鈍化	男	9	(0.1%)	0	0	0	0	6	2	1	胆嚢描出不良	男	69	(0.9%)	1	1	7	16	21	12	11	脾腫	男	21	(0.3%)	0	1	7	3	4	5	1
	女	1	(0.0%)	0	0	0	0	0	1	0		女	23	(0.4%)	0	0	2	6	1	7	7		女	15	(0.3%)	0	1	4	5	4	1	0
	計	10	(0.1%)	0	0	0	0	6	3	1		計	92	(0.7%)	1	1	9	22	22	19	18		計	36	(0.3%)	0	2	11	8	8	6	1
肝内エコー粗雑化	男	8	(0.1%)	0	0	0	3	4	1	0	脾管拡張の疑い	男	49	(0.6%)	0	0	0	6	17	18	8	脾嚢胞の疑い	男	7	(0.1%)	0	0	2	2	3	0	0
	女	4	(0.1%)	0	0	0	0	4	0	0		女	11	(0.2%)	0	0	1	1	3	3	3		女	11	(0.2%)	0	1	0	2	5	0	3
	計	12	(0.1%)	0	0	0	3	8	1	0		計	60	(0.5%)	0	0	1	7	20	21	11		計	18	(0.1%)	0	1	2	4	8	0	3
肝表面の不整	男	1	(0.0%)	0	0	0	0	1	0	0	脾嚢胞の疑い	男	19	(0.3%)	0	0	2	3	3	10	1	脾臓に高エコーのものあり	男	7	(0.1%)	0	0	1	0	4	1	1
	女	0	(0.0%)	0	0	0	0	0	0	0		女	32	(0.6%)	0	1	2	5	11	10	3		女	11	(0.2%)	0	1	1	1	0	6	2
	計	1	(0.0%)	0	0	0	0	1	0	0		計	51	(0.4%)	0	1	4	8	14	20	4		計	18	(0.1%)	0	1	2	1	4	7	3
慢性肝疾患の疑い	男	8	(0.1%)	0	0	0	2	4	2	0	脾占拠性病変の疑い	男	1	(0.0%)	0	0	0	0	0	1	0	副脾の疑い	男	187	(2.5%)	0	37	48	47	41	10	4
	女	0	(0.0%)	0	0	0	0	0	0	0		女	1	(0.0%)	0	0	0	1	0	0	0		女	227	(4.2%)	0	25	59	80	53	9	1
	計	8	(0.1%)	0	0	0	2	4	2	0		計	2	(0.0%)	0	0	0	1	0	1	0		計	414	(3.2%)	0	62	107	127	94	19	5
肝臓描出不良	男	158	(2.1%)	0	2	11	34	63	32	16	脾臓描出不良	男	4,089	(54.1%)	2	67	456	1,145	1,525	722	172	脾臓描出不良	男	20	(0.3%)	0	0	1	2	8	6	3
	女	15	(0.3%)	0	0	2	3	5	3	2		女	1,083	(20.0%)	0	9	68	234	349	295	128		女	8	(0.1%)	0	0	0	1	2	1	4
	計	173	(1.3%)	0	2	13	37	68	35	18		計	5,172	(39.9%)	2	76	524	1,379	1,874	1,017	300		計	28	(0.2%)	0	0	1	3	10	7	7
肝内胆管拡張の疑い	男	23	(0.3%)	0	0	0	2	8	8	5	その他の所見(肝臓・胆嚢・脾臓)	男	96	(1.3%)	0	1	5	22	39	21	8	脾血管腫の疑い	男	7	(0.1%)	0	0	3	3	1	0	0
	女	8	(0.1%)	0	0	0	1	1	5	1		女	43	(0.8%)	0	1	5	10	17	7	3		女	7	(0.1%)	0	0	1	1	2	2	1
	計	31	(0.2%)	0	0	0	3	9	13	6		計	139	(1.1%)	0	2	10	32	56	28	11		計	14	(0.1%)	0	0	4	4	3	2	1
肝外胆管拡張の疑い	男	63	(0.8%)	0	0	0	5	25	21	12	腎嚢胞の疑い	男	1,531	(20.3%)	0	20	78	303	607	409	114	腹部大動脈石灰化の疑い	男	1,231	(16.3%)	0	0	3	88	497	491	152
	女	56	(1.0%)	0	0	1	6	13	24	12		女	573	(10.6%)	0	3	43	114	169	157	87		女	688	(12.7%)	0	0	1	39	191	303	154
	計	119	(0.9%)	0	0	1	11	38	45	24		計	2,104	(16.2%)	0	23	121	417	776	566	201		計	1,919	(14.8%)	0	0	4	127	688	794	306
胆嚢ポリープの疑い	男	1,382	(18.3%)	1	48	226	460	446	176	25	腎結石の疑い	男	360	(4.8%)	0	7	31	97	155	60	10	腹部大動脈瘤の疑い	男	19	(0.3%)	0	0	0	1	7	6	5
	女	653	(12.1%)	1	16	104	201	195	116	20		女	123	(2.3%)	0	2	13	30	40	27	11		女	0	(0.0%)	0	0	0	0	0	0	0
	計	2,035	(15.7%)	2	64	330	661	641	292	45		計	483	(3.7%)	0	9	44	127	195	87	21		計	19	(0.1%)	0	0	0	1	7	6	5
胆石の疑い	男	278	(3.7%)	0	0	11	63	131	61	12	腎臓に高エコーのものあり	男	843	(11.2%)	0	20	77	220	335	161	30	その他の所見(脾臓・他)	男	28	(0.4%)	0	0	1	6	7	11	3
	女	165	(3.1%)	0	5	8	39	54	46	13		女	393	(7.3%)	0	8	33	111	145	78	18		女	24	(0.4%)	0	1	1	12	3	5	2
	計	443	(3.4%)	0	5	19	102	185	107	25		計	1,236	(9.5%)	0	28	110	331	480	239	48		計	52	(0.4%)	0	1	2	18	10	16	5
胆砂・胆泥の疑い	男	55	(0.7%)	0	1	6	8	25	8	7	腎血管筋脂肪腫の疑い	男	41	(0.5%)	0	3	2	15	14	6	1	検診者総数	男	7,552		7	386	1,193	2,185	2,440	1,089	252
	女	43	(0.8%)	0	1	3	8	15	12	4		女	66	(1.2%)	0	0	3	21	24	13	5		女	5,408		4	238	891	1,565	1,566	882	262
	計	98	(0.8%)	0	2	9	16	40	20	11		計	107	(0.8%)	0	3	5	36	38	19	6		計	12,960		11	624	2,084	3,750	4,006	1,971	5145

眼底写真判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	5,379	(60.8%)	7	412	1,179	1,846	1,516	395	24
	女	3,464	(71.0%)	5	239	781	1,256	936	225	22
	計	8,843	(64.4%)	12	651	1,960	3,102	2,452	620	46
放置可	男	585	(6.6%)	0	3	20	162	288	102	10
	女	334	(6.8%)	0	2	15	79	159	64	15
	計	919	(6.7%)	0	5	35	241	447	166	25
要経過観察	男	2,297	(26.0%)	0	8	60	465	1077	599	88
	女	789	(16.2%)	0	2	29	129	334	245	50
	計	3,086	(22.5%)	0	10	89	594	1411	844	138
要精密検査	男	418	(4.7%)	0	6	25	79	180	107	21
	女	184	(3.8%)	0	3	13	27	68	57	16
	計	602	(4.4%)	0	9	38	106	248	164	37
要治療	男	1	(0.0%)	0	0	0	0	1	0	0
	女	1	(0.0%)	0	0	0	0	0	1	0
	計	2	(0.0%)	0	0	0	0	1	1	0
治療中	男	165	(1.9%)	0	0	4	20	74	53	14
	女	104	(2.1%)	0	0	1	10	37	40	16
	計	269	(2.0%)	0	0	5	30	111	93	30
合計	男	8,845		7	429	1,288	2,572	3,136	1,256	157
	女	4,876		5	246	839	1,501	1,534	632	119
	計	13,721		12	675	2,127	4,073	4,670	1,888	276

眼底検査所見

所 見	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上	所 見	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
細動脈狭細化	男	283	(3.2%)	0	0	4	51	148	71	9	白内障の疑い	男	348	(3.9%)	0	0	0	20	115	153	60
	女	95	(1.9%)	0	0	2	19	48	19	7		女	213	(4.4%)	0	0	1	10	45	109	48
	計	378	(2.7%)	0	0	6	70	196	90	16		計	561	(4.1%)	0	0	1	30	160	262	108
細動脈口径不同	男	164	(1.8%)	0	0	4	41	84	30	5	視神経乳頭異常	男	18	(0.2%)	0	0	0	4	5	9	0
	女	53	(1.1%)	0	0	5	8	28	11	1		女	6	(0.1%)	0	0	0	1	3	2	0
	計	217	(1.6%)	0	0	9	49	112	41	6		計	24	(0.2%)	0	0	0	5	8	11	0
網膜浮腫	男	1	(0.0%)	0	0	0	0	0	1	0	視神経乳頭異常の疑い	男	371	(4.2%)	0	10	33	94	156	69	9
	女	0	(0.0%)	0	0	0	0	0	0	0		女	156	(3.2%)	0	3	16	43	60	28	6
	計	1	(0.0%)	0	0	0	0	0	1	0		計	527	(3.8%)	0	13	49	137	216	97	15
眼底出血	男	55	(0.6%)	0	0	3	13	24	13	2	光凝固後	男	28	(0.3%)	0	0	1	4	12	10	1
	女	21	(0.4%)	0	1	2	2	12	4	0		女	10	(0.2%)	0	0	0	3	2	4	1
	計	76	(0.6%)	0	1	5	15	36	17	2		計	38	(0.3%)	0	0	1	7	14	14	2
細動脈反射増強	男	615	(6.9%)	0	0	9	115	327	150	14	網膜有鞘神経繊維の疑い	男	60	(0.7%)	0	2	3	17	23	15	0
	女	245	(5.0%)	0	0	3	36	112	85	9		女	17	(0.3%)	0	1	1	6	6	3	0
	計	860	(6.3%)	0	0	12	151	439	235	23		計	77	(0.6%)	0	3	4	23	29	18	0
動静脈交叉現象	男	1,409	(15.9%)	0	0	21	312	726	321	29	網膜静脈叢出血の疑い	男	46	(0.5%)	0	0	3	13	17	9	4
	女	443	(9.1%)	0	1	10	79	209	122	22		女	21	(0.4%)	0	0	2	7	6	5	1
	計	1,852	(13.5%)	0	1	31	391	935	443	51		計	67	(0.5%)	0	0	5	20	23	14	5
毛細血管瘤（赤色点）	男	12	(0.1%)	0	0	1	2	7	2	0	瞳孔の開き不充分	男	150	(1.7%)	0	0	1	21	58	58	12
	女	3	(0.1%)	0	0	1	0	2	0	0		女	44	(0.9%)	0	0	1	8	16	16	3
	計	15	(0.1%)	0	0	2	2	9	2	0		計	194	(1.4%)	0	0	2	29	74	74	15
新生血管	男	1	(0.0%)	0	0	0	0	0	1	0	黄斑部異常の疑い	男	61	(0.7%)	0	0	1	14	19	26	1
	女	0	(0.0%)	0	0	0	0	0	0	0		女	41	(0.8%)	0	0	2	4	14	21	0
	計	1	(0.0%)	0	0	0	0	0	1	0		計	102	(0.7%)	0	0	3	18	33	47	1
静脈拡張	男	2	(0.0%)	0	0	0	0	1	1	0	右眼のみ撮影	男	21	(0.2%)	0	0	2	2	10	5	2
	女	0	(0.0%)	0	0	0	0	0	0	0		女	7	(0.1%)	0	0	0	0	4	3	0
	計	2	(0.0%)	0	0	0	0	1	1	0		計	28	(0.2%)	0	0	2	2	14	8	2
点状出血	男	57	(0.6%)	0	0	0	12	36	9	0	左眼のみ撮影	男	18	(0.2%)	0	0	3	6	5	3	1
	女	6	(0.1%)	0	0	0	2	3	1	0		女	7	(0.1%)	0	0	0	1	3	3	0
	計	63	(0.5%)	0	0	0	14	39	10	0		計	25	(0.2%)	0	0	3	7	8	6	1
斑状出血	男	55	(0.6%)	0	0	0	13	31	10	1	その他の所見	男	427	(4.8%)	0	6	30	89	191	97	14
	女	7	(0.1%)	0	0	0	1	1	5	0		女	232	(4.8%)	0	1	18	36	85	68	24
	計	62	(0.5%)	0	0	0	14	32	15	1		計	659	(4.8%)	0	7	48	125	276	165	38
網膜白斑	男	721	(8.1%)	0	0	10	96	361	217	37	検査者総数	男	8,873		8	440	1,294	2,577	3,139	1,258	157
	女	264	(5.4%)	0	0	2	32	126	88	16		女	4,877		5	246	839	1,502	1,534	632	119
	計	985	(7.2%)	0	0	12	128	487	305	53		計	13,750		13	686	2,133	4,079	4,673	1,890	276

肺機能判定結果

判 定	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	5,739 (93.5%)	149	389	1076	1,665	1,707	624	129
	女	2,938 (95.7%)	93	86	530	802	883	443	101
	計	8,677 (94.2%)	242	475	1,606	2,467	2,590	1067	230
放置可	男	332 (5.4%)	4	8	28	72	97	94	29
	女	112 (3.6%)	0	3	11	29	32	26	11
	計	444 (4.8%)	4	11	39	101	129	120	40
要経過観察	男	49 (0.8%)	1	0	0	9	13	17	9
	女	17 (0.6%)	0	1	0	2	2	9	3
	計	66 (0.7%)	1	1	0	11	15	26	12
要精密検査	男	15 (0.2%)	0	0	1	0	5	9	0
	女	4 (0.1%)	0	0	0	2	0	1	1
	計	19 (0.2%)	0	0	1	2	5	10	1
要治療	男	0 (0.0%)	0	0	0	0	0	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	0 (0.0%)	0	0	0	0	0	0	0
治療中	男	1 (0.0%)	0	0	0	0	0	0	1
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	1 (0.0%)	0	0	0	0	0	0	1
合 計	男	6,136	154	397	1105	1,746	1,822	744	168
	女	3,071	93	90	541	835	917	479	116
	計	9,207	247	487	1,646	2,581	2,739	1,223	284

肺機能検査の性年齢別平均値

検査項目	総数		29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
%肺活量 %	男	6,124 103.8	563 101.4	1107 107.4	1,737 105.9	1,814 103.7	734 98.2	169 94.0						
	女	3,048 103.1	183 97.5	534 103.4	821 105.2	916 103.6	476 101.6	118 98.2						
	計	9,172 103.6	746 100.5	1,641 106.1	2,558 105.7	2,730 103.7	1,210 99.5	287 95.7						
一秒率 %	男	6,124 81.1	563 85.6	1107 82.7	1,737 81.1	1,814 80.1	734 78.9	169 77.3						
	女	3,048 82.4	183 87.6	534 85.1	821 82.4	916 81.5	476 80.3	118 78.2						
	計	9,172 81.6	746 86.1	1,641 83.5	2,558 81.5	2,730 80.5	1,210 79.4	287 77.7						

聴力判定結果

判 定	総数			19歳以下		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
	男																
異常なし	男	24,117	(85.2%)	179	3,288	5,569	6,937	5,804	2,193	147							
	女	19,816	(93.6%)	108	3,216	4,307	5,580	4,724	1,759	122							
	計	43,933	(88.8%)	287	6,504	9,876	12,517	10,528	3,952	269							
放置可	男	1	(0.0%)	0	0	0	0	1	0	0							
	女	0	(0.0%)	0	0	0	0	0	0	0							
	計	1	(0.0%)	0	0	0	0	1	0	0							
要経過観察	男	4,179	(14.8%)	1	52	228	559	1,425	1,624	290							
	女	1,362	(6.4%)	3	36	79	178	417	510	139							
	計	5,541	(11.2%)	4	88	307	737	1,842	2,134	429							
要精密検査	男	1	(0.0%)	0	0	0	0	0	1	0							
	女	0	(0.0%)	0	0	0	0	0	0	0							
	計	1	(0.0%)	0	0	0	0	0	1	0							
要治療	男	0	(0.0%)	0	0	0	0	0	0	0							
	女	0	(0.0%)	0	0	0	0	0	0	0							
	計	0	(0.0%)	0	0	0	0	0	0	0							
治療中	男	0	(0.0%)	0	0	0	0	0	0	0							
	女	0	(0.0%)	0	0	0	0	0	0	0							
	計	0	(0.0%)	0	0	0	0	0	0	0							
合計	男	28,298		180	3,340	5,797	7,496	7,230	3,818	437							
	女	21,178		111	3,252	4,386	5,758	5,141	2,269	261							
	計	49,476		291	6,592	10,183	13,254	12,371	6,087	698							

聴力検査の性年齢別平均値

検査項目	総数			29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
	男														
聴力・右・ 5 0 0 H z d B	男	224	16.3	0	—	4	6.3	61	12.8	107	14.4	40	23.9	12	27.9
	女	82	16.5	0	—	2	15.0	25	10.2	36	15.6	13	22.7	6	35.0
	計	306	16.3	0	—	6	9.2	86	12.0	143	14.7	53	23.6	18	30.3
聴力・右・ 1 K H z d B	男	224	17.7	0	—	4	11.3	61	13.4	107	16.7	40	24.1	12	28.8
	女	82	17.4	0	—	2	20.0	25	11.6	36	16.1	13	23.1	6	35.8
	計	306	17.6	0	—	6	14.2	86	12.9	143	16.6	53	23.9	18	31.1
聴力・右・ 2 K H z d B	男	224	19.1	0	—	4	10.0	61	14.2	107	17.5	40	25.3	12	40.0
	女	82	17.9	0	—	2	10.0	25	11.6	36	16.1	13	26.5	6	38.3
	計	306	18.7	0	—	6	10.0	86	13.4	143	17.2	53	25.6	18	39.4
聴力・右・ 4 K H z d B	男	224	23.5	0	—	4	8.8	61	12.8	107	22.9	40	34.8	12	50.4
	女	82	13.5	0	—	2	12.5	25	6.8	36	13.2	13	20.0	6	30.0
	計	306	20.8	0	—	6	10.0	86	11.1	143	20.5	53	31.1	18	43.6
聴力・左・ 5 0 0 H z d B	男	224	15.7	0	—	4	3.8	61	10.7	107	14.6	40	24.0	12	27.1
	女	82	15.9	0	—	2	5.0	25	10.6	36	15.8	13	21.5	6	30.0
	計	306	15.8	0	—	6	4.2	86	10.7	143	14.9	53	23.4	18	28.1
聴力・左・ 1 K H z d B	男	224	16.7	0	—	4	7.5	61	11.7	107	15.8	40	23.5	12	30.0
	女	82	15.1	0	—	2	7.5	25	10.4	36	14.3	13	21.9	6	27.5
	計	306	16.3	0	—	6	7.5	86	11.3	143	15.5	53	23.1	18	29.2
聴力・左・ 2 K H z d B	男	224	18.3	0	—	4	8.8	61	11.7	107	17.4	40	24.6	12	41.7
	女	82	16.0	0	—	2	5.0	25	9.8	36	15.3	13	23.1	6	34.2
	計	306	17.7	0	—	6	7.5	86	11.2	143	16.9	53	24.3	18	39.2
聴力・左・ 4 K H z d B	男	224	23.2	0	—	4	8.8	61	13.1	107	22.2	40	34.1	12	51.7
	女	82	14.1	0	—	2	10.0	25	6.4	36	13.9	13	21.2	6	33.3
	計	306	20.7	0	—	6	9.2	86	11.2	143	20.1	53	30.9	18	45.6

精密聴力検査の性年齢別平均値

検査項目	総数			29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
	男														
精密聴力・右 2 5 0 H z d B	男	367	24.6	102	21.2	37	20.1	100	23.1	87	28.3	41	33.2	0	—
	女	0	—	0	—	0	—	0	—	0	—	0	—	0	—
	計	367	24.6	102	21.2	37	20.1	100	23.1	87	28.3	41	33.2	0	—
精密聴力・右 5 0 0 H z d B	男	367	20.8	102	17.4	37	17.2	100	18.6	87	24.7	41	29.2	0	—
	女	0	—	0	—	0	—	0	—	0	—	0	—	0	—
	計	367	20.8	102	17.4	37	17.2	100	18.6	87	24.7	41	29.2	0	—
精密聴力・右 1 K H z d B	男	367	18.5	102	13.1	37	15.7	100	16.0	87	25.5	41	25.7	0	—
	女	0	—	0	—	0	—	0	—	0	—	0	—	0	—
	計	367	18.5	102	13.1	37	15.7	100	16.0	87	25.5	41	25.7	0	—
精密聴力・右 2 K H z d B	男	367	18.6	102	11.7	37	13.1	100	16.6	87	26.6	41	28.9	0	—
	女	0	—	0	—	0	—	0	—	0	—	0	—	0	—
	計	367	18.6	102	11.7	37	13.1	100	16.6	87	26.6	41	28.9	0	—
精密聴力・右 4 K H z d B	男	367	20.9	102	7.2	37	14.6	100	15.6	87	35.0	41	43.5	0	—
	女	0	—	0	—	0	—	0	—	0	—	0	—	0	—
	計	367	20.9	102	7.2	37	14.6	100	15.6	87	35.0	41	43.5	0	—
精密聴力・右 8 K H z d B	男	367	24.2	102	9.3	37	14.6	100	18.6	87	39.6	41	50.6	0	—
	女	0	—	0	—	0	—	0	—	0	—	0	—	0	—
	計	367	24.2	102	9.3	37	14.6	100	18.6	87	39.6	41	50.6	0	—
精密聴力・左 2 5 0 H z d B	男	367	25.1	102	19.9	37	21.1	100	23.4	87	30.0	41	35.5	0	—
	女	0	—	0	—	0	—	0	—	0	—	0	—	0	—
	計	367	25.1	102	19.9	37	21.1	100	23.4	87	30.0	41	35.5	0	—
精密聴力・左 5 0 0 H z d B	男	367	21.2	102	16.3	37	16.4	100	19.2	87	26.7	41	30.5	0	—
	女	0	—	0	—	0	—	0	—	0	—	0	—	0	—
	計	367	21.2	102	16.3	37	16.4	100	19.2	87	26.7	41	30.5	0	—
精密聴力・左 1 K H z d B	男	367	18.4	102	12.2	37	13.8	100	16.0	87	25.9	41	27.8	0	—
	女	0	—	0	—	0	—	0	—	0	—	0	—	0	—
	計	367	18.4	102	12.2	37	13.8	100	16.0	87	25.9	41	27.8	0	—
精密聴力・左 2 K H z d B	男	367	18.5	102	10.7	37	13.8	100	15.6	87	26.6	41	32.3	0	—
	女	0	—	0	—	0	—	0	—	0	—	0	—	0	—
	計	367	18.5	102	10.7	37	13.8	100	15.6	87	26.6	41	32.3	0	—
精密聴力・左 4 K H z d B	男	367	22.4	102	7.2	37	14.5	100	17.8	87	37.9	41	46.1	0	—
	女	0	—	0	—	0	—	0	—	0	—	0	—	0	—
	計	367	22.4	102	7.2	37	14.5	100	17.8	87	37.9	41	46.1	0	—
精密聴力・左 8 K H z d B	男	367	23.9	102	8.0	37	12.8	100	19.3	87	39.1	41	52.7	0	—
	女	0	—	0	—	0	—	0	—	0	—	0	—	0	—
	計	367	23.9	102	8.0	37	12.8	100	19.3	87	39.1	41	52.7	0	—

血圧判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	18,308	(60.2%)	334	3,423	5,134	4,781	3,259	1,244	133
	女	18,807	(79.3%)	302	3,657	4,627	5,191	3,676	1,203	151
	計	37,115	(68.6%)	636	7,080	9,761	9,972	6,935	2,447	284
放置可	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
要経過観察	男	7,168	(23.6%)	33	599	1,315	1,945	2,076	1,077	123
	女	2,769	(11.7%)	4	95	289	713	1,020	566	82
	計	9,937	(18.4%)	37	694	1,604	2,658	3,096	1,643	205
要精密検査	男	690	(2.3%)	0	15	71	185	247	152	20
	女	238	(1.0%)	0	3	17	59	91	53	15
	計	928	(1.7%)	0	18	88	244	338	205	35
要治療	男	192	(0.6%)	0	2	29	56	70	32	3
	女	57	(0.2%)	0	1	8	13	24	9	2
	計	249	(0.5%)	0	3	37	69	94	41	5
治療中	男	4,060	(13.3%)	0	4	96	631	1,679	1,423	227
	女	1,831	(7.7%)	0	2	26	223	741	645	194
	計	5,891	(10.9%)	0	6	122	854	2,420	2,068	421
合 計	男	30,418		367	4,043	6,645	7,598	7,331	3,928	506
	女	23,702		306	3,758	4,967	6,199	5,552	2,476	444
	計	54,120		673	7,801	11,612	13,797	12,883	6,404	950

血圧（安静時・負荷時）の性年齢別平均値

検査項目	総数			29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
安静時血圧・最大 mmHg	男	31,098	124.3	4,621	119.6	6,766	120.5	7,738	122.9	7,486	127.7	3,980	131.5	507	131.5
	女	23,836	114.2	4,101	107.0	5,023	108.4	6,235	113.1	5,557	119.8	2,476	125.8	444	128.0
	計	54,934	119.9	8,722	113.7	11,789	115.3	13,973	118.5	13,043	124.3	6,456	129.3	951	129.8
安静時血圧・最小 mmHg	男	31,098	77.1	4,621	68.8	6,766	73.9	7,738	78.4	7,486	81.2	3,980	81.4	507	77.8
	女	23,836	70.2	4,101	64.7	5,023	67.3	6,235	70.5	5,557	74.2	2,476	75.1	444	72.7
	計	54,934	74.1	8,722	66.9	11,789	71.1	13,973	74.9	13,043	78.2	6,456	79.0	951	75.4
負荷後血圧・最大 mmHg	男	276	137.9	47	129.2	8	139.3	71	136.6	103	139.8	38	143.4	9	146.0
	女	78	129.5	0	0.0	3	118.0	22	127.0	40	131.0	9	130.0	4	136.0
	計	354	136.0	47	129.2	11	133.5	93	134.3	143	137.4	47	140.8	13	142.9
負荷後血圧・最小 mmHg	男	276	77.2	47	70.9	8	76.0	71	76.7	103	80.4	38	77.1	9	80.4
	女	78	71.1	0	0.0	3	66.0	22	68.6	40	72.7	9	72.9	4	69.0
	計	354	75.9	47	70.9	11	73.3	93	74.8	143	78.3	47	76.3	13	76.9

眼圧の性年齢別平均値

検査項目	総数			29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
眼圧・右 mmHg	男	4,201	14.5	380	14.8	796	14.5	1,191	14.7	1,370	14.2	426	14.5	38	12.9
	女	3,005	14.3	213	14.9	538	14.2	1,015	14.3	970	14.1	243	14.2	26	12.9
	計	7,206	14.4	593	14.8	1,334	14.4	2,206	14.5	2,340	14.2	669	14.4	64	12.9
眼圧・左 mmHg	男	4,195	14.6	380	14.9	795	14.6	1,191	14.8	1,366	14.4	425	14.7	38	13.8
	女	3,006	14.4	213	14.9	538	14.3	1,015	14.6	971	14.3	243	14.5	26	13.5
	計	7,201	14.6	593	14.9	1,333	14.5	2,206	14.7	2,337	14.4	668	14.7	64	13.7

内科診察判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	21,094	(97.1%)	215	1,530	3,730	6,333	5,956	2,995	335
	女	14,641	(95.7%)	110	992	2,491	4,729	4,275	1,793	251
	計	35,735	(96.5%)	325	2,522	6,221	11,062	10,231	4,788	586
放置可	男	364	(1.7%)	1	17	45	89	97	96	19
	女	281	(1.8%)	3	6	30	68	99	63	12
	計	645	(1.7%)	4	23	75	157	196	159	31
要経過観察	男	170	(0.8%)	0	3	23	37	55	44	8
	女	305	(2.0%)	1	5	43	118	92	41	5
	計	475	(1.3%)	1	8	66	155	147	85	13
要精密検査	男	35	(0.2%)	0	0	3	9	9	12	2
	女	32	(0.2%)	1	1	7	8	11	4	0
	計	67	(0.2%)	1	1	10	17	20	16	2
要治療	男	8	(0.0%)	0	0	0	1	3	2	2
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	8	(0.0%)	0	0	0	1	3	2	2
治療中	男	59	(0.3%)	0	0	1	8	24	18	8
	女	44	(0.3%)	0	1	10	13	10	8	2
	計	103	(0.3%)	0	1	11	21	34	26	10
合計	男	21,730		216	1,550	3,802	6,477	6,144	3,167	374
	女	15,303		115	1005	2,581	4,936	4,487	1,909	270
	計	37,033		331	2,555	6,383	11,413	10,631	5,076	644

内科診察所見

所 見	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上	所 見	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上	所 見	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
眼瞼結膜充血あり	男	2	(0.0%)	0	0	0	0	0	2	0	湿性ラ音	男	1	(0.0%)	0	0	0	0	0	0	1	浮腫	男	16	(0.1%)	0	0	1	3	8	2	2
	女	30	(0.2%)	0	0	2	18	8	2	0		女	3	(0.0%)	0	0	1	0	1	1	0		女	10	(0.1%)	0	0	0	3	4	3	0
	計	30	(0.1%)	0	0	2	18	8	2	0		計	4	(0.0%)	0	0	1	0	1	1	1		計	23	(0.1%)	0	0	1	3	12	5	2
眼球結膜黄斑あり	男	1	(0.0%)	0	0	0	1	0	0	0	呼吸音の減弱	男	2	(0.0%)	0	0	0	0	2	0	0	黄色腫	男	6	(0.0%)	0	0	1	2	3	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0		女	0	(0.0%)	0	0	0	0	0	0	0		女	3	(0.0%)	0	0	0	2	0	1	0
	計	1	(0.0%)	0	0	0	1	0	0	0		計	2	(0.0%)	0	0	0	0	2	0	0		計	7	(0.0%)	0	0	1	2	3	1	0
扁桃腺肥大	男	70	(0.3%)	1	12	25	23	6	3	0	肝触知	男	2	(0.0%)	0	0	0	1	1	0	0	術創あり	男	264	(1.2%)	0	3	20	51	79	94	17
	女	36	(0.2%)	0	7	10	11	7	1	0		女	0	(0.0%)	0	0	0	0	0	0	0		女	211	(1.4%)	1	1	16	51	80	51	11
	計	106	(0.3%)	1	19	35	34	13	4	0		計	2	(0.0%)	0	0	0	1	1	0	0		計	424	(1.1%)	1	4	36	51	159	145	28
頸部リンパ節腫脹	男	6	(0.0%)	0	1	0	5	0	0	0	脾触知	男	0	(0.0%)	0	0	0	0	0	0	0	皮膚の異常	男	10	(0.0%)	0	2	5	0	2	1	0
	女	9	(0.1%)	0	0	3	2	2	2	0		女	1	(0.0%)	0	0	0	0	0	1	0		女	2	(0.0%)	0	1	1	0	0	0	0
	計	15	(0.0%)	0	1	3	7	2	2	0		計	1	(0.0%)	0	0	0	0	0	1	0		計	11	(0.0%)	0	3	5	0	2	1	0
甲状腺腫大	男	12	(0.1%)	0	0	2	2	7	1	0	腹部圧痛	男	33	(0.2%)	0	0	5	6	13	8	1	知覚異常	男	1	(0.0%)	0	0	0	0	1	0	0
	女	160	(1.0%)	0	2	36	61	47	14	0		女	60	(0.4%)	0	0	10	20	20	8	2		女	0	(0.0%)	0	0	0	0	0	0	0
	計	172	(0.5%)	0	2	38	63	54	15	0		計	85	(0.2%)	0	0	15	26	33	8	3		計	1	(0.0%)	0	0	0	0	1	0	0
胸郭変形	男	9	(0.0%)	0	1	2	3	3	0	0	腹部腫瘍触知	男	4	(0.0%)	0	0	0	0	2	1	1	皮膚の乾燥・癢じわ	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0		女	28	(0.2%)	0	0	4	15	7	2	0		女	1	(0.0%)	0	0	0	1	0	0	0
	計	9	(0.0%)	0	1	2	3	3	0	0		計	32	(0.1%)	0	0	4	15	9	3	1		計	1	(0.0%)	0	0	0	1	0	0	0
心拍不整	男	112	(0.5%)	0	1	3	21	33	43	11	下肢静脈瘤	男	11	(0.1%)	0	0	1	3	4	1	2	爪の異常	男	1	(0.0%)	0	0	0	1	0	0	0
	女	37	(0.2%)	4	1	1	11	14	4	2		女	29	(0.2%)	0	0	1	2	13	11	2		女	0	(0.0%)	0	0	0	0	0	0	0
	計	147	(0.4%)	4	1	4	32	47	47	12		計	37	(0.1%)	0	0	1	5	17	12	2		計	1	(0.0%)	0	0	0	1	0	0	0
収縮期雑音	男	41	(0.2%)	0	1	6	5	8	10	3	関節変形	男	2	(0.0%)	0	0	0	2	0	0	0	皮膚・粘膜の異常	男	4	(0.0%)	0	1	2	1	0	0	0
	女	35	(0.2%)	0	2	4	4	8	8	2		女	2	(0.0%)	0	0	0	0	1	1	0		女	0	(0.0%)	0	0	0	0	0	0	0
	計	76	(0.1%)	0	3	10	9	8	18	5		計	4	(0.0%)	0	0	0	2	1	1	0		計	4	(0.0%)	0	1	2	1	0	0	0
拡張期雑音	男	5	(0.0%)	0	0	2	3	1	0	1	関節腫脹	男	0	(0.0%)	0	0	0	0	0	0	0	軽度の腰痛症状	男	6	(0.0%)	0	0	1	4	1	0	0
	女	4	(0.0%)	0	0	0	0	0	0	0		女	2	(0.0%)	0	0	0	0	2	0	0		女	6	(0.0%)	0	0	1	2	2	1	0
	計	5	(0.0%)	0	0	2	3	1	0	1		計	2	(0.0%)	0	0	0	0	2	0	0		計	12	(0.0%)	0	0	2	6	3	1	0
連続性雑音	男	0	(0.0%)	0	0	0	0	0	0	0	脊椎異常	男	3	(0.0%)	0	1	0	1	0	1	0	前屈制限・前屈時疼痛	男	2	(0.0%)	0	0	0	2	0	0	0
	女	1	(0.0%)	0	0	0	0	1	0	0		女	2	(0.0%)	0	0	0	0	1	1	0		女	1	(0.0%)	0	0	0	0	1	0	0
	計	1	(0.0%)	0	0	0	0	1	0	0		計	4	(0.0%)	0	1	0	1	1	1	0		計	3	(0.0%)	0	0	0	2	1	0	0
3音、4音	男	41	(0.0%)	0	0	0	1	1	0	0	知覚障害	男	0	(0.0%)	0	0	0	0	0	0	0	その他の所見	男	84	(0.4%)	3	21	16	17	18	6	3
	女	35	(0.0%)	0	0	0	0	0	0	0		女	1	(0.0%)	0	0	1	0	0	0	0		女	24	(0.2%)	0	1	2	10	5	6	0
	計	76	(0.0%)	0	0	0	1	1	0	0		計	1	(0.0%)	0	0	1	0	0	0	0		計	107	(0.3%)	3	22	18	27	23	11	3
乾性ラ音	男	5	(0.1%)	0	0	2	7	8	7	2	麻痺	男	4	(0.0%)	0	0	1	0	3	0	0	検診者総数	男	21,730		216	1,550	3,802	6,477	6,144	3,167	374
	女	4	(0.0%)	2	0	0	1	1	2	1		女	2	(0.0%)	0	0	1	0	1	0	0		女	15,303		115	1005	2,581	4,936	4,487	1,909	270
	計	5	(0.1%)	2	0	2	8	9	9	3		計	6	(0.0%)	0	0	2	0	4	0	0		計	37,033		331	2,555	6,383	11,413	10,631	5,076	644

大腸検査判定結果

判 定	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	408 (92.1%)	0	0	36	126	138	94	14
	女	210 (92.5%)	0	0	22	58	81	39	10
	計	618 (92.2%)	0	0	58	184	219	133	24
放置可	男	12 (2.7%)	0	0	1	5	4	2	0
	女	7 (3.1%)	0	0	0	2	2	3	0
	計	19 (2.8%)	0	0	1	7	6	5	0
要経過観察	男	21 (4.7%)	0	0	2	4	5	6	4
	女	8 (3.5%)	0	0	1	1	2	3	1
	計	29 (4.3%)	0	0	3	5	7	9	5
要精密検査	男	2 (0.5%)	0	0	0	1	0	1	0
	女	2 (0.9%)	0	0	0	0	0	2	0
	計	4 (0.6%)	0	0	0	1	0	3	0
要治療	男	0 (0.0%)	0	0	0	0	0	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	0 (0.0%)	0	0	0	0	0	0	0
治療中	男	0 (0.0%)	0	0	0	0	0	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	0 (0.0%)	0	0	0	0	0	0	0
合 計	男	443	0	0	39	136	147	103	18
	女	227	0	0	23	61	85	47	11
	計	670	0	0	62	197	232	150	29

直腸診所見

所 見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
外痔核	男	6 (1.4%)	0	0	1	1	3	1	0
	女	10 (4.4%)	0	0	0	2	3	4	1
	計	16 (2.4%)	0	0	1	3	6	5	1
外痔核の疑い	男	1 (0.2%)	0	0	0	0	0	1	0
	女	1 (0.4%)	0	0	0	0	0	1	0
	計	2 (0.3%)	0	0	0	0	0	2	0
内痔核の疑い	男	2 (0.5%)	0	0	0	1	0	1	0
	女	3 (1.3%)	0	0	0	1	1	1	0
	計	5 (0.7%)	0	0	0	2	1	2	0
痔瘻	男	2 (0.5%)	0	0	1	1	0	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	2 (0.3%)	0	0	1	1	0	0	0
肛門ポリープ	男	2 (0.5%)	0	0	0	1	0	1	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	2 (0.3%)	0	0	0	1	0	1	0
肛門ポリープの疑い	男	1 (0.2%)	0	0	0	1	0	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	1 (0.1%)	0	0	0	1	0	0	0
前立腺肥大	男	3 (0.7%)	0	0	0	0	0	2	1
前立腺肥大の疑い	男	9 (2.0%)	0	0	1	1	2	2	3
前立腺腫瘍の疑い	男	3 (0.7%)	0	0	0	1	2	0	0
直腸腫瘍の疑い	男	1 (0.2%)	0	0	0	0	0	1	0
	女	2 (0.9%)	0	0	0	0	0	2	0
	計	3 (0.4%)	0	0	0	0	0	3	0
肛門周囲湿疹	男	2 (0.5%)	0	0	0	1	1	0	0
	女	1 (0.4%)	0	0	1	0	0	0	0
	計	3 (0.4%)	0	0	1	1	1	0	0
痔疾患術後	男	1 (0.2%)	0	0	0	1	0	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	1 (0.1%)	0	0	0	1	0	0	0
その他の所見	男	2 (0.5%)	0	0	0	1	1	0	0
	女	1 (0.4%)	0	0	0	0	1	0	0
	計	3 (0.4%)	0	0	0	1	2	0	0
検診者総数	男	443	0	0	39	136	147	103	18
	女	227	0	0	23	61	85	47	11
	計	670	0	0	62	197	232	150	29

乳房検査判定結果

判 定	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	9,493	(81.6%)	4	546	1,528	2,808	2,715	1,515	377
放置可	513	(4.4%)	0	22	100	128	169	76	18
要経過観察	1,014	(8.7%)	0	83	265	354	201	85	26
要精密検査	612	(5.3%)	1	27	93	222	153	87	29
要治療	0	(0.0%)	0	0	0	0	0	0	0
治療中	7	(0.1%)	0	0	0	3	3	1	0
合計	11,639		5	678	1,986	3,515	3,241	1,764	450

乳房触診所見

所 見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
乳腺症	5	(0.0%)	0	0	2	1	2	0	0
乳腺症の疑い	313	(2.7%)	0	29	87	135	44	13	5
良性乳腺腫瘍	24	(0.2%)	0	3	7	8	6	0	0
良性乳腺腫瘍の疑い	179	(1.6%)	0	18	45	63	36	14	3
乳腺腫瘍	12	(0.1%)	0	0	5	2	2	1	2
乳腺腫瘍の疑い	80	(0.7%)	0	5	15	35	18	4	3
手術後	24	(0.2%)	0	0	0	5	11	3	5
術創あり	45	(0.4%)	0	0	1	12	16	14	2
その他の所見	17	(0.1%)	0	0	5	7	2	3	0
検診者総数	11,414		5	678	1,956	3,487	3,216	1,675	397

マンモグラフィ所見

所 見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
カテゴリー1	9,186	(89.4%)	3	411	1,248	2,935	2,767	1,470	352
カテゴリー2	545	(5.3%)	0	11	87	162	192	78	15
カテゴリー3	512	(5.0%)	1	14	63	193	136	84	21
カテゴリー4	28	(0.3%)	0	0	1	8	11	4	4
カテゴリー5	3	(0.0%)	0	0	0	2	1	0	0
腫瘍	224	(2.2%)	0	5	32	81	62	32	12
石灰化	570	(5.5%)	1	14	87	175	199	77	17
非対称性	303	(2.9%)	0	5	34	108	93	52	11
構築の乱れ	43	(0.4%)	0	1	3	16	12	7	4
その他の所見	15	(0.1%)	0	0	0	6	5	3	1
検診者総数	10,274		4	436	1,399	3,300	3,107	1,636	392

乳房超音波検査所見

所 見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
カテゴリー1	1,204	(69.2%)	1	188	449	224	175	117	50
カテゴリー2	436	(25.0%)	0	51	174	118	49	31	13
カテゴリー3	98	(5.6%)	0	13	34	27	11	6	7
カテゴリー4	3	(0.2%)	0	0	0	2	0	1	0
嚢胞疑い	328	(18.8%)	0	41	108	110	40	22	7
濃縮嚢胞疑い	2	(0.1%)	0	0	0	1	0	0	1
嚢胞内病変疑い	11	(0.6%)	0	1	3	5	1	0	1
充実性病変疑い	63	(3.6%)	0	7	22	15	9	5	5
乳腺内低エコー域	23	(1.3%)	0	3	3	6	5	3	3
乳管拡張（乳輪の範囲を超える）	115	(6.6%)	0	14	68	19	8	6	0
線維腺腫の疑い	12	(0.7%)	0	1	4	3	3	1	0
その他の所見	113	(6.5%)	0	12	46	27	15	6	7
検診者総数	1,741		1	252	657	371	235	155	70

婦人科判定結果

判 定	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	9,301	(83.9%)	12	840	1721	2552	2592	1302	282
放置可	821	(7.4%)	1	73	139	217	223	133	35
要経過観察	444	(4.0%)	0	29	84	185	120	24	2
要精密検査	507	(4.6%)	0	71	116	176	110	29	5
要治療	11	(0.1%)	0	0	4	3	3	0	1
治療中	0	(0.0%)	0	0	0	0	0	0	0
合計	11,084		13	1013	2,064	3,133	3,048	1,488	325

子宮頸部細胞診

所 見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
N I L M（異常なし）	10,679	(96.4%)	13	944	1,958	3,013	2,967	1,464	320
A S C－U S	235	(2.1%)	0	48	65	68	46	7	1
A S C－H	80	(0.7%)	0	5	18	19	23	12	3
L S I L	57	(0.5%)	0	12	15	20	6	4	0
H S I L	28	(0.3%)	0	4	7	11	4	1	1
S C C	1	(0.0%)	0	0	1	0	0	0	0
A G C	2	(0.0%)	0	0	0	1	1	0	0
A d e n o c a .	1	(0.0%)	0	0	0	0	1	0	0
トリコモナス	8	(0.1%)	0	0	2	3	3	0	0
カンジダ	243	(2.2%)	1	58	81	79	22	2	0
その他の所見	9	(0.1%)	0	0	1	6	2	0	0
検診者総数	11,083		13	1013	2,064	3,132	3,048	1,488	325

H P V検査所見

所 見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
陰 性	112	(96.6%)	0	0	90	7	12	3	0
陽 性	4	(3.4%)	0	0	2	2	0	0	0
1 6 型 陰性	115	(99.1%)	0	0	91	9	12	3	0
1 6 型 陽性	1	(0.9%)	0	0	1	0	0	0	0
1 8 型 陰性	115	(99.1%)	0	0	92	8	12	3	0
1 8 型 陽性	1	(0.9%)	0	0	0	1	0	0	0
その他の ハイリスク型陰性	114	(98.3%)	0	0	91	8	12	3	0
その他の ハイリスク型陽性	2	(1.7%)	0	0	1	1	0	0	0
検診者総数	116		0	0	92	9	12	3	0

婦人科診察所見

所 見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
外陰異常	2	(0.0%)	0	0	0	1	1	0	0
外陰異常疑い	1	(0.0%)	0	0	0	0	0	1	0
外陰炎	7	(0.1%)	0	1	1	1	1	3	0
外陰炎疑い	2	(0.0%)	0	0	0	0	1	1	0
膣炎	2	(0.0%)	0	0	0	1	1	0	0
膣炎疑い	3	(0.0%)	0	1	1	0	1	0	0
萎縮性膣炎	33	(0.3%)	0	0	0	0	8	16	9
カンジダ疑い	16	(0.1%)	0	3	6	3	3	1	0
バルトリン腺異常	4	(0.0%)	0	0	1	3	0	0	0
バルトリン腺異常疑い	4	(0.0%)	0	0	2	1	1	0	0
子宮膣部びらん	332	(3.0%)	0	45	110	142	34	1	0
子宮膣部びらん疑い	24	(0.2%)	0	0	13	8	3	0	0
頸管粘膜ポリープ	114	(1.0%)	0	0	9	44	51	10	0
頸管粘膜ポリープ疑い	28	(0.3%)	0	1	1	12	10	3	1
子宮頸部腫瘍疑い	3	(0.0%)	0	0	1	0	2	0	0
子宮体部腫瘍	207	(1.9%)	0	0	15	101	81	10	0
子宮体部腫瘍疑い	72	(0.6%)	0	1	9	40	19	2	1
子宮附属器炎	1	(0.0%)	0	0	1	0	0	0	0
子宮附属器炎疑い	4	(0.0%)	0	0	3	1	0	0	0
良性卵巣腫瘍	1	(0.0%)	0	0	0	1	0	0	0
良性卵巣腫瘍疑い	1	(0.0%)	0	0	1	0	0	0	0
卵巣腫瘍	2	(0.0%)	0	0	1	1	0	0	0
卵巣腫瘍疑い	3	(0.0%)	0	0	2	0	1	0	0
子宮下垂	9	(0.1%)	0	0	0	0	2	5	2
子宮下垂疑い	3	(0.0%)	0	0	0	1	2	0	0
子宮術後	323	(2.9%)	0	0	3	39	146	110	25
その他の所見	12	(0.1%)	0	0	0	8	3	0	1
検診者総数	11,079		13	1013	2,061	3,131	3,048	1,488	325

便潜血判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	18,978	(95.6%)	13	497	3,140	6,142	6,057	2,805	324
	女	12,204	(95.9%)	4	379	2,063	4,052	3,876	1,629	201
	計	31,182	(95.7%)	17	876	5,203	10,194	9,933	4,434	525
放置可	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
要経過観察	男	0	(0.0%)	0	0	0	0	0	0	0
	女	1	(0.0%)	0	0	1	0	0	0	0
	計	1	(0.0%)	0	0	1	0	0	0	0
要精密検査	男	866	(4.4%)	0	11	114	239	299	172	31
	女	517	(4.1%)	1	21	100	191	117	73	14
	計	1,383	(4.2%)	1	32	214	430	416	245	45
要治療	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
治療中	男	16	(0.1%)	0	0	4	6	5	1	0
	女	5	(0.0%)	0	0	0	1	3	1	0
	計	21	(0.1%)	0	0	4	7	8	2	0
合計	男	19,860		13	508	3,258	6,387	6,361	2,978	355
	女	12,727		5	400	2,164	4,244	3,996	1,703	215
	計	32,587		18	908	5,422	10,631	10,357	4,681	570

便潜血の性年齢別陽性率

検査項目	性別	総数			29歳以下			30-39歳			40-49歳			50-59歳			60-69歳			70歳以上		
		受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率
便潜血(一日法)	男	512	19	3.7	11	1	9.1	79	1	1.3	135	4	3.0	154	6	3.9	102	3	2.9	31	4	12.9
	女	563	23	4.1	5	0	—	77	3	3.9	135	4	3.0	190	8	4.2	142	5	3.5	14	3	21.4
	計	1,075	42	3.9	16	1	6.3	156	4	2.6	270	8	3.0	344	14	4.1	244	8	3.3	45	7	15.6
便潜血(二日法)	男	19,348	863	4.5	510	10	2.0	3,179	117	3.7	6,252	241	3.9	6,207	298	4.8	2,876	170	5.9	324	27	8.3
	女	12,165	500	4.1	400	22	5.5	2,088	98	4.7	4,109	188	4.6	3,806	112	2.9	1,561	69	4.4	201	11	5.5
	計	31,513	1,363	4.3	910	32	3.5	5,267	215	4.1	10,361	429	4.1	10,013	410	4.1	4,437	239	5.4	525	38	7.2

陽性率%

脂質検査判定結果

平成26年4月1日より、判定基準を変更しています。

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	9,332	(33.4%)	186	1,781	2,332	2,059	1,760	1,053	161
	女	10,446	(47.6%)	172	2,071	2,883	3,219	1,482	502	117
	計	19,778	(39.6%)	358	3,852	5,215	5,278	3,242	1,555	278
放置可	男	6,083	(21.7%)	26	563	1,267	1,671	1,581	862	113
	女	4,240	(19.3%)	30	381	748	1,211	1,277	505	88
	計	10,323	(20.7%)	56	944	2,015	2,882	2,858	1,367	201
要経過観察	男	8,418	(30.1%)	75	551	1,769	2,517	2,306	1,080	120
	女	4,233	(19.3%)	33	212	479	1,083	1,632	687	107
	計	12,651	(25.3%)	108	763	2,248	3,600	3,938	1,767	227
要精密検査	男	1,660	(5.9%)	19	227	361	481	410	149	13
	女	1,507	(6.9%)	9	332	353	337	349	108	19
	計	3,167	(6.3%)	28	559	714	818	759	257	32
要治療	男	409	(1.5%)	0	21	93	140	115	38	2
	女	184	(0.8%)	0	3	16	36	85	42	2
	計	593	(1.2%)	0	24	109	176	200	80	4
治療中	男	2,078	(7.4%)	0	10	77	440	915	556	80
	女	1,349	(6.1%)	0	2	21	125	550	550	101
	計	3,427	(6.9%)	0	12	98	565	1,465	1,106	181
合計	男	27,980		306	3,153	5,899	7,308	7,087	3,738	489
	女	21,959		244	3,001	4,500	6,011	5,375	2,394	434
	計	49,939		550	6,154	10,399	13,319	12,462	6,132	923

脂質検査の性年齢別平均値

検査項目	総数			29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
総コレステロール mg/dL	男	21,459	198.1	1,941	176.3	4,265	193.7	5,993	202.4	5,861	203.9	2,985	199.7	414	192.8
	女	14,571	199.3	1,731	178.2	2,959	184.4	4,156	195.8	3,716	217.2	1,685	215.0	324	203.5
	計	36,030	198.6	3,672	177.2	7,224	189.9	10,149	199.7	9,577	209.0	4,670	205.2	738	197.5
中性脂肪 mg/dL	男	27,812	117.4	3,398	82.4	5,817	113.2	7,291	126.8	7,082	127.8	3,735	118.5	489	108.2
	女	21,905	75.7	3,209	57.4	4,483	65.2	6,010	72.5	5,375	88.1	2,394	95.5	434	99.2
	計	49,717	99.0	6,607	70.3	10,300	92.3	13,301	102.3	12,457	110.7	6,129	109.5	923	104.0
HDLコレステロール mg/dL	男	27,952	55.8	3,458	56.2	5,893	54.9	7,294	55.2	7,082	56.8	3,736	56.4	489	55.3
	女	21,959	66.7	3,245	66.2	4,500	66.1	6,011	66.8	5,375	68.5	2,394	65.3	434	59.7
	計	49,911	60.6	6,703	61.1	10,393	59.8	13,305	60.5	12,457	61.9	6,130	59.9	923	57.3
LDLコレステロール mg/dL	男	27,907	120.9	3,435	106.5	5,880	120.0	7,290	126.0	7,081	124.1	3,732	120.4	489	114.4
	女	21,803	114.5	3,180	97.7	4,445	103.4	5,997	113.0	5,359	128.8	2,388	128.5	434	118.4
	計	49,710	118.1	6,615	102.3	10,325	112.9	13,287	120.1	12,440	126.2	6,120	123.5	923	116.3

糖尿病判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
	男	女	計							
異常なし	21,464	19,655	41,119	368	3,847	5,761	5,506	4,016	1,765	201
	(70.3%)	(82.6%)	(75.7%)							
	325	4,676	5,001	3,662	10,437	10,824	8,058	3,202	396	
放置可	5,144	2,972	8,116	11	175	661	1,402	1,744	1,014	137
	(16.8%)	(12.5%)	(14.9%)							
	8	93	101	268	917	2,131	2,814	1,664	303	
要経過観察	1,682	566	2,248	0	15	138	324	690	448	67
	(5.5%)	(2.4%)	(4.1%)							
	0	3	3	18	165	423	905	629	108	
要精密検査	325	100	425	0	5	22	54	144	89	11
	(1.1%)	(0.4%)	(0.8%)							
	0	1	1	6	25	73	187	116	18	
要治療	326	73	399	0	3	32	87	120	76	8
	(1.1%)	(0.3%)	(0.7%)							
	0	1	1	4	40	101	151	92	11	
治療中	1,597	436	2,033	0	6	61	258	652	538	82
	(5.2%)	(1.8%)	(3.7%)							
	0	3	3	9	75	312	818	705	114	
合計	30,538	23,802	54,340	379	4,051	6,675	7,631	7,366	3,930	506
	333	4,984	5,317	7,814	11,659	13,864	12,933	6,408	950	

糖尿病検査の性年齢別平均値

検査項目	性別	総数			29歳以下			30-39歳			40-49歳			50-59歳			60-69歳			70歳以上		
		人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差
血糖 mg/dL	男	27,897	97.8	3,376	89.3	5,901	92.7	7,311	97.1	7,087	102.4	3,733	105.6	489	105.6							
	女	21,847	91.7	3,167	86.9	4,475	88.3	6,008	91.0	5,370	94.4	2,393	98.1	434	102.4							
	計	49,744	95.1	6,543	88.1	10,376	90.8	13,319	94.3	12,457	98.9	6,126	102.7	923	104.1							
HbA1c (NGSP) %	男	13,595	5.4	1,309	5.1	2,311	5.2	3,940	5.4	4,214	5.6	1,599	5.7	222	5.8							
	女	7,850	5.4	684	5.1	1,449	5.2	2,498	5.3	2,084	5.5	941	5.7	194	5.7							
	計	21,445	5.4	1,993	5.1	3,760	5.2	6,438	5.4	6,298	5.5	2,540	5.7	416	5.8							

糖負荷試験の判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
	男	女	計							
正常型	125	61	186	0	0	3	48	55	14	5
	(62.8%)	(84.7%)	(68.6%)							
	0	0	0	0	0	5	66	85	21	9
境界型	64	10	74	0	0	1	8	34	18	3
	(32.2%)	(13.9%)	(27.3%)							
	0	0	0	0	0	1	9	38	21	5
糖尿病型	10	1	11	0	0	0	1	5	1	3
	(5.0%)	(1.4%)	(4.1%)							
	0	0	0	0	0	0	1	5	2	3
合計	199	72	271	0	0	4	57	94	33	11
	0	0	0	0	0	6	76	128	44	17

尿糖の性年齢別陽性率

検査項目	性別	総数			29歳以下			30-39歳			40-49歳			50-59歳			60-69歳			70歳以上		
		受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率
尿糖定性	男	31,088	631	2.0	4,632	11	0.2	6,765	65	1.0	7,737	155	2.0	7,478	230	3.1	3,971	150	3.8	505	20	4.0
	女	23,840	104	0.4	4,124	7	0.2	5,015	12	0.2	6,233	23	0.4	5,556	32	0.6	2,472	29	1.2	440	1	0.2
	計	54,928	735	1.3	8,756	18	0.2	11,780	77	0.7	13,970	178	1.3	13,034	262	2.0	6,443	179	2.8	945	21	2.2

陽性率%

痛風判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	18,270	(75.3%)	185	1,676	3,706	4,947	4,810	2,616	330
	女	15,727	(94.1%)	191	1,790	3,323	4,569	3,838	1,735	281
	計	33,997	(83.0%)	376	3,466	7,029	9,516	8,648	4,351	611
放置可	男	1,795	(7.4%)	19	179	392	539	438	208	20
	女	459	(2.7%)	3	30	68	92	172	75	19
	計	2,254	(5.5%)	22	209	460	631	610	283	39
要経過観察	男	2,191	(9.0%)	23	192	469	661	583	239	24
	女	407	(2.4%)	6	27	50	90	142	66	26
	計	2,598	(6.3%)	29	219	519	751	725	305	50
要精密検査	男	537	(2.2%)	5	49	123	181	129	39	11
	女	82	(0.5%)	0	4	8	17	32	16	5
	計	619	(1.5%)	5	53	131	198	161	55	16
要治療	男	45	(0.2%)	1	8	8	16	7	5	0
	女	11	(0.1%)	0	1	0	1	5	1	3
	計	56	(0.1%)	1	9	8	17	12	6	3
治療中	男	1,419	(5.8%)	0	10	108	340	626	293	42
	女	30	(0.2%)	0	0	1	5	8	10	6
	計	1,449	(3.5%)	0	10	109	345	634	303	48
合 計	男	24,257		233	2,114	4,806	6,684	6,593	3,400	427
	女	16,716		200	1,852	3,450	4,774	4,197	1,903	340
	計	40,973		433	3,966	8,256	11,458	10,790	5,303	767

尿酸の性年齢別平均値

検査項目	総数			29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
尿酸・ mg / dL	男	24,257	6.09	2,347	6.13	4,806	6.16	6,684	6.15	6,593	6.06	3,400	5.89	427	5.92
	女	16,716	4.41	2,052	4.31	3,450	4.21	4,774	4.25	4,197	4.63	1,903	4.68	340	4.87
	計	40,973	5.40	4,399	5.28	8,256	5.34	11,458	5.36	10,790	5.50	5,303	5.46	767	5.45

肝機能・血液蛋白検査判定結果平成 26 年 4 月 1 日より、判定基準を変更しています。

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	14,049	(49.0%)	125	2,067	3,132	3,363	3,154	1,960	248
	女	17,335	(77.5%)	272	2,692	3,826	4,867	3,835	1,638	205
	計	31,384	(61.5%)	397	4,759	6,958	8,230	6,989	3,598	453
放置可	男	6,974	(24.3%)	61	710	1,442	1,859	1,901	880	121
	女	3,009	(13.5%)	38	306	504	781	887	416	77
	計	9,983	(19.6%)	99	1016	1,946	2,640	2,788	1,296	198
要経過観察	男	3,338	(11.7%)	120	314	612	866	883	473	70
	女	1,217	(5.4%)	27	80	177	250	375	208	100
	計	4,555	(8.9%)	147	394	789	1,116	1,258	681	170
要精密検査	男	3,916	(13.7%)	17	299	881	1221	1049	404	45
	女	698	(3.1%)	0	34	75	161	271	118	39
	計	4,614	(9.0%)	17	333	956	1382	1320	522	84
要治療	男	30	(0.1%)	1	5	11	5	7	1	0
	女	5	(0.0%)	0	1	0	1	3	0	0
	計	35	(0.1%)	1	6	11	6	10	1	0
治療中	男	341	(1.2%)	0	5	28	78	168	56	6
	女	93	(0.4%)	0	3	12	12	35	17	14
	計	434	(0.9%)	0	8	40	90	203	73	20
合計	男	28,648		324	3,400	6,106	7,392	7,162	3,774	490
	女	22,357		337	3,116	4,594	6,072	5,406	2,397	435
	計	51,005		661	6,516	10,700	13,464	12,568	6,171	925

肝炎ウイルス検査の性年齢別陽性率

検査項目	性別	総数			29歳以下			30-39歳			40-49歳			50-59歳			60-69歳			70歳以上		
		受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率
HBs抗原	男	3,495	27	0.8	262	0	—	634	1	0.2	956	5	0.5	1,089	11	1.0	448	8	1.8	106	2	1.9
	女	2,840	15	0.5	391	0	—	548	3	0.5	712	1	0.1	728	8	1.1	380	2	0.5	81	1	1.2
	計	6,335	42	0.7	653	0	—	1,182	4	0.3	1,668	6	0.4	1,817	19	1.0	828	10	1.2	187	3	1.6
HCV抗体	男	1,567	5	0.3	156	0	—	304	0	—	424	1	0.2	484	3	0.6	187	0	—	12	1	8.3
	女	1,270	3	0.2	181	0	—	297	0	—	318	0	—	367	1	0.3	100	2	2.0	7	0	—
	計	2,837	8	0.3	337	0	—	601	0	—	742	1	0.1	851	4	0.5	287	2	0.7	19	1	5.3

陽性率%

肝機能・血液蛋白検査の性年齢別平均値

検査項目	総数			29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
TTT マクラガン	男	4,394	1.44	161	1.40	768	1.39	1,411	1.44	1,412	1.44	526	1.47	116	1.54
	女	2,438	1.61	131	1.63	435	1.76	765	1.65	720	1.53	294	1.39	93	1.84
	計	6,832	1.50	292	1.50	1,203	1.53	2,176	1.52	2,132	1.47	820	1.44	209	1.67
ZTT クンケル	男	5,697	6.93	346	7.08	965	6.90	1,690	6.70	1,707	6.74	790	7.55	199	7.85
	女	3,160	7.91	163	8.20	518	8.50	915	8.03	880	7.50	516	7.57	168	8.40
	計	8,857	7.28	509	7.44	1,483	7.46	2,605	7.17	2,587	7.00	1,306	7.56	367	8.10
総ビリルビン mg／dL	男	14,126	1.01	1,905	1.04	2,864	1.01	3,758	1.00	3,947	0.99	1,428	0.99	224	1.00
	女	8,993	0.89	1,504	0.87	1,932	0.86	2,531	0.88	2,028	0.92	812	0.89	186	0.89
	計	23,119	0.96	3,409	0.97	4,796	0.95	6,289	0.95	5,975	0.97	2,240	0.96	410	0.95
AST (GOT) U／L	男	28,092	24.4	3,530	23.5	5,975	24.0	7,298	24.4	7,069	25.1	3,732	24.7	488	25.1
	女	21,937	19.9	3,271	17.7	4,501	18.1	5,982	18.9	5,360	22.0	2,389	23.2	434	24.2
	計	50,029	22.4	6,801	20.7	10,476	21.5	13,280	21.9	12,429	23.8	6,121	24.1	922	24.7
ALT (GPT) U／L	男	28,092	28.1	3,530	26.2	5,975	30.5	7,298	29.7	7,069	27.7	3,732	24.5	488	22.4
	女	21,937	17.3	3,271	13.9	4,501	15.3	5,982	16.6	5,360	20.2	2,389	20.6	434	19.1
	計	50,029	23.4	6,801	20.3	10,476	24.0	13,280	23.8	12,429	24.5	6,121	23.0	922	20.9
アルフォス U／L	男	23,413	217.8	2,185	230.5	4,593	213.3	6,546	216.8	6,452	215.7	3,225	221.6	412	220.9
	女	15,764	195.5	1,758	175.1	3,126	172.1	4,606	173.5	4,135	223.4	1,768	236.4	371	259.1
	計	39,177	208.8	3,943	205.8	7,719	196.6	11,152	198.9	10,587	218.7	4,993	226.8	783	239.0
コリンエステラーゼ U／L	男	4,794	362.4	495	347.5	886	368.4	1,438	372.9	1,434	359.0	499	349.1	42	327.4
	女	3,047	317.5	352	287.9	615	294.8	992	305.3	810	348.7	258	361.5	20	306.8
	計	7,841	344.9	847	322.7	1,501	338.2	2,430	345.3	2,244	355.3	757	353.3	62	320.7
LDH U／L	男	13,234	180.2	1,624	182.4	2,575	177.9	3,638	178.4	3,836	180.6	1,350	184.5	211	189.2
	女	7,305	173.5	1,028	163.7	1,542	161.0	2,084	167.2	1,711	184.5	763	197.5	177	205.6
	計	20,539	177.8	2,652	175.1	4,117	171.5	5,722	174.3	5,547	181.8	2,113	189.2	388	196.7
γ-GT (γ-GTP) U／L	男	28,061	45.9	3,517	27.3	5,959	40.2	7,297	50.1	7,069	55.2	3,732	47.1	487	41.2
	女	21,779	22.1	3,196	15.5	4,440	18.1	5,962	21.5	5,359	27.4	2,389	27.3	433	24.9
	計	49,840	35.5	6,713	21.7	10,399	30.8	13,259	37.3	12,428	43.2	6,121	39.4	920	33.5
LAP U／L	男	3,153	61.0	30	62.2	552	61.2	924	61.8	1,149	60.8	400	60.1	98	58.1
	女	2,014	53.9	29	53.2	341	52.8	577	49.8	713	56.2	264	57.6	90	56.4
	計	5,167	58.2	59	57.8	893	58.0	1,501	57.2	1,862	59.0	664	59.1	188	57.3
総蛋白 g／dL	男	15,543	7.27	2,173	7.40	3,121	7.32	4,147	7.24	4,319	7.22	1,539	7.23	244	7.18
	女	10,488	7.21	1,764	7.35	2,166	7.24	2,919	7.14	2,443	7.21	891	7.21	305	6.98
	計	26,031	7.25	3,937	7.38	5,287	7.29	7,066	7.20	6,762	7.21	2,430	7.22	549	7.07
アルブミン g／dL	男	14,561	4.58	1,991	4.81	2,929	4.69	3,886	4.56	4,101	4.48	1,411	4.41	243	4.21
	女	10,101	4.39	1,744	4.56	2,088	4.42	2,809	4.32	2,330	4.37	827	4.35	303	3.92
	計	24,662	4.50	3,735	4.70	5,017	4.58	6,695	4.46	6,431	4.44	2,238	4.39	546	4.05
A／G比	男	5,044	1.72	621	1.87	957	1.79	1,384	1.73	1,461	1.66	503	1.56	118	1.49
	女	4,499	1.58	787	1.64	823	1.59	1,279	1.55	1,153	1.56	358	1.55	99	1.54
	計	9,543	1.65	1,408	1.74	1,780	1.70	2,663	1.65	2,614	1.62	861	1.55	217	1.51
アルブミン %	男	24	65.3	0	—	2	65.7	11	65.2	9	65.3	2	66.2	0	—
	女	3	61.6	0	—	1	57.3	0	—	1	66.9	1	60.5	0	—
	計	27	64.9	0	—	3	62.9	11	65.2	10	65.4	3	64.3	0	—
α1-グロブリン %	男	24	2.7	0	—	2	2.5	11	2.7	9	2.7	2	2.6	0	—
	女	3	2.9	0	—	1	3.6	0	—	1	2.7	1	2.3	0	—
	計	27	2.7	0	—	3	2.9	11	2.7	10	2.7	3	2.5	0	—
α2-グロブリン %	男	24	7.5	0	—	2	7.4	11	7.7	9	7.1	2	8.2	0	—
	女	3	8.0	0	—	1	10.0	0	—	1	7.7	1	6.4	0	—
	計	27	7.5	0	—	3	8.2	11	7.7	10	7.2	3	7.6	0	—
β-グロブリン %	男	24	9.5	0	—	2	9.0	11	9.8	9	9.4	2	9.1	0	—
	女	3	10.7	0	—	1	12.8	0	—	1	10.5	1	8.8	0	—
	計	27	9.7	0	—	3	10.3	11	9.8	10	9.6	3	9.0	0	—
γ-グロブリン %	男	24	15.0	0	—	2	15.5	11	14.7	9	15.5	2	14.0	0	—
	女	3	16.8	0	—	1	16.3	0	—	1	12.2	1	22.0	0	—
	計	27	15.2	0	—	3	15.7	11	14.7	10	15.1	3	16.7	0	—
αフェトпротеイン定量 ng／mL	男	1,003	3.3	3	3.2	176	2.7	252	3.4	344	3.5	208	3.6	20	2.7
	女	361	3.9	3	1.6	51	4.5	95	3.5	138	3.9	60	4.4	14	3.2
	計	1,364	3.5	6	2.4	227	3.1	347	3.4	482	3.6	268	3.8	34	2.9

貧血・血球数判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	20,284	(73.1%)	218	2,581	4,532	5,241	4,881	2,538	293
	女	15,593	(71.4%)	184	2,424	3,187	3,899	3,836	1,780	283
	計	35,877	(72.3%)	402	5,005	7,719	9,140	8,717	4,318	576
放置可	男	4,686	(16.9%)	54	408	903	1,312	1,300	636	73
	女	3,470	(15.9%)	34	396	721	913	939	405	62
	計	8,156	(16.4%)	88	804	1,624	2,225	2,239	1,041	135
要経過観察	男	2,497	(9.0%)	23	125	350	655	773	475	96
	女	2,022	(9.3%)	16	241	505	679	373	150	58
	計	4,519	(9.1%)	39	366	855	1,334	1,146	625	154
要精密検査	男	199	(0.7%)	1	3	27	40	56	55	17
	女	431	(2.0%)	3	26	99	229	51	3	20
	計	630	(1.3%)	4	29	126	269	107	58	37
要治療	男	23	(0.1%)	0	1	1	1	9	7	4
	女	63	(0.3%)	0	4	10	39	9	0	1
	計	86	(0.2%)	0	5	11	40	18	7	5
治療中	男	58	(0.2%)	3	1	3	10	26	12	3
	女	263	(1.2%)	7	10	29	128	71	11	7
	計	321	(0.6%)	10	11	32	138	97	23	10
合 計	男	27,747		299	3,119	5,816	7,259	7,045	3,723	486
	女	21,842		244	3,101	4,551	5,887	5,279	2,349	431
	計	49,589		543	6,220	10,367	13,146	12,324	6,072	917

貧血・血球数検査の性年齢別平均値

検査項目		総数		29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
赤血球数 万 / μ L	男	27,828	492.8	3,435	512.3	5,835	505.0	7,278	495.3	7,059	483.1	3,735	474.5	486	456.2
	女	21,851	437.7	3,349	442.0	4,551	436.8	5,891	434.6	5,280	440.8	2,349	438.3	431	415.7
	計	49,679	468.6	6,784	477.6	10,386	475.1	13,169	468.1	12,339	465.0	6,084	460.5	917	437.2
ヘモグロビン g / dL	男	27,829	15.35	3,435	15.60	5,836	15.52	7,278	15.40	7,059	15.25	3,735	15.05	486	14.51
	女	21,853	13.08	3,350	13.13	4,552	12.92	5,890	12.80	5,280	13.35	2,350	13.46	431	12.89
	計	49,682	14.35	6,785	14.38	10,388	14.38	13,168	14.24	12,339	14.44	6,085	14.43	917	13.75
ヘマトクリット %	男	27,014	45.04	3,204	45.64	5,635	45.49	7,083	45.22	6,926	44.77	3,685	44.32	481	42.90
	女	21,296	39.29	3,153	39.31	4,420	38.85	5,777	38.61	5,195	39.99	2,321	40.31	430	38.74
	計	48,310	42.51	6,357	42.50	10,055	42.57	12,860	42.25	12,121	42.72	6,006	42.77	911	40.94
M C V fL	男	3,534	91.70	104	89.30	633	89.91	1,052	91.18	1,241	92.53	427	93.33	77	94.28
	女	2,688	89.94	122	89.62	527	89.04	821	88.96	896	90.65	276	92.13	46	91.74
	計	6,222	90.94	226	89.47	1,160	89.52	1,873	90.21	2,137	91.74	703	92.86	123	93.33
M C H pg	男	3,533	31.26	104	30.57	632	30.74	1,052	31.08	1,241	31.57	427	31.67	77	31.81
	女	2,687	29.95	121	30.14	527	29.68	821	29.47	896	30.24	276	30.74	46	30.42
	計	6,220	30.69	225	30.34	1,159	30.26	1,873	30.37	2,137	31.01	703	31.30	123	31.29
M C H C %	男	3,521	34.09	104	34.23	626	34.20	1,050	34.08	1,237	34.12	427	33.93	77	33.75
	女	2,673	33.26	120	33.61	525	33.30	813	33.07	894	33.33	276	33.37	45	33.12
	計	6,194	33.73	224	33.90	1,151	33.79	1,863	33.64	2,131	33.79	703	33.71	122	33.52
血清鉄 μ g / dL	男	3,712	118.6	326	111.6	628	121.9	1,020	119.9	1,249	118.9	421	116.7	68	108.3
	女	2,473	103.0	156	102.1	443	102.8	764	97.7	845	107.6	213	105.1	52	100.4
	計	6,185	112.4	482	108.6	1,071	114.0	1,784	110.4	2,094	114.4	634	112.8	120	104.9
白血球数 千 / μ L	男	24,791	6.29	2,449	6.08	4,841	6.34	6,844	6.43	6,782	6.25	3,437	6.18	438	6.03
	女	18,041	5.79	2,293	6.38	3,699	6.11	5,136	5.81	4,549	5.35	1,966	5.48	398	5.80
	計	42,832	6.08	4,742	6.23	8,540	6.24	11,980	6.16	11,331	5.89	5,403	5.93	836	5.92
血小板数 万 / μ L	男	7,813	23.99	598	24.39	1,372	24.78	2,156	24.79	2,324	23.70	1,138	22.58	225	20.60
	女	6,013	24.90	419	25.33	1,108	25.48	1,696	26.14	1,683	24.54	806	23.27	301	21.63
	計	13,826	24.39	1,017	24.78	2,480	25.09	3,852	25.38	4,007	24.05	1,944	22.87	526	21.19

炎症反応判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	3,095	(91.2%)	0	21	499	898	1087	467	123
	女	2,095	(92.1%)	0	34	364	555	679	371	92
	計	5,190	(91.6%)	0	55	863	1,453	1,766	838	215
放置可	男	0	(0.0%)	0	0	0	0	0	0	0
	女	1	(0.0%)	0	0	0	0	0	1	0
	計	1	(0.0%)	0	0	0	0	0	1	0
要経過観察	男	298	(8.8%)	0	2	41	87	106	46	16
	女	178	(7.8%)	0	1	36	45	56	27	13
	計	476	(8.4%)	0	3	77	132	162	73	29
要精密検査	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
要治療	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
治療中	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
合 計	男	3,393		0	23	540	985	1,193	513	139
	女	2,274		0	35	400	600	735	399	105
	計	5,667		0	58	940	1,585	1,928	912	244

炎症反応検査の性年齢別陽性率

検査項目	性 別	総数			29歳以下			30-39歳			40-49歳			50-59歳			60-69歳			70歳以上		
		受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率
C R P	男	3,363	295	8.8	10	1	10.0	538	41	7.6	981	86	8.8	1,189	106	8.9	506	45	8.9	139	16	11.5
	女	2,221	173	7.8	23	0	—	390	35	9.0	591	43	7.3	721	55	7.6	391	27	6.9	105	13	12.4
	計	5,584	468	8.4	33	1	3.0	928	76	8.2	1,572	129	8.2	1,910	161	8.4	897	72	8.0	244	29	11.9
血 沈	男	18	0	—	0	—	—	1	0	—	6	0	0.0	10	0	—	0	—	—	1	0	—
	女	10	1	10.0	0	—	—	2	0	—	2	1	50.0	5	0	—	1	0	—	0	—	—
	計	28	1	3.6	0	—	—	3	0	—	8	1	12.5	15	0	—	1	0	—	1	0	—

陽性率%

腎機能判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	23,397	(76.9%)	355	3,571	5,608	5,862	5,172	2,573	256
	女	16,146	(68.1%)	261	3,069	3,672	3,897	3,536	1,522	189
	計	39,543	(73.0%)	616	6,640	9,280	9,759	8,708	4,095	445
放置可	男	2,324	(7.6%)	4	264	422	593	606	362	73
	女	2,747	(11.6%)	24	389	559	917	537	250	71
	計	5,071	(9.4%)	28	653	981	1,510	1,143	612	144
要経過観察	男	4,073	(13.4%)	14	184	547	994	1,357	833	144
	女	4,582	(19.3%)	46	272	703	1,332	1,421	658	150
	計	8,655	(16.0%)	60	456	1,250	2,326	2,778	1,491	294
要精密検査	男	368	(1.2%)	6	20	42	92	113	82	13
	女	139	(0.6%)	1	20	23	23	27	22	23
	計	507	(0.9%)	7	40	65	115	140	104	36
要治療	男	2	(0.0%)	0	0	0	2	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	2	(0.0%)	0	0	0	2	0	0	0
治療中	男	261	(0.9%)	0	4	27	55	81	76	18
	女	106	(0.4%)	1	7	7	28	31	20	12
	計	367	(0.7%)	1	11	34	83	112	96	30
合計	男	30,425		379	4,043	6,646	7,598	7,329	3,926	504
	女	23,720		333	3,757	4,964	6,197	5,552	2,472	445
	計	54,145		712	7,800	11,610	13,795	12,881	6,398	949

腎機能検査の性年齢別平均値

検査項目	総数			29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
尿素窒素 mg / dL	男	23,133	13.95	2,200	13.29	4,550	13.14	6,441	13.52	6,356	14.36	3,174	15.31	412	16.64
	女	15,616	12.46	1,810	11.28	3,084	11.16	4,553	11.74	4,070	13.52	1,723	14.49	376	16.64
	計	38,749	13.35	4,010	12.38	7,634	12.34	10,994	12.78	10,426	14.03	4,897	15.02	788	16.64
クレアチニン mg / dL	男	24,327	0.88	2,289	0.86	4,775	0.86	6,786	0.88	6,704	0.89	3,347	0.90	426	0.93
	女	16,848	0.65	1,919	0.63	3,395	0.62	4,942	0.64	4,357	0.66	1,854	0.66	381	0.70
	計	41,175	0.79	4,208	0.76	8,170	0.76	11,728	0.78	11,061	0.80	5,201	0.82	807	0.82

尿検査の性年齢別陽性率

検査項目	性別	総数			29歳以下			30-39歳			40-49歳			50-59歳			60-69歳			70歳以上		
		受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率
尿蛋白定性	男	31,401	1,523	4.9	4,695	166	3.5	6,834	214	3.1	7,815	335	4.3	7,543	479	6.4	4,009	290	7.2	505	39	7.7
	女	23,907	750	3.1	4,140	182	4.4	5,035	160	3.2	6,244	197	3.2	5,564	125	2.2	2,484	57	2.3	440	29	6.6
	計	55,308	2,273	4.1	8,835	348	3.9	11,869	374	3.2	14,059	532	3.8	13,107	604	4.6	6,493	347	5.3	945	68	7.2
尿潜血	男	24,904	1,667	6.7	2,547	60	2.4	4,885	220	4.5	6,844	405	5.9	6,795	605	8.9	3,401	329	9.7	432	48	11.1
	女	17,913	4,069	22.7	2,336	425	18.2	3,686	788	21.4	5,088	1,297	25.5	4,485	1,008	22.5	1,931	461	23.9	387	90	23.3
	計	42,817	5,736	13.4	4,883	485	9.9	8,571	1,008	11.8	11,932	1,702	14.3	11,280	1,613	14.3	5,332	790	14.8	819	138	16.8

(+) 以上の人を陽性として判定している 陽性率%

膵機能判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	12,741	(92.6%)	55	1,430	2,559	3,580	3,697	1,251	169
	女	6,458	(92.2%)	30	833	1,405	1,905	1,496	633	156
	計	19,199	(92.5%)	85	2,263	3,964	5,485	5,193	1,884	325
放置可	男	373	(2.7%)	3	29	81	127	114	19	0
	女	102	(1.5%)	0	11	14	36	28	11	2
	計	475	(2.3%)	3	40	95	163	142	30	2
要経過観察	男	608	(4.4%)	3	41	61	129	200	137	37
	女	433	(6.2%)	1	50	82	95	115	71	19
	計	1,041	(5.0%)	4	91	143	224	315	208	56
要精密検査	男	25	(0.2%)	0	1	5	5	11	3	0
	女	11	(0.2%)	0	1	3	1	4	2	0
	計	36	(0.2%)	0	2	8	6	15	5	0
要治療	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
治療中	男	10	(0.1%)	0	0	0	0	6	3	1
	女	2	(0.0%)	0	0	0	0	2	0	0
	計	12	(0.1%)	0	0	0	0	8	3	1
合 計	男	13,757		61	1,501	2,706	3,841	4,028	1,413	207
	女	7,006		31	895	1,504	2,037	1,645	717	177
	計	20,763		92	2,396	4,210	5,878	5,673	2,130	384

膵機能検査の性年齢別平均値

検査項目	総数			29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
血清アミラーゼ IU/L	男	13,758	72.7	1,562	70.7	2,707	69.5	3,841	70.0	4,028	74.1	1,413	81.3	207	91.2
	女	7,006	78.0	926	77.5	1,504	78.1	2,037	75.4	1,645	78.8	717	82.7	177	82.6
	計	20,764	74.5	2,488	73.2	4,211	72.6	5,878	71.9	5,673	75.5	2,130	81.8	384	87.2

電解質判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	2,193	(85.7%)	137	58	349	597	664	301	87
	女	1,228	(81.3%)	78	31	209	299	308	183	120
	計	3,421	(84.1%)	215	89	558	896	972	484	207
放置可	男	280	(10.9%)	5	17	41	75	94	39	9
	女	216	(14.3%)	10	3	28	47	83	28	17
	計	496	(12.2%)	15	20	69	122	177	67	26
要経過観察	男	85	(3.3%)	3	3	6	18	34	18	3
	女	65	(4.3%)	1	0	7	13	21	10	13
	計	150	(3.7%)	4	3	13	31	55	28	16
要精密検査	男	1	(0.0%)	0	0	0	0	0	0	1
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	1	(0.0%)	0	0	0	0	0	0	1
要治療	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
治療中	男	0	(0.0%)	0	0	0	0	0	0	0
	女	1	(0.1%)	0	0	0	0	1	0	0
	計	1	(0.0%)	0	0	0	0	1	0	0
合 計	男	2,559		145	78	396	690	792	358	100
	女	1,510		89	34	244	359	413	221	150
	計	4,069		234	112	640	1,049	1,205	579	250

電解質検査の性年齢別平均値

検査項目	総数			29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
ナトリウム・ mEq/L	男	2,559	140.8	223	141.0	397	140.9	690	140.9	792	140.8	357	140.8	100	140.2
	女	1,509	140.3	123	140.3	244	139.2	359	139.5	413	141.0	220	141.4	150	140.0
	計	4,068	140.6	346	140.8	641	140.3	1,049	140.4	1,205	140.9	577	141.0	250	140.1
カリウム・ mEq/L	男	2,560	4.13	223	4.23	397	4.05	690	4.09	792	4.11	358	4.22	100	4.24
	女	1,510	4.08	123	4.20	244	3.98	359	4.03	413	4.07	221	4.14	150	4.19
	計	4,070	4.11	346	4.22	641	4.02	1,049	4.07	1,205	4.10	579	4.19	250	4.21
クロール・ mEq/L	男	2,559	105.8	223	104.8	397	105.8	690	106.0	792	106.2	357	105.8	100	104.9
	女	1,509	106.0	123	105.5	244	106.1	359	106.3	413	106.3	220	106.1	150	104.7
	計	4,068	105.9	346	105.1	641	105.9	1,049	106.1	1,205	106.2	577	105.9	250	104.8
無機リン・ mg/dL	男	2,254	3.24	80	3.61	392	3.35	664	3.26	768	3.17	299	3.15	51	3.08
	女	1,174	3.59	29	3.61	241	3.55	354	3.45	382	3.74	136	3.64	32	3.59
	計	3,428	3.36	109	3.61	633	3.43	1,018	3.33	1,150	3.36	435	3.30	83	3.28
カルシウム・ mg/dL	男	2,261	9.43	80	9.74	392	9.48	666	9.43	773	9.39	299	9.38	51	9.32
	女	1,183	9.33	29	9.41	241	9.24	354	9.21	388	9.45	139	9.42	32	9.40
	計	3,444	9.39	109	9.65	633	9.39	1,020	9.35	1,161	9.41	438	9.39	83	9.35

その他の血液検査の判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	4,461	(91.8%)	47	378	743	1062	1,280	776	175
	女	3,816	(94.5%)	121	521	716	847	956	535	120
	計	8,277	(93.0%)	168	899	1,459	1,909	2,236	1,311	295
放置可	男	110	(2.3%)	33	11	9	18	23	13	3
	女	93	(2.3%)	22	10	9	19	22	9	2
	計	203	(2.3%)	55	21	18	37	45	22	5
要経過観察	男	197	(4.1%)	82	10	26	23	30	21	5
	女	73	(1.8%)	21	3	6	13	20	5	5
	計	270	(3.0%)	103	13	32	36	50	26	10
要精密検査	男	90	(1.9%)	0	1	2	14	47	23	3
	女	53	(1.3%)	0	6	6	13	19	7	2
	計	143	(1.6%)	0	7	8	27	66	30	5
要治療	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
治療中	男	2	(0.0%)	0	0	0	0	0	1	1
	女	2	(0.0%)	0	0	2	0	0	0	0
	計	4	(0.0%)	0	0	2	0	0	1	1
合計	男	4,860		162	400	780	1,117	1,380	834	187
	女	4,037		164	540	739	892	1017	556	129
	計	8,897		326	940	1,519	2,009	2,397	1,390	316

その他の血液検査の性年齢別陽性率

検査項目	性別	総数			29歳以下			30-39歳			40-49歳			50-59歳			60-69歳			70歳以上		
		受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率
CEA	男	1,277	52	4	3	0	—	81	2	3	390	9	2	536	27	5	238	14	6	29	0	—
	女	789	5	1	6	0	—	102	0	—	238	0	—	343	3	1	85	1	1	15	1	7
	計	2,066	57	3	9	0	—	183	2	1	628	9	1	879	30	3	323	15	5	44	1	2
RF	男	2,592	133	5	353	3	1	518	6	1	564	26	5	662	58	9	414	31	8	81	9	11
	女	2,436	132	5	587	16	3	559	17	3	517	33	6	505	48	10	226	14	6	42	4	10
	計	5,028	265	5	940	19	2	1,077	23	2	1,081	59	6	1,167	106	9	640	45	7	123	13	11
CPK	男	230	146	64	189	134	71	14	6	43	14	4	29	12	2	17	1	0	—	0	—	—
	女	97	42	43	94	42	45	2	0	—	0	—	—	1	0	—	0	—	—	0	—	—
	計	327	188	58	283	176	62	16	6	38	14	4	29	13	2	15	1	0	—	0	—	—
CA1919	男	796	8	1	3	0	—	48	0	—	173	1	1	289	3	1	252	1	0	31	3	10
	女	435	7	2	7	0	—	45	0	—	93	1	1	185	4	2	87	1	1	18	1	6
	計	1,231	15	1	10	0	—	93	0	—	266	2	1	474	7	2	339	2	1	49	4	8
CA125	男	2	0	—	0	—	—	0	—	—	1	0	—	1	0	—	0	—	—	0	—	—
	女	504	20	4	21	3	14	82	2	2	136	9	7	172	4	2	77	2	3	16	0	—
	計	506	20	4	21	3	14	82	2	2	137	9	7	173	4	2	77	2	3	16	0	—
シフラ	男	310	6	1.9	1	0	0.0	21	0	—	78	1	1.3	118	3	2.5	81	2	2.5	11	0	—
	女	91	0	—	0	—	—	8	0	—	15	0	—	40	0	—	23	0	—	5	0	—
	計	401	6	1.5	1	0	—	29	0	—	93	1	1.1	158	3	1.9	104	2	1.9	16	0	—
PROGRP	男	316	4	1.3	1	0	—	21	0	—	79	1	1.3	120	1	0.8	84	2	2.4	11	0	—
	女	92	0	—	0	—	—	8	0	—	15	0	—	40	0	—	24	0	—	5	0	—
	計	408	4	1.0	1	0	—	29	0	—	94	1	1.1	160	1	0.6	108	2	1.9	16	0	—
NT-proBNP	男	241	14	5.8	15	0	—	18	0	—	55	1	1.8	84	6	7.1	62	7	11.3	7	0	—
	女	117	4	3.4	2	0	—	11	0	—	30	0	—	41	1	2.4	26	1	3.8	7	2	28.6
	計	358	18	5.0	17	0	—	29	0	—	85	1	1.2	125	7	5.6	88	8	9.1	14	2	14.3

前立腺検査の判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし		4,324	(95.1%)	0	10	63	354	2,735	1,026	136
放置可		0	(0.0%)	0	0	0	0	0	0	0
要経過観察		12	(0.3%)	0	0	0	0	3	8	1
要精密検査		122	(2.7%)	0	0	1	4	64	45	8
要治療		0	(0.0%)	0	0	0	0	0	0	0
治療中		88	(1.9%)	0	0	0	2	33	35	18
合計		4,546		0	10	64	360	2,835	1,114	163

P S A の年齢別平均値

検査項目	総数		29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
P S A n g / m L	4,547	1.30	10	1.03	64	1.10	361	0.96	2,835	1.20	1,114	1.56	163	2.13

骨密度検査判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	112	(51.4%)	0	0	8	21	40	36	7
	女	801	(51.4%)	88	21	141	223	270	52	6
	計	913	(51.4%)	88	21	149	244	310	88	13
放置可	男	0	(0.0%)	0	0	0	0	0	0	0
	女	66	(4.2%)	1	6	32	27	0	0	0
	計	66	(3.7%)	1	6	32	27	0	0	0
要経過観察	男	79	(36.2%)	0	1	4	11	32	21	10
	女	519	(33.3%)	0	7	48	101	211	129	23
	計	598	(33.7%)	0	8	52	112	243	150	33
要精密検査	男	23	(10.6%)	0	0	0	2	7	6	8
	女	136	(8.7%)	0	0	1	11	64	47	13
	計	159	(8.9%)	0	0	1	13	71	53	21
要治療	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
治療中	男	4	(1.8%)	0	0	0	0	1	3	0
	女	37	(2.4%)	0	0	0	2	12	14	9
	計	41	(2.3%)	0	0	0	2	13	17	9
合計	男	218		0	1	12	34	80	66	25
	女	1,559		89	34	222	364	557	242	51
	計	1,777		89	35	234	398	637	308	76

超音波骨密度検査の性年齢別平均値

検査項目	総数		29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		
骨密度 スティフネス	男	218	94.2	1	77.0	12	109.3	34	97.8	80	94.4	66	93.7	25	83.2
	女	1,559	87.1	123	117.4	222	93.6	364	89.2	557	83.5	242	74.7	51	70.8
	計	1,777	88.0	124	117.1	234	94.4	398	89.9	637	84.8	308	78.8	76	74.9

動脈硬化検査判定結果

判 定	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	109 (16.9%)	0	0	11	57	35	6	0
	女	130 (25.9%)	0	0	11	64	45	10	0
	計	239 (20.9%)	0	0	22	121	80	16	0
放置可	男	61 (9.5%)	0	0	3	25	27	6	0
	女	59 (11.8%)	0	0	2	13	35	8	1
	計	120 (10.5%)	0	0	5	38	62	14	1
要経過観察	男	400 (62.1%)	0	0	4	85	172	105	34
	女	280 (55.8%)	0	0	2	33	124	106	15
	計	680 (59.3%)	0	0	6	118	296	211	49
要精密検査	男	66 (10.2%)	0	0	0	5	17	38	6
	女	33 (6.6%)	0	0	0	2	4	16	11
	計	99 (8.6%)	0	0	0	7	21	54	17
要治療	男	0 (0.0%)	0	0	0	0	0	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	0 (0.0%)	0	0	0	0	0	0	0
治療中	男	8 (1.2%)	0	0	0	0	1	5	2
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	8 (0.7%)	0	0	0	0	1	5	2
合計	男	644	0	0	18	172	252	160	42
	女	502	0	0	15	112	208	140	27
	計	1,146	0	0	33	284	460	300	69

頸部動脈超音波検査所見

所 見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
頸動脈の内腔狭窄 (50-75%)	男	17 (2.6%)	0	0	0	0	2	12	3
	女	3 (0.6%)	0	0	0	0	1	0	2
	計	20 (1.7%)	0	0	0	0	3	12	5
頸動脈の高度狭窄 (75%以上)	男	2 (0.3%)	0	0	0	0	0	1	1
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	2 (0.2%)	0	0	0	0	0	1	1
総頸動脈内径拡大	男	4 (0.6%)	0	0	0	0	1	3	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	4 (0.3%)	0	0	0	0	1	3	0
頸動脈描出不良	男	33 (5.1%)	0	0	1	9	10	9	4
	女	16 (3.2%)	0	0	0	3	3	8	2
	計	49 (4.3%)	0	0	1	12	13	17	6
不安定プラーク (III b)	男	3 (0.5%)	0	0	0	0	0	3	0
	女	3 (0.6%)	0	0	0	1	0	1	1
	計	6 (0.5%)	0	0	0	1	0	4	1
不安定プラーク (IV)	男	1 (0.2%)	0	0	0	0	0	1	0
	女	1 (0.2%)	0	0	0	0	0	1	0
	計	2 (0.2%)	0	0	0	0	0	2	0
総頸動脈壁肥厚	男	59 (9.2%)	0	0	0	4	14	29	12
	女	19 (3.8%)	0	0	0	0	3	11	5
	計	78 (6.8%)	0	0	0	4	17	40	17
その他の所見	男	4 (0.6%)	0	0	0	1	1	2	0
	女	1 (0.2%)	0	0	0	0	0	1	0
	計	5 (0.4%)	0	0	0	1	1	3	0
検診者総数	男	644	0	0	18	172	252	160	42
	女	502	0	0	15	112	208	140	27
	計	1,146	0	0	33	284	460	300	69

頸部動脈超音波検査の計測値の性年齢別平均値

検査項目	総数			29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
平均IMT 右mm	男	644	0.740	0	—	18	0.590	172	0.620	252	0.720	160	0.850	42	0.940
	女	502	0.690	0	—	15	0.510	112	0.590	208	0.670	140	0.760	27	0.890
	計	1,146	0.720	0	—	33	0.560	284	0.610	460	0.700	300	0.810	69	0.920
平均IMT 左mm	男	644	0.790	0	—	18	0.620	172	0.670	252	0.790	160	0.880	42	1.050
	女	502	0.720	0	—	15	0.530	112	0.610	208	0.710	140	0.790	27	0.950
	計	1,146	0.760	0	—	33	0.580	284	0.640	460	0.750	300	0.840	69	1.010
最大IMT 右mm	男	644	1.360	0	—	18	0.960	172	1.130	252	1.300	160	1.630	42	1.830
	女	502	1.230	0	—	15	1.010	112	1.060	208	1.230	140	1.330	27	1.470
	計	1,146	1.300	0	—	33	0.980	284	1.100	460	1.270	300	1.490	69	1.690
最大IMT 左mm	男	644	1.400	0	—	18	0.980	172	1.140	252	1.360	160	1.690	42	1.770
	女	502	1.250	0	—	15	0.910	112	1.050	208	1.260	140	1.370	27	1.530
	計	1,146	1.330	0	—	33	0.950	284	1.100	460	1.310	300	1.540	69	1.670

血圧脈波検査の性年齢別平均値

検査項目	総数			29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
ABI検査 右	男	641	1.150	0	—	18	1.130	172	1.140	251	1.150	159	1.160	41	1.150
	女	498	1.120	0	—	15	1.090	111	1.120	207	1.120	139	1.130	26	1.130
	計	1,139	1.140	0	—	33	1.120	283	1.130	458	1.140	298	1.150	67	1.140
ABI検査 左	男	641	1.140	0	—	18	1.110	172	1.120	251	1.140	159	1.150	41	1.140
	女	498	1.110	0	—	15	1.090	111	1.110	207	1.110	139	1.120	26	1.120
	計	1,139	1.130	0	—	33	1.100	283	1.120	458	1.130	298	1.140	67	1.130
PWV検査 右 cm/s	男	641	1434.6	0	—	18	1213.3	172	1313.2	251	1390.1	159	1583.1	41	1737.6
	女	498	1348.9	0	—	15	1139.1	111	1156.3	207	1303.8	139	1516.3	26	1756.7
	計	1,139	1397.1	0	—	33	1179.6	283	1251.7	458	1351.1	298	1551.9	67	1745.0
PWV検査 左 cm/s	男	641	1425.0	0	—	18	1199.4	172	1303.2	251	1385.2	159	1577.7	41	1686.0
	女	498	1360.0	0	—	15	1134.3	111	1160.4	207	1317.2	139	1531.6	26	1764.9
	計	1,139	1396.6	0	—	33	1169.9	283	1247.2	458	1354.5	298	1556.2	67	1716.6

甲状腺検査判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
	男	女	計							
異常なし	男	116	(54.2%)	0	0	13	24	54	24	1
	女	108	(33.8%)	0	4	19	46	24	14	1
	計	224	(41.9%)	0	4	32	70	78	38	2
放置可	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
要経過観察	男	69	(32.2%)	0	0	5	9	24	26	5
	女	127	(39.7%)	0	0	16	34	39	31	7
	計	196	(36.7%)	0	0	21	43	63	57	12
要精密検査	男	29	(13.6%)	0	0	1	6	19	1	2
	女	84	(26.3%)	0	1	8	26	31	16	2
	計	113	(21.2%)	0	1	9	32	50	17	4
要治療	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
治療中	男	0	(0.0%)	0	0	0	0	0	0	0
	女	1	(0.3%)	0	0	0	0	1	0	0
	計	1	(0.2%)	0	0	0	0	1	0	0
合計	男	214		0	0	19	39	97	51	8
	女	320		0	5	43	106	95	61	10
	計	534		0	5	62	145	192	112	18

甲状腺超音波検査所見

所 見	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
	男	女	計							
右葉に甲状腺結節あり	男	70	(37.8%)	0	0	2	9	34	18	7
	女	145	(52.7%)	0	1	14	37	51	35	7
	計	215	(46.7%)	0	1	16	46	85	53	14
左葉に甲状腺結節あり	男	60	(32.4%)	0	0	5	5	25	18	7
	女	147	(53.5%)	0	1	13	46	46	34	7
	計	207	(45.0%)	0	1	18	51	71	52	14
峡部に甲状腺結節あり	男	5	(2.7%)	0	0	0	1	3	1	0
	女	8	(2.9%)	0	0	2	3	0	3	0
	計	13	(2.8%)	0	0	2	4	3	4	0
充実性腫瘍の疑いあり	男	37	(20.0%)	0	0	0	3	24	6	4
	女	100	(36.4%)	0	1	7	27	36	25	4
	計	137	(29.8%)	0	1	7	30	60	31	8
嚢胞性病変の疑いあり	男	114	(61.6%)	0	0	7	12	51	33	11
	女	249	(90.5%)	0	2	25	69	79	62	12
	計	363	(78.9%)	0	2	32	81	130	95	23
副甲状腺腫大あり	男	0	(0.0%)	0	0	0	0	0	0	0
	女	1	(0.4%)	0	0	0	1	0	0	0
	計	1	(0.2%)	0	0	0	1	0	0	0
甲状腺腫大あり	男	4	(2.2%)	0	0	0	0	4	0	0
	女	16	(5.8%)	0	0	1	9	4	2	0
	計	20	(4.3%)	0	0	1	9	8	2	0
その他の所見	男	12	(6.5%)	0	0	1	1	5	3	2
	女	30	(10.9%)	0	0	2	9	11	8	0
	計	42	(9.1%)	0	0	3	10	16	11	2
検診者総数	男	185		0	0	14	28	90	45	8
	女	275		0	4	30	84	89	58	10
	計	460		0	4	44	112	179	103	18

甲状腺機能検査の性年齢別平均値

検査項目	総数			29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
T S H μ U / m L	男	72	1.7	0	—	8	2.0	18	1.4	27	1.4	18	2.1	1	6.6
	女	164	1.9	1	2.7	31	2.1	69	1.7	40	1.8	18	1.9	5	3.3
	計	236	1.8	1	2.7	39	2.1	87	1.6	67	1.6	36	2.0	6	3.8
F T 3 p g / m L	男	72	3.1	0	—	8	3.3	18	3.1	27	3.1	18	3.0	1	3.2
	女	163	3.2	1	3.1	30	2.9	69	3.2	40	3.7	18	2.8	5	2.9
	計	235	3.2	1	3.1	38	3.0	87	3.1	67	3.4	36	2.9	6	2.9
F T 4 n g / d L	男	72	1.3	0	—	8	1.4	18	1.2	27	1.3	18	1.3	1	1.5
	女	163	1.3	1	1.3	30	1.2	69	1.3	40	1.4	18	1.2	5	1.2
	計	235	1.3	1	1.3	38	1.2	87	1.3	67	1.3	36	1.2	6	1.3

脳検査判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	302	(41.0%)	0	1	35	76	149	25	16
	女	225	(48.4%)	0	1	22	52	92	39	19
	計	527	(43.9%)	0	2	57	128	241	64	35
放置可	男	387	(52.6%)	0	0	10	53	182	108	34
	女	209	(44.9%)	0	0	0	19	80	67	43
	計	596	(49.6%)	0	0	10	72	262	175	77
要経過観察	男	23	(3.1%)	0	0	2	9	8	4	0
	女	16	(3.4%)	0	0	1	2	7	3	3
	計	39	(3.2%)	0	0	3	11	15	7	3
要精密検査	男	14	(1.9%)	0	0	0	2	9	0	3
	女	13	(2.8%)	0	0	0	3	5	3	2
	計	27	(2.2%)	0	0	0	5	14	3	5
要治療	男	1	(0.1%)	0	0	0	0	0	1	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	1	(0.1%)	0	0	0	0	0	1	0
治療中	男	9	(1.2%)	0	0	0	3	2	2	2
	女	2	(0.4%)	0	0	0	0	0	2	0
	計	11	(0.9%)	0	0	0	3	2	4	2
合計	男	736		0	1	47	143	350	140	55
	女	465		0	1	23	76	184	114	67
	計	1,201		0	2	70	219	534	254	122

頭頸部MR I 検査所見

所 見	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
無症候性脳梗塞巣	男	37	(5.0%)	0	0	0	0	14	17	6
	女	9	(1.9%)	0	0	0	0	3	2	4
	計	46	(3.8%)	0	0	0	0	17	19	10
脳内異常陰影	男	0	(0.0%)	0	0	0	0	0	0	0
	女	2	(0.4%)	0	0	0	0	1	0	1
	計	2	(0.2%)	0	0	0	0	1	0	1
中耳炎症所見	男	2	(0.3%)	0	0	0	1	1	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	2	(0.2%)	0	0	0	1	1	0	0
副鼻腔炎	男	38	(5.2%)	0	0	4	12	13	8	1
	女	14	(3.0%)	0	0	1	3	8	1	1
	計	52	(4.3%)	0	0	5	15	21	9	2
硬膜下水腫	男	1	(0.1%)	0	0	0	0	0	0	1
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	1	(0.1%)	0	0	0	0	0	0	1
くも膜嚢胞	男	6	(0.8%)	0	0	0	1	4	1	0
	女	3	(0.6%)	0	0	0	0	2	1	0
	計	9	(0.7%)	0	0	0	1	6	2	0
その他の所見	男	16	(2.2%)	0	0	0	3	8	4	1
	女	9	(1.9%)	0	0	0	1	4	3	1
	計	25	(2.1%)	0	0	0	4	12	7	2
検診者総数	男	736		0	1	47	143	350	140	55
	女	465		0	1	23	76	184	114	67
	計	1,201		0	2	70	219	534	254	122

頭頸部MR A 検査所見

所 見	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
脳動脈硬化	男	16	(2.2%)	0	0	0	0	2	4	10
	女	6	(1.3%)	0	0	0	0	0	0	6
	計	22	(1.8%)	0	0	0	0	2	4	16
脳動脈瘤	男	7	(1.0%)	0	0	0	0	5	0	2
	女	4	(0.9%)	0	0	0	1	2	0	1
	計	11	(0.9%)	0	0	0	1	7	0	3
内頸動脈狭窄	男	3	(0.4%)	0	0	0	0	1	1	1
	女	2	(0.4%)	0	0	0	0	1	0	1
	計	5	(0.4%)	0	0	0	0	2	1	2
脳動脈狭窄	男	1	(0.1%)	0	0	0	0	1	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	1	(0.1%)	0	0	0	0	1	0	0
脳動脈閉塞	男	1	(0.1%)	0	0	0	0	1	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	1	(0.1%)	0	0	0	0	1	0	0
遺残性三叉動脈	男	1	(0.1%)	0	0	0	0	1	0	0
	女	2	(0.4%)	0	0	0	1	0	0	1
	計	3	(0.2%)	0	0	0	1	1	0	1
その他の所見	男	4	(0.5%)	0	0	0	0	2	2	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	4	(0.3%)	0	0	0	0	2	2	0
検診者総数	男	736		0	1	47	143	350	140	55
	女	465		0	1	23	76	184	114	67
	計	1,201		0	2	70	219	534	254	122

大脳高次機能検査所見

所 見	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
MMSE (脳の認知機能評価の簡易検査) 低下	男	0	(0.0%)	0	0	0	0	0	0	0
	女	1	(0.6%)	0	0	0	0	0	0	1
	計	1	(0.3%)	0	0	0	0	0	0	1
検診者総数	男	125		0	0	5	9	10	58	43
	女	170		0	0	2	5	24	81	58
	計	295		0	0	7	14	34	139	101

心理検査所見

所 見	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
抑うつ状態	男	27	(21.6%)	0	0	1	6	1	11	8
	女	26	(15.3%)	0	0	0	1	6	14	5
	計	53	(18.0%)	0	0	1	7	7	25	13
検診者総数	男	125		0	0	5	9	10	58	43
	女	170		0	0	2	5	24	81	58
	計	295		0	0	7	14	34	139	101

肥満やせ判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	16,896	(55.8%)	272	2,837	3,931	3,943	3,673	1,974	266
	女	14,997	(63.8%)	213	2,492	3,181	3,949	3,435	1,541	186
	計	31,893	(59.3%)	485	5,329	7,112	7,892	7,108	3,515	452
放置可	男	1,113	(3.7%)	34	272	293	198	186	110	20
	女	4,453	(18.9%)	39	913	1,122	1,094	945	290	50
	計	5,566	(10.3%)	73	1,185	1,415	1,292	1,131	400	70
要経過観察	男	12,271	(40.5%)	52	908	2,345	3,451	3,472	1,843	200
	女	4,069	(17.3%)	16	329	662	1,156	1,172	642	92
	計	16,340	(30.4%)	68	1,237	3,007	4,607	4,644	2,485	292
要精密検査	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
要治療	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
治療中	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
合 計	男	30,280		358	4,017	6,569	7,592	7,331	3,927	486
	女	23,519		268	3,734	4,965	6,199	5,552	2,473	328
	計	53,799		626	7,751	11,534	13,791	12,883	6,400	814

身体計測の性年齢別平均値

検査項目	総数			29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
身長 c m	男	30,635	170.27	4,459	171.50	6,613	171.74	7,698	171.16	7,409	169.51	3,969	166.94	487	163.95
	女	23,624	157.34	4,016	158.32	5,017	158.48	6,233	158.25	5,557	156.58	2,473	153.60	328	151.26
	計	54,259	164.64	8,475	165.25	11,630	166.02	13,931	165.39	12,966	163.97	6,442	161.82	815	158.84
体重 k g	男	30,890	69.06	4,505	66.95	6,674	70.11	7,764	71.13	7,459	69.26	4,001	66.00	487	62.91
	女	23,658	54.07	4,022	52.09	5,022	53.68	6,236	55.30	5,565	54.79	2,485	53.63	328	52.10
	計	54,548	62.56	8,527	59.94	11,696	63.05	14,000	64.08	13,024	63.08	6,486	61.26	815	58.56
B M I (体格指数)	男	30,635	23.80	4,459	22.75	6,613	23.75	7,698	24.27	7,409	24.08	3,969	23.67	487	23.39
	女	23,624	21.84	4,016	20.76	5,017	21.36	6,233	22.08	5,557	22.35	2,473	22.74	328	22.77
	計	54,259	22.95	8,475	21.81	11,630	22.72	13,931	23.29	12,966	23.34	6,442	23.31	815	23.14
体脂肪率 %	男	3,997	21.74	253	18.11	732	20.50	1,103	22.02	1,336	22.54	487	22.52	86	22.73
	女	3,366	30.01	419	28.47	689	29.08	948	29.90	975	30.89	284	31.68	51	31.54
	計	7,363	25.52	672	24.57	1,421	24.66	2,051	25.66	2,311	26.06	771	25.89	137	26.01
腹囲 c m	男	25,318	83.73	2,108	78.33	4,825	82.55	7,225	84.50	6,999	84.86	3,707	84.63	454	84.27
	女	18,557	77.72	1,498	71.66	3,198	74.87	5,835	77.41	5,328	79.46	2,388	81.57	310	82.98
	計	43,875	81.19	3,606	75.56	8,023	79.49	13,060	81.33	12,327	82.52	6,095	83.43	764	83.75

特定健診判定結果

判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
該当	男	3,380	(15.6%)	0	16	281	886	1,321	789	87
	女	438	(3.1%)	0	0	13	91	166	138	30
	計	3,818	(10.6%)	0	16	294	977	1,487	927	117
予備群該当	男	3,449	(15.9%)	2	74	480	1,156	1,104	574	59
	女	607	(4.3%)	0	3	40	186	237	120	21
	計	4,056	(11.3%)	2	77	520	1,342	1,341	694	80
非該当	男	14,497	(67.0%)	46	885	2709	4,521	4,091	1,995	250
	女	12,606	(88.8%)	25	410	1797	4536	3999	1637	202
	計	27,103	(75.6%)	71	1295	4506	9,057	8,090	3,632	452
判定不能	男	325	(1.5%)	2	76	72	78	62	24	11
	女	549	(3.9%)	2	166	226	107	37	5	6
	計	874	(2.4%)	4	242	298	185	99	29	17
合計	男	21,651		50	1051	3,542	6,641	6,578	3,382	407
	女	14,200		27	579	2,076	4,920	4,439	1,900	259
	計	35,851		77	1,630	5,618	11,561	11,017	5,282	666

体力測定 of 性年齢別平均値

検査項目		総数		29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
全身反応時間 m 秒	男	827	334.1	185	304.4	190	340.4	238	345.5	172	337.1	36	357.9	6	370.8
	女	280	366.0	93	322.8	51	389.1	69	389.5	45	375.3	18	401.8	4	405.3
	計	1,107	342.2	278	310.6	241	350.7	307	355.4	217	345.0	54	372.5	10	384.6
立位体前屈 c m	男	825	4.66	184	10.57	190	3.61	237	2.81	172	2.60	36	1.64	6	6.78
	女	279	8.93	94	11.64	50	9.56	69	5.75	44	7.45	18	9.67	4	5.38
	計	1,104	5.74	278	10.93	240	4.85	306	3.47	216	3.59	54	4.31	10	6.22
上体おこし 回	男	817	22.6	182	28.9	189	24.1	236	20.7	169	18.5	35	13.9	6	14.5
	女	277	16.0	94	26.9	50	13.8	67	11.5	44	8.5	18	2.8	4	4.5
	計	1094	20.9	276	28.2	239	21.9	303	18.7	213	16.5	53	10.2	10	10.5
閉眼片足立ち 秒	男	689	57.55	47	104.02	190	80.17	238	55.53	172	32.65	36	18.17	6	7.83
	女	187	46.25	0	0.00	51	64.43	69	55.25	45	28.93	18	11.22	4	11.75
	計	876	55.14	47	104.02	241	76.84	307	55.47	217	31.88	54	15.85	10	9.40
最大酸素摂取量 mL/分/ k g	男	679	35.26	47	42.72	189	36.94	235	34.71	167	33.02	36	31.38	5	29.82
	女	183	31.50	0	0.00	51	33.00	68	31.86	44	31.87	16	24.17	4	31.80
	計	862	34.46	47	42.72	240	36.10	303	34.07	211	32.78	52	29.16	9	30.70

V D T 健診管理区分										
判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	3,128	(78.1%)	14	365	714	1,002	870	163	0
	女	1,352	(82.8%)	6	309	423	397	193	24	0
	計	4,480	(79.5%)	20	674	1,137	1,399	1063	187	0
要経過観察	男	561	(14.0%)	0	45	44	119	275	78	0
	女	169	(10.3%)	0	31	31	43	56	8	0
	計	730	(13.0%)	0	76	75	162	331	86	0
二次検査後 要経過観察	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
要就労制限	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
就労により 悪化の可能性あり	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
就労以外の 原因による 異常	男	264	(6.6%)	1	8	32	90	117	15	1
	女	87	(5.3%)	0	9	21	33	20	4	0
	計	351	(6.2%)	1	17	53	123	137	19	1
要二次検査	男	50	(1.2%)	1	12	8	9	15	5	0
	女	25	(1.5%)	1	7	8	5	3	1	0
	計	75	(1.3%)	2	19	16	14	18	6	0
合 計	男	4,003		16	430	798	1,220	1,277	261	1
	女	1,633		7	356	483	478	272	37	0
	計	5,636		23	786	1,281	1,698	1,549	298	1

特定化学物質管理区分										
判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	360	(94.5%)	5	86	100	95	54	20	0
	女	18	(100.0%)	0	4	1	5	4	4	0
	計	378	(94.7%)	5	90	101	100	58	24	0
要経過観察	男	20	(5.2%)	1	3	6	3	5	2	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	20	(5.0%)	1	3	6	3	5	2	0
二次検査後 要経過観察	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
要就労制限	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
就労により 悪化の可能性あり	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
就労以外の 原因による 異常	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
要二次検査	男	1	(0.3%)	0	0	0	0	1	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	1	(0.3%)	0	0	0	0	1	0	0
合 計	男	381		6	89	106	98	60	22	0
	女	18		0	4	1	5	4	4	0
	計	399		6	93	107	103	64	26	0

騒音・高気圧業務健診管理区分										
判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	139	(37.9%)	3	64	17	45	10	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	139	(37.9%)	3	64	17	45	10	0	0
要経過観察	男	15	(4.1%)	0	8	5	2	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	15	(4.1%)	0	8	5	2	0	0	0
二次検査後 要経過観察	男	210	(57.2%)	0	27	14	52	76	41	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	210	(57.2%)	0	27	14	52	76	41	0
要就労制限	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
就労により 悪化の可能性あり	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
就労以外の 原因による 異常	男	3	(0.8%)	0	0	1	1	1	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	3	(0.8%)	0	0	1	1	1	0	0
要二次検査	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
合 計	男	367		3	99	37	100	87	41	0
	女	0		0	0	0	0	0	0	0
	計	367		3	99	37	100	87	41	0

電離放射線健診管理区分										
判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	61	(92.4%)	1	11	18	18	8	5	0
	女	2	(40.0%)	0	1	0	0	0	1	0
	計	63	(88.7%)	1	12	18	18	8	6	0
要経過観察	男	5	(7.6%)	0	2	1	0	2	0	0
	女	2	(40.0%)	0	0	0	1	0	1	0
	計	7	(9.9%)	0	2	1	1	2	1	0
二次検査後 要経過観察	男	0	(0.0%)	0	0	0	0	0	0	0
	女	1	(20.0%)	0	0	0	1	0	0	0
	計	1	(1.4%)	0	0	0	1	0	0	0
要就労制限	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
就労により 悪化の可能性あり	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
就労以外の 原因による 異常	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
要二次検査	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
合 計	男	66		1	13	19	18	10	5	0
	女	5		0	1	0	2	0	2	0
	計	71		1	14	19	20	10	7	0

有機溶剤健診管理区分										
判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	580	(80.7%)	1	119	132	142	125	61	0
	女	88	(82.2%)	0	17	19	19	12	21	0
	計	668	(80.9%)	1	136	151	161	137	82	0
要経過観察	男	93	(12.9%)	2	22	14	24	25	6	0
	女	19	(17.8%)	0	4	3	6	4	2	0
	計	112	(13.6%)	2	26	17	30	29	8	0
二次検査後 要経過観察	男	6	(0.8%)	0	1	1	3	0	1	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	6	(0.7%)	0	1	1	3	0	1	0
要就労制限	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
就労により 悪化の可能性あり	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
就労以外の 原因による 異常	男	16	(2.2%)	0	2	0	5	8	1	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	16	(1.9%)	0	2	0	5	8	1	0
要二次検査	男	24	(3.3%)	0	4	8	8	3	1	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	24	(2.9%)	0	4	8	8	3	1	0
合 計	男	719		3	148	155	182	161	70	0
	女	107		0	21	22	25	16	23	0
	計	826		3	169	177	207	177	93	0

じん肺・アスベスト健診管理区分										
判 定	総数			19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
異常なし	男	71	(84.5%)	0	4	7	3	30	26	1
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	71	(84.5%)	0	4	7	3	30	26	1
要経過観察	男	7	(8.3%)	0	0	0	0	4	3	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	7	(8.3%)	0	0	0	0	4	3	0
二次検査後 要経過観察	男	1	(1.2%)	0	0	0	0	0	1	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	1	(1.2%)	0	0	0	0	0	1	0
要就労 制限	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
就労により 悪化の可能 性あり	男	0	(0.0%)	0	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0	0
就労以外の 原因による 異常	男	4	(4.8%)	0	0	0	0	2	2	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	4	(4.8%)	0	0	0	0	2	2	0
要二次 検査	男	1	(1.2%)	0	1	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0
	計	1	(1.2%)	0	1	0	0	0	0	0
合 計	男	84		0	5	7	3	36	32	1
	女	0		0	0	0	0	0	0	0
	計	84		0	5	7	3	36	32	1

精密検査判定者の他医療機関からの回答状況

精査項目	対象者	回答数	回答率	回答結果				
				異常なし	放置可	経過観察	治療中	要治療
視 機 能	2	1	50.0%	0 0.0%	0 0.0%	0 0.0%	0 0.0%	1 100.0%
肺 機 能	19	14	73.7%	1 7.1%	3 21.4%	4 28.6%	3 21.4%	3 21.4%
血 圧	1,177	592	50.3%	36 6.1%	42 7.1%	187 31.6%	30 5.1%	297 50.2%
脂 質	3,760	2,117	56.3%	461 21.8%	487 23.0%	592 28.0%	72 3.4%	503 23.8%
糖 尿 病	824	463	56.2%	37 8.0%	15 3.2%	164 35.4%	47 10.2%	195 42.1%
痛 風	675	380	56.3%	39 10.3%	24 6.3%	131 34.5%	24 6.3%	162 42.6%
肝機能・血清蛋白	4,649	2,823	60.7%	287 10.2%	476 16.9%	1,648 58.4%	183 6.5%	226 8.0%
甲 状 腺 検 査	113	103	91.2%	2 1.9%	25 24.3%	62 60.2%	1 1.0%	6 5.8%
脾 機 能	36	28	77.8%	10 35.7%	4 14.3%	12 42.9%	2 7.1%	0 0.0%
腎 機 能	509	321	63.1%	66 20.6%	36 11.2%	149 46.4%	45 14.0%	25 7.8%
貧 血 ・ 血 球 数	716	424	59.2%	28 6.6%	47 11.1%	110 25.9%	30 7.1%	206 48.6%
電 解 質	1	1	100.0%	0 0.0%	0 0.0%	0 0.0%	1 100.0%	0 0.0%
その他の血液検査	143	83	58.0%	18 21.7%	28 33.7%	26 31.3%	2 2.4%	4 4.8%
便 潜 血	1,383	880	63.6%	263 29.9%	204 23.2%	143 16.3%	8 0.9%	131 14.9%
眼 底 検 査	604	459	76.0%	92 20.0%	99 21.6%	163 35.5%	7 1.5%	90 19.6%
胸 部 検 査	676	516	76.3%	220 42.6%	127 24.6%	133 25.8%	8 1.6%	19 3.7%
心 電 図	240	163	67.9%	22 13.5%	45 27.6%	62 38.0%	11 6.7%	22 13.5%
上 部 消 化 管	1,019	799	78.4%	69 8.6%	277 34.7%	188 23.5%	9 1.1%	244 30.5%
大 腸 検 査	4	4	100.0%	1 25.0%	0 0.0%	2 50.0%	0 0.0%	0 0.0%
腹 部 超 音 波	577	440	76.3%	37 8.4%	124 28.2%	245 55.7%	2 0.5%	19 4.3%
内 科 診 察	75	44	58.7%	7 15.9%	7 15.9%	16 36.4%	0 0.0%	10 22.7%
乳 房 検 査	612	546	89.2%	185 33.9%	170 31.1%	155 28.4%	8 1.5%	3 0.5%
婦 人 科 検 査	518	416	80.3%	20 4.8%	37 8.9%	305 73.3%	6 1.4%	35 8.4%
前 立 腺 検 査	122	86	70.5%	9 10.5%	12 14.0%	47 54.7%	3 3.5%	12 14.0%
骨 密 度	159	91	57.2%	21 23.1%	14 15.4%	18 19.8%	2 2.2%	36 39.6%
動 脈 硬 化 検 査	99	72	72.7%	5 6.9%	8 11.1%	33 45.8%	8 11.1%	18 25.0%
脳 検 査	28	22	78.6%	1 4.5%	4 18.2%	16 72.7%	1 4.5%	0 0.0%
睡眠時無呼吸検査	64	42	65.6%	1 2.4%	7 16.7%	10 23.8%	0 0.0%	23 54.8%

市民健診総合判定結果

判 定	総数			35-39歳	40-49歳	50-59歳	60-69歳	70-79歳	80歳以上
異常なし	男	32	(0.3%)	5	10	6	6	2	3
	女	179	(1.0%)	45	99	14	17	4	0
	計	211	(0.7%)	50	109	20	23	6	3
放置可	男	602	(5.7%)	14	82	54	205	211	36
	女	2,395	(13.5%)	161	555	466	780	391	42
	計	2,997	(10.6%)	175	637	520	985	602	78
要経過観察	男	3,610	(33.9%)	31	252	242	1,222	1,480	383
	女	6,120	(34.6%)	86	392	768	2,591	1,869	414
	計	9,730	(34.3%)	117	644	1,010	3,813	3,349	797
要精密検査	男	1,023	(9.6%)	8	86	78	352	378	121
	女	1,483	(8.4%)	29	111	198	599	414	132
	計	2,506	(8.8%)	37	197	276	951	792	253
要治療	男	396	(3.7%)	4	40	40	152	121	39
	女	512	(2.9%)	3	24	67	240	153	25
	計	908	(3.2%)	7	64	107	392	274	64
治療中	男	4,972	(46.8%)	4	52	132	1,431	2,534	819
	女	7,013	(39.6%)	2	45	325	2,505	3,129	1,007
	計	11,985	(42.3%)	6	97	457	3,936	5,663	1,826
合計	男	10,635		66	522	552	3,368	4,726	1,401
	女	17,702		326	1,226	1,838	6,732	5,960	1,620
	計	28,337		392	1,748	2,390	10,100	10,686	3,021

市民健診メタボ判定結果

判 定	総数			40-49歳	50-59歳	60-69歳	70-74歳
該当	男	1,543	(21.1%)	79	108	740	616
	女	830	(6.2%)	16	62	447	305
	計	2,373	(11.4%)	95	170	1,187	921
予備群該当	男	1,313	(18.0%)	107	105	604	497
	女	783	(5.8%)	41	76	388	278
	計	2,096	(10.1%)	148	181	992	775
非該当	男	4,442	(60.8%)	336	339	2,018	1,749
	女	11,863	(87.9%)	1,168	1,700	5,890	3,105
	計	16,305	(78.4%)	1,504	2,039	7,908	4,854
判定不能	男	10	(0.1%)	0	0	4	6
	女	17	(0.1%)	1	0	3	13
	計	27	(0.1%)	1	0	7	19
合計	男	7,308		522	552	3,366	2,868
	女	13,493		1,226	1,838	6,728	3,701
	計	20,801		1,748	2,390	10,094	6,569

市民健診肥満判定結果

判 定	総数			35-39歳	40-49歳	50-59歳	60-69歳	70-79歳	80歳以上
異常なし	男	5,967	(56.1%)	36	236	267	1,701	2,718	1,009
	女	12,233	(69.1%)	212	883	1,249	4,726	4,055	1,108
	計	18,200	(64.2%)	248	1,119	1,516	6,427	6,773	2,117
放置可	男	426	(4.0%)	2	19	18	89	194	104
	女	1,817	(10.3%)	77	184	277	605	483	191
	計	2,243	(7.9%)	79	203	295	694	677	295
要経過観察	男	4,242	(39.9%)	28	267	267	1,578	1,814	288
	女	3,652	(20.6%)	37	159	312	1,401	1,422	321
	計	7,894	(27.9%)	65	426	579	2,979	3,236	609
要精密検査	男	0	(0.0%)	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0
要治療	男	0	(0.0%)	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0
治療中	男	0	(0.0%)	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0
合 計	男	10,635		66	522	552	3,368	4,726	1,401
	女	17,702		326	1,226	1,838	6,732	5,960	1,620
	計	28,337		392	1,748	2,390	10,100	10,686	3,021

市民健診身体計測の性年齢別平均値

検査項目	総数			35-39歳		40-49歳		50-59歳		60-69歳		70-79歳		80歳以上	
身長 c m	男	10,635	164.14	66	171.19	522	169.78	552	168.34	3,368	165.66	4,726	163.15	1,401	159.73
	女	17,702	152.09	326	158.39	1,226	158.06	1,838	155.65	6,732	152.97	5,960	150.00	1,620	146.34
	計	28,337	156.61	392	160.54	1,748	161.56	2,390	158.58	10,100	157.20	10,686	155.82	3,021	152.55
体重 k g	男	10,635	63.13	66	69.59	522	70.01	552	67.60	3,368	64.57	4,726	62.31	1,401	57.81
	女	17,702	51.48	326	52.24	1,226	53.60	1,838	52.74	6,732	52.09	5,960	50.90	1,620	47.84
	計	28,337	55.85	392	55.16	1,748	58.50	2,390	56.17	10,100	56.25	10,686	55.95	3,021	52.46
B M I (体格指数)	男	10,635	23.39	66	23.75	522	24.24	552	23.83	3,368	23.50	4,726	23.38	1,401	22.63
	女	17,702	22.25	326	20.82	1,226	21.45	1,838	21.78	6,732	22.26	5,960	22.62	1,620	22.33
	計	28,337	22.68	392	21.31	1,748	22.28	2,390	22.25	10,100	22.67	10,686	22.96	3,021	22.47
腹囲 c m	男	7,371	84.19	66	83.07	522	85.08	552	84.60	3,366	84.32	2,865	83.81	0	—
	女	13,808	80.37	326	73.84	1,226	76.19	1,838	78.28	6,726	80.91	3,692	82.40	0	—
	計	21,179	81.70	392	75.39	1,748	78.85	2,390	79.74	10,092	82.04	6,557	83.02	0	—

市民健診血圧判定結果

判 定	総数			35-39歳	40-49歳	50-59歳	60-69歳	70-79歳	80歳以上
異常なし	男	2,806	(26.4%)	50	320	243	967	1020	206
	女	7,017	(39.6%)	299	1,043	1,127	2,791	1,555	202
	計	9,823	(34.7%)	349	1,363	1,370	3,758	2,575	408
放置可	男	0	(0.0%)	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0
要経過観察	男	3,188	(30.0%)	11	146	174	1,067	1,394	396
	女	4,840	(27.3%)	22	136	455	1,990	1,813	424
	計	8,028	(28.3%)	33	282	629	3,057	3,207	820
要精密検査	男	561	(5.3%)	0	22	30	177	245	87
	女	700	(4.0%)	3	20	60	250	268	99
	計	1,261	(4.5%)	3	42	90	427	513	186
要治療	男	141	(1.3%)	1	5	13	49	54	19
	女	134	(0.8%)	0	5	14	46	56	13
	計	275	(1.0%)	1	10	27	95	110	32
治療中	男	3,939	(37.0%)	4	29	92	1,108	2,013	693
	女	5,011	(28.3%)	2	22	182	1,655	2,268	882
	計	8,950	(31.6%)	6	51	274	2,763	4,281	1,575
合 計	男	10,635		66	522	552	3,368	4,726	1,401
	女	17,702		326	1,226	1,838	6,732	5,960	1,620
	計	28,337		392	1,748	2,390	10,100	10,686	3,021

市民健診安静時血圧の性年齢別平均値

検査項目	総数			35-39歳		40-49歳		50-59歳		60-69歳		70-79歳		80歳以上	
安静時血圧・ 最大・ mmHg	男	10,635	136.6	66	121.0	522	124.7	552	129.5	3,368	135.4	4,726	138.3	1,401	142.0
	女	17,702	131.7	326	111.4	1,226	115.3	1,838	123.9	6,732	130.9	5,960	136.6	1,620	142.4
	計	28,337	133.5	392	113.0	1,748	118.1	2,390	125.2	10,100	132.4	10,686	137.3	3,021	142.2
安静時血圧・ 最小・ mmHg	男	10,635	77.3	66	73.1	522	79.6	552	81.3	3,368	79.9	4,726	76.1	1,401	72.4
	女	17,702	73.3	326	68.3	1,226	70.3	1,838	74.3	6,732	74.5	5,960	73.0	1,620	71.9
	計	28,337	74.8	392	69.1	1,748	73.1	2,390	75.9	10,100	76.3	10,686	74.4	3,021	72.1

市民健診肝機能検査判定結果

判 定	総数			35-39歳	40-49歳	50-59歳	60-69歳	70-79歳	80歳以上
異常なし	男	7,029	(66.3%)	39	274	313	2,135	3,231	1,037
	女	14,689	(83.1%)	310	1,092	1,501	5,518	4,906	1,362
	計	21,718	(76.8%)	349	1,366	1,814	7,653	8,137	2,399
放置可	男	1,844	(17.4%)	14	91	112	599	812	216
	女	1,862	(10.5%)	9	66	199	738	680	170
	計	3,706	(13.1%)	23	157	311	1,337	1,492	386
要経過観察	男	1,104	(10.4%)	6	76	79	420	434	89
	女	717	(4.1%)	6	44	76	315	220	56
	計	1,821	(6.4%)	12	120	155	735	654	145
要精密検査	男	420	(4.0%)	6	61	40	146	138	29
	女	182	(1.0%)	1	16	42	70	48	5
	計	602	(2.1%)	7	77	82	216	186	34
要治療	男	49	(0.5%)	1	12	3	19	14	0
	女	41	(0.2%)	0	3	7	18	13	0
	計	90	(0.3%)	1	15	10	37	27	0
治療中	男	161	(1.5%)	0	6	5	42	87	21
	女	181	(1.0%)	0	5	11	63	81	21
	計	342	(1.2%)	0	11	16	105	168	42
合 計	男	10,607		66	520	552	3,361	4,716	1,392
	女	17,672		326	1,226	1,836	6,722	5,948	1,614
	計	28,279		392	1,746	2,388	10,083	10,664	3,006

市民健診肝機能検査の性年齢別平均値

検査項目	総数			35-39歳		40-49歳		50-59歳		60-69歳		70-79歳		80歳以上	
GOT・IU/L	男	10,607	25.9	66	23.4	520	25.9	552	24.7	3,361	25.7	4,716	26.2	1,392	25.7
	女	17,672	23.8	326	18.4	1,226	20.8	1,836	23.1	6,722	23.8	5,948	24.8	1,614	24.6
	計	28,279	24.6	392	19.3	1,746	22.3	2,388	23.5	10,083	24.4	10,664	25.4	3,006	25.1
GPT・IU/L	男	10,607	23.1	66	30.2	520	32.3	552	26.1	3,361	24.3	4,716	22.0	1,392	19.0
	女	17,672	19.3	326	14.8	1,226	17.9	1,836	20.7	6,722	20.1	5,948	19.2	1,614	17.0
	計	28,279	20.8	392	17.4	1,746	22.2	2,388	22.0	10,083	21.5	10,664	20.4	3,006	17.9
γ-GTP・IU/L	男	10,607	42.8	66	43.5	520	53.4	552	53.2	3,361	46.1	4,716	40.8	1,392	33.1
	女	17,672	24.2	326	18.7	1,226	23.6	1,836	27.1	6,722	25.7	5,948	23.0	1,614	20.6
	計	28,279	31.2	392	22.9	1,746	32.5	2,388	33.1	10,083	32.5	10,664	30.9	3,006	26.4

市民健診腎機能検査判定結果

判 定	総数			35-39歳	40-49歳	50-59歳	60-69歳	70-79歳	80歳以上
異常なし	男	489	(4.6%)	21	111	61	158	119	19
	女	1,256	(7.1%)	133	305	202	406	182	28
	計	1,745	(6.2%)	154	416	263	564	301	47
放置可	男	6,313	(59.4%)	42	367	415	2,281	2,653	555
	女	11,867	(67.0%)	183	857	1,455	5,009	3,684	679
	計	18,180	(64.2%)	225	1,224	1,870	7,290	6,337	1,234
要経過観察	男	3,072	(28.9%)	2	36	65	665	1,645	659
	女	3,978	(22.5%)	7	58	150	1,078	1,909	776
	計	7,050	(24.9%)	9	94	215	1,743	3,554	1,435
要精密検査	男	586	(5.5%)	1	7	8	223	211	136
	女	485	(2.7%)	3	4	20	205	131	122
	計	1,071	(3.8%)	4	11	28	428	342	258
要治療	男	0	(0.0%)	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0
	計	0	(0.0%)	0	0	0	0	0	0
治療中	男	170	(1.6%)	0	1	3	40	97	29
	女	113	(0.6%)	0	2	10	34	53	14
	計	283	(1.0%)	0	3	13	74	150	43
合 計	男	10,630		66	522	552	3,367	4,725	1,398
	女	17,699		326	1,226	1,837	6,732	5,959	1,619
	計	28,329		392	1,748	2,389	10,099	10,684	3,017

市民健診腎機能検査の性年齢別平均値

検査項目	総数			35-39歳		40-49歳		50-59歳		60-69歳		70-79歳		80歳以上	
クレアチニン・ mg / dL	男	10,607	0.92	66	0.83	520	0.84	552	0.85	3,361	0.89	4,716	0.93	1,392	1.00
	女	17,672	0.67	326	0.62	1,226	0.63	1,836	0.64	6,722	0.66	5,948	0.69	1,614	0.75
	計	28,279	0.76	392	0.65	1,746	0.69	2,388	0.69	10,083	0.74	10,664	0.79	3,006	0.86

市民健診尿監査の性年齢別陽性率

検査項目	性 別	総数			35-39歳			40-49歳			50-59歳			60-69歳			70-79歳			80歳以上		
		受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率
尿蛋白定性	男	10,538	929	8.8	66	2	3.0	520	30	5.8	548	29	5.3	3,345	255	7.6	4,684	438	9.4	1,375	175	12.7
	女	17,596	612	3.5	323	9	2.8	1,221	36	2.9	1,833	46	2.5	6,714	185	2.8	5,915	227	3.8	1,590	109	6.9
	計	28,134	1,541	5.5	389	11	2.8	1,741	66	3.8	2,381	75	3.1	10,059	440	4.4	10,599	665	6.3	2,965	284	9.6

陽性率%

市民健診脂質検査判定結果

判 定	総数			35-39歳	40-49歳	50-59歳	60-69歳	70-79歳	80歳以上
異常なし	男	3,446	(32.5%)	23	146	133	976	1,537	631
	女	4,682	(26.5%)	188	624	492	1,340	1,482	556
	計	8,128	(28.7%)	211	770	625	2,316	3,019	1,187
放置可	男	2,541	(24.0%)	14	110	129	781	1,207	300
	女	3,920	(22.2%)	72	313	434	1,418	1,302	381
	計	6,461	(22.8%)	86	423	563	2,199	2,509	681
要経過観察	男	3,123	(29.4%)	24	188	204	1,106	1,290	311
	女	4,921	(27.8%)	45	204	590	2,194	1,529	359
	計	8,044	(28.4%)	69	392	794	3,300	2,819	670
要精密検査	男	333	(3.1%)	3	34	33	123	117	23
	女	789	(4.5%)	18	59	130	351	199	32
	計	1,122	(4.0%)	21	93	163	474	316	55
要治療	男	137	(1.3%)	2	21	15	55	37	7
	女	343	(1.9%)	2	13	47	190	83	8
	計	480	(1.7%)	4	34	62	245	120	15
治療中	男	1,027	(9.7%)	0	21	38	320	528	120
	女	3,017	(17.1%)	1	13	143	1,229	1,353	278
	計	4,044	(14.3%)	1	34	181	1,549	1,881	398
合 計	男	10,607		66	520	552	3,361	4,716	1,392
	女	17,672		326	1,226	1,836	6,722	5,948	1,614
	計	28,279		392	1,746	2,388	10,083	10,664	3,006

市民健診脂質検査の性年齢別平均値

検査項目	総数			35-39歳		40-49歳		50-59歳		60-69歳		70-79歳		80歳以上	
中性脂肪・ mg / dL	男	10,607	120.8	66	119.3	520	143.9	552	153.7	3,361	125.2	4,716	116.6	1,392	102.7
	女	17,672	100.4	326	71.7	1,226	74.5	1,836	94.9	6,722	103.0	5,948	105.2	1,614	103.1
	計	28,279	108.0	392	79.7	1,746	95.2	2,388	108.5	10,083	110.4	10,664	110.2	3,006	102.9
HDLコレステ ロール・ mg / dL	男	10,607	55.7	66	55.0	520	54.2	552	56.3	3,361	56.5	4,716	55.6	1,392	54.6
	女	17,672	65.3	326	66.1	1,226	68.4	1,836	69.8	6,722	65.8	5,948	63.6	1,614	61.8
	計	28,279	61.7	392	64.3	1,746	64.2	2,388	66.7	10,083	62.7	10,664	60.0	3,006	58.4
LDLコレステ ロール・ mg / dL	男	10,607	118.7	66	124.7	520	128.7	552	127.7	3,361	122.7	4,716	116.2	1,392	110.2
	女	17,672	128.3	326	106.9	1,226	115.4	1,836	133.2	6,722	134.2	5,948	126.3	1,614	119.5
	計	28,279	124.7	392	109.9	1,746	119.4	2,388	131.9	10,083	130.4	10,664	121.8	3,006	115.2

市民健診糖尿病判定結果

判 定	総数			35-39歳	40-49歳	50-59歳	60-69歳	70-79歳	80歳以上
異常なし	男	4,908	(46.2%)	54	368	297	1,457	2,089	643
	女	10,178	(57.5%)	291	1,067	1,328	3,584	3,050	858
	計	15,086	(53.3%)	345	1,435	1,625	5,041	5,139	1,501
放置可	男	2,848	(26.8%)	8	96	147	959	1,279	359
	女	4,812	(27.2%)	32	133	367	2,071	1,761	448
	計	7,660	(27.0%)	40	229	514	3,030	3,040	807
要経過観察	男	1,203	(11.3%)	1	31	52	401	552	166
	女	1,450	(8.2%)	3	17	79	594	594	163
	計	2,653	(9.4%)	4	48	131	995	1,146	329
要精密検査	男	259	(2.4%)	2	4	13	89	112	39
	女	235	(1.3%)	0	1	8	93	101	32
	計	494	(1.7%)	2	5	21	182	213	71
要治療	男	203	(1.9%)	1	10	15	75	75	27
	女	127	(0.7%)	0	2	10	50	44	21
	計	330	(1.2%)	1	12	25	125	119	48
治療中	男	1,209	(11.4%)	0	13	28	386	618	164
	女	897	(5.1%)	0	6	45	340	409	97
	計	2,106	(7.4%)	0	19	73	726	1,027	261
合計	男	10,630		66	522	552	3,367	4,725	1,398
	女	17,699		326	1,226	1,837	6,732	5,959	1,619
	計	28,329		392	1,748	2,389	10,099	10,684	3,017

市民健診糖尿病検査の性年齢別平均値

検査項目		総数		35-39歳		40-49歳		50-59歳		60-69歳		70-79歳		80歳以上	
血糖 mg / dL	男	10,607	103.1	66	92.1	520	96.3	552	100.1	3,361	102.9	4,716	103.7	1,392	105.3
	女	17,671	96.9	326	87.9	1,226	88.8	1,836	93.0	6,721	96.8	5,948	98.9	1,614	102.4
	計	28,278	99.2	392	88.6	1,746	91.1	2,388	94.7	10,082	98.9	10,664	101.0	3,006	103.7
ヘモグロビンA1c %	男	7,628	5.65	66	5.28	511	5.42	548	5.55	3,308	5.67	3,041	5.68	154	5.69
	女	11,745	5.57	305	5.22	634	5.29	1,042	5.47	5,848	5.58	3,731	5.64	185	5.69
	計	19,373	5.60	371	5.23	1,145	5.35	1,590	5.50	9,156	5.62	6,772	5.66	339	5.69

HbA1cは選択検査として実施されている

市民健診尿糖の性年齢別陽性率

検査項目	性別	総数			35-39歳			40-49歳			50-59歳			60-69歳			70-79歳			80歳以上		
		受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率	受診者数	陽性者数	陽性率
尿糖定性	男	10,538	342	3.2	66	1	1.5	520	10	1.9	548	14	2.6	3,345	107	3.2	4,684	164	3.5	1,375	46	3.3
	女	17,596	105	0.6	323	0	—	1,221	3	0.2	1,833	8	0.4	6,714	46	0.7	5,915	35	0.6	1,590	13	0.8
	計	28,134	447	1.6	389	1	0.3	1,741	13	0.7	2,381	22	0.9	10,059	153	1.5	10,599	199	1.9	2,965	59	2.0

陽性率%

市民健診貧血・血球数判定結果

判 定	総数			35-39歳	40-49歳	50-59歳	60-69歳	70-74歳
異常なし	男	95	(65.5%)	0	15	14	34	32
	女	907	(79.6%)	38	139	218	390	122
	計	1,002	(78.0%)	38	154	232	424	154
放置可	男	8	(5.5%)	0	0	0	5	3
	女	78	(6.8%)	4	11	24	25	14
	計	86	(6.7%)	4	11	24	30	17
要経過観察	男	26	(17.9%)	1	0	0	9	16
	女	104	(9.1%)	2	51	12	28	11
	計	130	(10.1%)	3	51	12	37	27
要精密検査	男	14	(9.7%)	0	0	1	5	8
	女	36	(3.2%)	7	18	3	4	4
	計	50	(3.9%)	7	18	4	9	12
要治療	男	2	(1.4%)	0	2	0	0	0
	女	7	(0.6%)	1	4	2	0	0
	計	9	(0.7%)	1	6	2	0	0
治療中	男	0	(0.0%)	0	0	0	0	0
	女	8	(0.7%)	0	3	1	4	0
	計	8	(0.6%)	0	3	1	4	0
合 計	男	145		1	17	15	53	59
	女	1,140		52	226	260	451	151
	計	1,285		53	243	275	504	210

市民健診貧血・血球数検査の性年齢別平均値

検査項目	総数			35-39歳		40-49歳		50-59歳		60-69歳		70-74歳	
赤血球数 万 / μL	男	145	460.0	1	423.0	17	480.2	15	491.3	53	460.6	59	446.3
	女	1,140	436.5	52	437.1	226	438.7	260	442.2	451	435.5	151	425.8
	計	1,285	439.1	53	436.8	243	441.6	275	444.9	504	438.2	210	431.5
ヘモグロビン g / dL	男	145	14.47	1	13.40	17	14.82	15	15.27	53	14.55	59	14.11
	女	1,140	12.96	52	12.33	226	12.24	260	13.16	451	13.26	151	13.04
	計	1,285	13.13	53	12.35	243	12.42	275	13.28	504	13.39	210	13.34
ヘマトクリット %	男	145	43.15	1	39.00	17	44.24	15	45.45	53	43.40	59	42.09
	女	1,140	39.52	52	37.91	226	37.94	260	40.07	451	40.14	151	39.65
	計	1,285	39.93	53	37.93	243	38.38	275	40.37	504	40.48	210	40.33

市民健診心電図判定結果

判 定	総数			40-49歳	50-59歳	60-69歳	70-74歳
異常なし	男	100	(60.6%)	10	9	49	32
	女	42	(68.9%)	2	3	26	11
	計	142	(62.8%)	12	12	75	43
放置可	男	8	(4.8%)	0	0	2	6
	女	4	(6.6%)	0	1	2	1
	計	12	(5.3%)	0	1	4	7
要経過観察	男	45	(27.3%)	1	1	20	23
	女	11	(18.0%)	0	2	9	0
	計	56	(24.8%)	1	3	29	23
要精密検査	男	6	(3.6%)	0	1	1	4
	女	1	(1.6%)	0	0	0	1
	計	7	(3.1%)	0	1	1	5
要治療	男	0	(0.0%)	0	0	0	0
	女	0	(0.0%)	0	0	0	0
	計	0	(0.0%)	0	0	0	0
治療中	男	6	(3.6%)	0	0	3	3
	女	3	(4.9%)	0	0	0	3
	計	9	(4.0%)	0	0	3	6
合 計	男	165		11	11	75	68
	女	61		2	6	37	16
	計	226		13	17	112	84

市民健診安静心電図検査所見

所 見	総数			40-49歳	50-59歳	60-69歳	70-74歳	所 見	総数			40-49歳	50-59歳	60-69歳	70-74歳
洞性徐脈	男	6	(3.6%)	0	0	2	4	左室肥大疑い	男	12	(7.3%)	0	0	7	5
	女	0	(0.0%)	0	0	0	0		女	3	(4.9%)	0	0	1	2
	計	6	(2.7%)	0	0	2	4		計	15	(6.6%)	0	0	8	7
左軸偏位 (-30° 以下)	男	5	(3.0%)	0	0	2	3	陳旧性心筋梗塞疑い	男	2	(1.2%)	0	0	0	2
	女	2	(3.3%)	0	0	2	0		女	0	(0.0%)	0	0	0	0
	計	7	(3.1%)	0	0	4	3		計	2	(0.9%)	0	0	0	2
第一度房室ブロック	男	12	(7.3%)	0	2	4	6	S T 低下	男	3	(1.8%)	0	0	1	2
	女	1	(1.6%)	0	1	0	0		女	0	(0.0%)	0	0	0	0
	計	13	(5.8%)	0	3	4	6		計	3	(1.3%)	0	0	1	2
不完全右脚ブロック	男	1	(0.6%)	0	0	1	0	S T - T 異常	男	2	(1.2%)	0	0	0	2
	女	1	(1.6%)	0	0	0	1		女	2	(3.3%)	0	0	1	1
	計	2	(0.9%)	0	0	1	1		計	4	(1.8%)	0	0	1	3
完全右脚ブロック	男	11	(6.7%)	0	0	5	6	異常Q波の疑い	男	5	(3.0%)	0	0	3	2
	女	1	(1.6%)	0	0	1	0		女	2	(3.3%)	0	0	2	0
	計	12	(5.3%)	0	0	6	6		計	7	(3.1%)	0	0	5	2
完全左脚ブロック	男	0	(0.0%)	0	0	0	0	陰性T波	男	4	(2.4%)	0	0	1	3
	女	1	(1.6%)	0	0	0	1		女	5	(8.2%)	0	1	4	0
	計	1	(0.4%)	0	0	0	1		計	9	(4.0%)	0	1	5	3
上室性期外収縮	男	2	(1.2%)	0	0	1	1	T波平低化	男	3	(1.8%)	0	0	1	2
	女	0	(0.0%)	0	0	0	0		女	3	(4.9%)	0	1	2	0
	計	2	(0.9%)	0	0	1	1		計	6	(2.7%)	0	1	3	2
心室性期外収縮	男	4	(2.4%)	0	1	0	3	S T 上昇	男	2	(1.2%)	1	0	1	0
	女	0	(0.0%)	0	0	0	0		女	0	(0.0%)	0	0	0	0
	計	4	(1.8%)	0	1	0	3		計	2	(0.9%)	1	0	1	0
心房細動	男	3	(1.8%)	0	0	3	0	Q T 延長	男	1	(0.6%)	0	0	1	0
	女	0	(0.0%)	0	0	0	0		女	0	(0.0%)	0	0	0	0
	計	3	(1.3%)	0	0	3	0		計	1	(0.4%)	0	0	1	0
左房負荷疑い	男	1	(0.6%)	0	0	0	1	検診者総数	男	165		11	11	75	68
	女	0	(0.0%)	0	0	0	0		女	61		2	6	37	16
	計	1	(0.4%)	0	0	0	1		計	226		13	17	112	84

市民健診眼底写真判定結果

判 定	総数			40-49歳	50-59歳	60-69歳	70-74歳
異常なし	男	63	(38.2%)	9	8	27	19
	女	25	(39.7%)	0	1	17	7
	計	88	(38.6%)	9	9	44	26
放置可	男	5	(3.0%)	0	2	3	0
	女	1	(1.6%)	0	1	0	0
	計	6	(2.6%)	0	3	3	0
要経過観察	男	80	(48.5%)	2	2	37	39
	女	32	(50.8%)	2	3	17	10
	計	112	(49.1%)	4	5	54	49
要精密検査	男	16	(9.7%)	0	0	8	8
	女	4	(6.3%)	0	1	2	1
	計	20	(8.8%)	0	1	10	9
要治療	男	0	(0.0%)	0	0	0	0
	女	0	(0.0%)	0	0	0	0
	計	0	(0.0%)	0	0	0	0
治療中	男	1	(0.6%)	0	0	0	1
	女	1	(1.6%)	0	0	1	0
	計	2	(0.9%)	0	0	1	1
合 計	男	165		11	12	75	67
	女	63		2	6	37	18
	計	228		13	18	112	85

市民健診眼底検査所見

所 見	総数			40-49歳	50-59歳	60-69歳	70-74歳
細動脈狭細化	男	7	(4.2%)	1	0	3	3
	女	4	(6.3%)	0	0	4	0
	計	11	(4.8%)	1	0	7	3
細動脈口径不同	男	7	(4.2%)	0	0	5	2
	女	3	(4.8%)	1	0	1	1
	計	10	(4.4%)	1	0	6	3
眼底出血	男	3	(1.8%)	0	0	1	2
	女	0	(0.0%)	0	0	0	0
	計	3	(1.3%)	0	0	1	2
細動脈反射増強	男	6	(3.6%)	0	0	4	2
	女	2	(3.2%)	0	1	1	0
	計	8	(3.5%)	0	1	5	2
動静脈交叉現象	男	30	(18.2%)	1	2	15	12
	女	15	(23.8%)	1	3	8	3
	計	45	(19.7%)	2	5	23	15
点状出血	男	1	(0.6%)	0	0	1	0
	女	0	(0.0%)	0	0	0	0
	計	1	(0.4%)	0	0	1	0
斑状出血	男	1	(0.6%)	0	0	1	0
	女	0	(0.0%)	0	0	0	0
	計	1	(0.4%)	0	0	1	0
網膜白斑	男	13	(7.9%)	0	0	9	4
	女	2	(3.2%)	0	0	2	0
	計	15	(6.6%)	0	0	11	4
白内障の疑い	男	41	(24.8%)	0	0	13	28
	女	13	(20.6%)	0	1	6	6
	計	54	(23.7%)	0	1	19	34
視神経乳頭異常の疑い	男	3	(1.8%)	0	0	2	1
	女	0	(0.0%)	0	0	0	0
	計	3	(1.3%)	0	0	2	1
網脈絡膜瘢痕巣の疑い	男	1	(0.6%)	0	1	0	0
	女	0	(0.0%)	0	0	0	0
	計	1	(0.4%)	0	1	0	0
瞳孔の開き不十分	男	6	(3.6%)	0	0	3	3
	女	1	(1.6%)	0	0	1	0
	計	7	(3.1%)	0	0	4	3
その他の所見	男	8	(4.8%)	0	1	6	1
	女	4	(6.3%)	0	1	2	1
	計	12	(5.3%)	0	2	8	2
検診者総数	男	165		11	12	75	67
	女	63		2	6	37	18
	計	228		13	18	112	85

結核・肺がん検診判定結果

判 定		総数		40-49歳	50-59歳	60-69歳	70-79歳	80歳以上
異常なし		男	6,876 (68.5%)	425	440	2,460	2,921	630
		女	12,612 (76.2%)	1,080	1,503	5,177	4,027	825
		計	19,488 (73.3%)	1,505	1,943	7,637	6,948	1,455
異常所見あるも 精査不要		男	2,721 (27.1%)	54	76	626	1,369	596
		女	3,456 (20.9%)	74	240	1126	1,395	621
		計	6,177 (23.2%)	128	316	1,752	2,764	1,217
要精密検査	結核の疑い	男	0 (0.0%)	0	0	0	0	0
		女	0 (0.0%)	0	0	0	0	0
		計	0 (0.0%)	0	0	0	0	0
	結核以外の疑い	男	286 (2.8%)	26	14	110	106	30
		女	268 (1.6%)	21	25	99	101	22
		計	554 (2.1%)	47	39	209	207	52
	肺がんの疑い	男	154 (1.5%)	2	7	41	70	34
		女	208 (1.3%)	4	21	73	80	30
		計	362 (1.4%)	6	28	114	150	64
合計		男	10,037	507	537	3,237	4,466	1,290
		女	16,544	1,179	1,789	6,475	5,603	1,498
		計	26,581	1,686	2,326	9,712	10,069	2,788

肺がん検診喀痰細胞診検査所見

所 見	総数		40-49歳	50-59歳	60-69歳	70-79歳	80歳以上
B 1 (C l a s s I (異常なし))	男	29	(12.4%)	0	1	17	9
	女	2	(14.3%)	0	0	1	0
	計	31	(12.5%)	0	1	18	2
B 2 (C l a s s II (悪性所見なし))	男	204	(87.2%)	2	11	91	86
	女	12	(85.7%)	0	2	6	2
	計	216	(87.1%)	2	13	97	16
E 2 (C l a s s V)	男	1	(0.4%)	0	0	0	0
	女	0	(0.0%)	0	0	0	0
	計	1	(0.4%)	0	0	0	1
検診者総数	男	234		2	12	108	95
	女	14		0	2	7	3
	計	248		2	14	115	98

結核・肺がん胸部X線検査所見

所 見	総数				40-49歳	50-59歳	60-69歳	70-79歳	80歳以上	所 見	総数				40-49歳	50-59歳	60-69歳	70-79歳	80歳以上	
心陰影拡大	男	149	(1.5%)	7	5	31	66	40	胸膜癒着	男	474	(4.7%)	8	12	93	236	125			
	女	176	(1.1%)	1	7	49	81	38		女	374	(2.3%)	14	29	128	132	71			
	計	325	(1.2%)	8	12	80	147	78		計	848	(3.2%)	22	41	221	368	196			
肺門陰影拡大	男	50	(0.5%)	6	2	20	11	11	胸水陰影	男	27	(0.3%)	0	0	11	7	9			
	女	68	(0.4%)	1	1	19	23	24		女	20	(0.1%)	0	0	3	13	4			
	計	118	(0.4%)	7	3	39	34	35		計	47	(0.2%)	0	0	14	20	13			
肺紋理増強	男	13	(0.1%)	0	1	2	6	4	上葉縮小	男	31	(0.3%)	1	1	6	18	5			
	女	7	(0.0%)	0	0	1	5	1		女	20	(0.1%)	1	2	6	7	4			
	計	20	(0.1%)	0	1	3	11	5		計	51	(0.2%)	2	3	12	25	9			
肺紋理減少	男	2	(0.0%)	0	0	2	0	0	横隔膜高位	男	171	(1.7%)	12	3	33	78	45			
	女	0	(0.0%)	0	0	0	0	0		女	122	(0.7%)	0	3	38	50	31			
	計	2	(0.0%)	0	0	2	0	0		計	293	(1.1%)	12	6	71	128	76			
線状陰影	男	330	(3.3%)	16	17	80	145	72	胸郭形成	男	11	(0.1%)	0	0	1	4	6			
	女	345	(2.1%)	16	8	124	146	51		女	8	(0.0%)	0	0	0	5	3			
	計	675	(2.5%)	32	25	204	291	123		計	19	(0.1%)	0	0	1	9	9			
帯状陰影	男	105	(1.0%)	2	4	40	42	17	陈旧性肺結核の疑い	男	34	(0.3%)	1	1	6	14	12			
	女	124	(0.7%)	2	6	41	48	27		女	30	(0.2%)	0	1	4	21	4			
	計	229	(0.9%)	4	10	81	90	44		計	64	(0.2%)	1	2	10	35	16			
大動脈蛇行	男	65	(0.6%)	0	2	10	38	15	肺気腫の疑い	男	220	(2.2%)	6	6	57	102	49			
	女	144	(0.9%)	0	0	18	68	58		女	55	(0.3%)	4	11	28	8	4			
	計	209	(0.8%)	0	2	28	106	73		計	275	(1.0%)	10	17	85	110	53			
微細斑点状陰影	男	161	(1.6%)	2	3	32	80	44	慢性気管支炎の疑い	男	12	(0.1%)	1	1	3	7	0			
	女	221	(1.3%)	5	24	64	90	38		女	5	(0.0%)	0	0	2	2	1			
	計	382	(1.4%)	7	27	96	170	82		計	17	(0.1%)	1	1	5	9	1			
心臓手術後	男	27	(0.3%)	0	0	6	17	4	肺炎の疑い	男	2	(0.0%)	0	0	0	1	1			
	女	19	(0.1%)	0	0	5	11	3		女	2	(0.0%)	0	1	0	1	0			
	計	46	(0.2%)	0	0	11	28	7		計	4	(0.0%)	0	1	0	2	1			
粒状陰影	男	10	(0.1%)	0	0	3	3	4	気管支拡張症の疑い	男	17	(0.2%)	0	0	2	10	5			
	女	12	(0.1%)	1	0	4	3	4		女	28	(0.2%)	0	1	13	12	2			
	計	22	(0.1%)	1	0	7	6	8		計	45	(0.2%)	0	1	15	22	7			
結節陰影	男	573	(5.7%)	9	18	152	291	103	肺囊胞症の疑い	男	1	(0.0%)	0	0	0	1	0			
	女	771	(4.7%)	32	53	256	337	93		女	0	(0.0%)	0	0	0	0	0			
	計	1,344	(5.1%)	41	71	408	628	196		計	1	(0.0%)	0	0	0	1	0			
円形陰影	男	20	(0.2%)	0	0	6	12	2	間質性肺炎の疑い	男	152	(1.5%)	0	0	32	71	49			
	女	37	(0.2%)	2	4	13	17	1		女	42	(0.3%)	0	0	4	17	21			
	計	57	(0.2%)	2	4	19	29	3		計	194	(0.7%)	0	0	36	88	70			
塊状陰影	男	38	(0.4%)	0	0	12	19	7	塵肺の疑い	男	1	(0.0%)	0	0	1	0	0			
	女	57	(0.3%)	1	5	16	22	13		女	1	(0.0%)	0	0	1	0	0			
	計	95	(0.4%)	1	5	28	41	20		計	2	(0.0%)	0	0	2	0	0			
浸潤陰影	男	279	(2.8%)	6	8	65	139	61	胸椎側弯	男	93	(0.9%)	2	5	13	44	29			
	女	299	(1.8%)	3	11	104	130	51		女	414	(2.5%)	10	32	110	160	102			
	計	578	(2.2%)	9	19	169	269	112		計	507	(1.9%)	12	37	123	204	131			
網目状陰影	男	153	(1.5%)	1	0	36	76	40	キライディティ症候群	男	19	(0.2%)	0	0	4	10	5			
	女	45	(0.3%)	1	3	9	22	10		女	3	(0.0%)	0	0	0	3	0			
	計	198	(0.7%)	2	3	45	98	50		計	22	(0.1%)	0	0	4	13	5			
石灰化陰影	男	426	(4.2%)	4	5	70	226	121	縦隔腫大	男	8	(0.1%)	0	1	3	3	1			
	女	514	(3.1%)	3	15	140	241	115		女	5	(0.0%)	0	0	2	3	0			
	計	940	(3.5%)	7	20	210	467	236		計	13	(0.0%)	0	1	5	6	1			
空洞陰影	男	8	(0.1%)	1	0	3	3	1	横隔膜ヘルニア	男	0	(0.0%)	0	0	0	0	0			
	女	6	(0.0%)	1	0	3	1	1		女	1	(0.0%)	0	0	0	0	1			
	計	14	(0.1%)	2	0	6	4	2		計	1	(0.0%)	0	0	0	0	1			
嚢胞陰影	男	75	(0.7%)	5	4	25	32	9	食道裂孔ヘルニア	男	16	(0.2%)	0	0	3	7	6			
	女	17	(0.1%)	1	2	6	8	0		女	86	(0.5%)	0	1	11	45	29			
	計	92	(0.3%)	6	6	31	40	9		計	102	(0.4%)	0	1	14	52	35			
無気肺陰影	男	10	(0.1%)	0	0	4	3	3	右大動脈弓	男	9	(0.1%)	0	0	2	6	1			
	女	12	(0.1%)	0	1	2	8	1		女	5	(0.0%)	0	1	0	3	1			
	計	22	(0.1%)	0	1	6	11	4		計	14	(0.1%)	0	1	2	9	2			
気腫様陰影	男	8	(0.1%)	0	0	4	4	0	肺手術後	男	19	(0.2%)	0	0	4	10	5			
	女	0	(0.0%)	0	0	0	0	0		女	19	(0.1%)	0	0	8	8	3			
	計	8	(0.0%)	0	0	4	4	0		計	38	(0.1%)	0	0	12	18	8			
胸膜肥厚	男	678	(6.8%)	17	23	177	319	142	その他の所見	男	110	(1.1%)	1	3	27	64	15			
	女	832	(5.0%)	16	98	349	256	113		女	118	(0.7%)	3	11	44	48	12			
	計	1,510	(5.7%)	33	121	526	575	255		計	228	(0.9%)	4	14	71	112	27			
										検診者総数	男	10,037		507	537	3,237	4,466	1,290		
											女	16,544		1,179	1,789	6,475	5,603	1,498		
											計	26,581		1,686	2,326	9,712	10,069	2,788		

前立腺がん検診判定結果

判 定	総数		50-59歳	60-69歳	70-79歳	80歳以上
異常なし	4,934	(88.5%)	277	1,748	2,388	521
放置可	0	(0.0%)	0	0	0	0
要経過観察	0	(0.0%)	0	0	0	0
要精密検査	344	(6.2%)	3	90	190	61
要治療	0	(0.0%)	0	0	0	0
治療中	297	(5.3%)	3	64	172	58
合計	5,575		283	1,902	2,750	640

前立腺がん検診P S Aの年齢別平均値

検査項目	総数		50-59歳		60-69歳		70-79歳		80歳以上	
P S A n g / m L	5,576	1.80	283	1.06	1,902	1.57	2,749	1.88	640	2.48

長寿スマイルチェック結果

判 定	総数		65-69歳	70-79歳	80歳以上
該 当	男	6,843 (92.1%)	2038	3914	891
	女	8,570 (87.8%)	3327	4467	776
	計	15,413 (89.7%)	5365	8,381	1667
非該 当	男	586 (7.9%)	111	290	185
	女	1,187 (12.2%)	210	642	335
	計	1,773 (10.3%)	321	932	520
合 計	男	7,429	2,149	4,204	1,076
	女	9,757	3,537	5,109	1,111
	計	17,186	5,686	9,313	2,187

長寿スマイルチェック基本チェックリスト

項目名称	総数		65-69歳	70-79歳	80歳以上
運動機能	男	310 (4.2%)	37	139	134
	女	786 (8.1%)	106	403	277
	計	1,096 (6.4%)	143	542	411
栄養状態	男	13 (0.2%)	1	5	7
	女	22 (0.2%)	4	12	6
	計	35 (0.2%)	5	17	13
口腔機能	男	360 (4.8%)	79	194	87
	女	562 (5.8%)	128	322	112
	計	922 (5.4%)	207	516	199
閉じこもり傾向	男	164 (2.2%)	27	70	67
	女	406 (4.2%)	60	201	145
	計	570 (3.3%)	87	271	212
ものわすれ	男	305 (4.1%)	54	154	97
	女	465 (4.8%)	94	242	129
	計	770 (4.5%)	148	396	226
うつ傾向	男	223 (3.0%)	42	106	75
	女	427 (4.4%)	68	224	135
	計	650 (3.8%)	110	330	210
検診者総数	男	7,429	2,149	4,204	1,076
	女	9,757	3,537	5,109	1,111
	計	17,186	5,686	9,313	2,187

児童・生徒を対象とした健診

児童・生徒の腎臓健診受診者の尿所見

検査項目	性別	総数		
		受診者数	陽性者数	陽性率
尿糖	男	22,237	22	0.1%
	女	21,623	15	0.1%
	計	43,860	37	0.1%
尿蛋白	男	22,237	321	1.4%
	女	21,623	278	1.3%
	計	43,860	599	1.4%
尿潜血	男	22,237	210	0.9%
	女	21,623	657	3.0%
	計	43,860	867	2.0%

児童・生徒の心臓検査所見（経過観察者を除く）

判定		人数	
心臓の異常	心房中隔欠損	16	0.1%
	心室中隔欠損	33	0.2%
	肺動脈狭窄	8	0.0%
	動脈管開存	5	0.0%
	ファロー四徴	0	0.0%
	先天性心疾患（位置異常）	0	0.0%
	先天性心疾患（その他）	16	0.1%
	大動脈弁疾患	2	0.0%
	僧帽弁疾患	8	0.0%
	三尖弁疾患	12	0.1%
	心筋症	0	0.0%
	虚血性心疾患	0	0.0%
	心臓手術後	78	0.5%
	その他	6	0.0%
計		184	1.1%
その他の異常	胸郭変形	0	0.0%
	先天性心疾患の自然閉鎖	44	0.3%
	川崎病後	187	1.1%
	高血圧	0	0.0%
	その他	5	0.0%
	計	236	1.4%
受診者数合計		16,689	

児童・生徒のすこやか検診（血液検査）結果

	性別	肥満	血清・脂質				肝機能	貧血			白血球		血小板
		肥満度	総コレステロール	中性脂肪	H D L コレステロール	動脈硬化	G P T	赤血球数	ヘモグロビン	ヘマトクリット	白血球数		血小板数
											4.3未満	15.1以上	
受診者数	男	1,966	1,966	1,966	1,966	1,963	1,966	1,966	1,966	1,966	1,966	1,966	1,965
	女	1,869	1,868	1,868	1,868	1,867	1,868	1,869	1,869	1,869	1,869	1,869	1,869
	計	3,835	3,834	3,834	3,834	3,830	3,834	3,835	3,835	3,835	3,835	3,835	3,834
要経過観察者数 (%)	男	173 (8.8)	240 (12.2)	43 (2.2)	91 (4.6)	- (1.5)	29 (0.5)	9 (1.1)	22 (0.7)	14 (0.7)	-	-	-
		137 (7.3)	216 (11.6)	49 (2.6)	113 (6.0)	- (0.9)	16 (0.4)	8 (1.4)	27 (0.6)	11 (0.6)	-	-	-
	女	310 (8.1)	456 (11.9)	92 (2.4)	204 (5.3)	- (1.2)	45 (0.4)	17 (0.4)	49 (1.3)	25 (0.7)	-	-	-
	計	47 (2.4)	52 (2.6)	23 (1.2)	0 (0.0)	22 (1.1)	39 (2.0)	0 (0.0)	4 (0.2)	3 (0.2)	9 (0.5)	4 (0.2)	3 (0.2)
		30 (1.6)	50 (2.7)	12 (0.6)	0 (0.0)	28 (1.5)	5 (0.3)	0 (0.0)	1 (0.1)	2 (0.1)	19 (1.0)	2 (0.1)	3 (0.2)
要精密検査数 (%)	計	77 (2.0)	102 (2.7)	35 (0.9)	0 (0.0)	50 (1.3)	44 (1.1)	0 (0.0)	5 (0.1)	5 (0.1)	28 (0.7)	6 (0.2)	6 (0.2)

判定		人数	
心電図の異常	右室肥大	0	0.0%
	左室肥大	1	0.0%
	心筋障害	1	0.0%
	洞性頻脈	0	0.0%
	洞性徐脈	69	0.4%
	上室性期外収縮	37	0.2%
	心室性期外収縮	41	0.2%
	上室性・心室性期外収縮	2	0.0%
	第1度房室ブロック	14	0.1%
	第2度房室ブロック	3	0.0%
	房室解離	0	0.0%
	完全ブロック	20	0.1%
	不完全右脚ブロック	8	0.0%
	ヘミブロック	0	0.0%
	刺激発生異常	256	1.5%
	W P W 症候群	21	0.1%
	P Q 短縮・L G L 症候群	1	0.0%
	Q T 延長症候群	5	0.0%
	その他	14	0.1%
	計	493	3.0%
X線検査の異常心拡大		4	0.0%

（経過観察者を除く）