

上部消化管検査判定結果

判定	総数		年齢							
	男	女	19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上	
異常なし	男	14,924 (82.2%)	11	332	2,915	4,938	4,331	2,077	320	
	女	8,353 (78.8%)	0	88	1,540	2,891	2,457	1,194	183	
	計	23,277 (80.9%)	11	420	4,455	7,829	6,788	3,271	503	
放置可	男	951 (5.2%)	0	2	99	302	351	168	29	
	女	552 (5.2%)	0	1	58	214	191	77	11	
	計	1,503 (5.2%)	0	3	157	516	542	245	40	
要経過観察	男	1,391 (7.7%)	0	3	152	380	516	284	56	
	女	1,226 (11.6%)	0	2	146	438	428	182	30	
	計	2,617 (9.1%)	0	5	298	818	944	466	86	
要精密検査	男	821 (4.5%)	0	6	87	222	238	210	58	
	女	456 (4.3%)	0	5	54	150	134	88	25	
	計	1,277 (4.4%)	0	11	141	372	372	298	83	
要治療	男	33 (0.2%)	0	0	3	12	14	4	0	
	女	13 (0.1%)	0	0	4	2	6	0	1	
	計	46 (0.2%)	0	0	7	14	20	4	1	
治療中	男	38 (0.2%)	0	0	2	8	16	9	3	
	女	4 (0.0%)	0	0	0	0	2	2	0	
	計	42 (0.1%)	0	0	2	8	18	11	3	
合計	男	18,158	11	343	3,258	5,862	5,466	2,752	466	
	女	10,604	0	96	1,802	3,695	3,218	1,543	250	
	計	28,762	11	439	5,060	9,557	8,684	4,295	716	

胃内視鏡検査所見

所見	総数	年齢							
		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上	
十二指腸潰瘍	男	8 (0.5%)	0	0	2	4	2	0	0
	女	5 (0.6%)	0	0	0	2	3	0	0
	計	13 (0.5%)	0	0	2	6	5	0	0
慢性胃炎	男	248 (15.3%)	0	0	15	77	120	30	6
	女	99 (11.0%)	0	0	10	30	45	12	2
	計	347 (13.8%)	0	0	25	107	165	42	8
食道裂孔ヘルニア	男	355 (22.0%)	0	1	34	113	147	55	5
	女	83 (9.3%)	0	0	15	33	28	5	2
	計	438 (17.4%)	0	1	49	146	175	60	7
十二指腸潰瘍瘢痕	男	77 (4.8%)	0	0	5	20	35	17	0
	女	32 (3.6%)	0	0	1	11	13	7	0
	計	109 (4.3%)	0	0	6	31	48	24	0
萎縮性胃炎	男	295 (18.3%)	0	1	16	60	125	83	10
	女	135 (15.1%)	0	0	8	43	48	31	5
	計	430 (17.1%)	0	1	24	103	173	114	15
逆流性食道炎	男	230 (14.2%)	0	0	23	80	93	31	3
	女	21 (2.3%)	0	0	2	8	9	2	0
	計	251 (10.0%)	0	0	25	88	102	33	3
出血性胃炎	男	6 (0.4%)	0	0	0	2	2	2	0
	女	1 (0.1%)	0	0	0	0	1	0	0
	計	7 (0.3%)	0	0	0	2	3	2	0
慢性十二指腸炎	男	7 (0.4%)	0	0	1	3	3	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	7 (0.3%)	0	0	1	3	3	0	0
びらん性胃炎	男	141 (8.7%)	0	0	19	46	51	23	2
	女	57 (6.4%)	0	1	5	24	18	9	0
	計	198 (7.9%)	0	1	24	70	69	32	2
タコイボびらん	男	19 (1.2%)	0	0	2	5	8	3	1
	女	7 (0.8%)	0	0	0	0	5	2	0
	計	26 (1.0%)	0	0	2	5	13	5	1
胃憩室	男	0 (0.0%)	0	0	0	0	0	0	0
	女	2 (0.2%)	0	0	1	0	1	0	0
	計	2 (0.1%)	0	0	1	0	1	0	0
胃潰瘍	男	5 (0.3%)	0	0	0	0	5	0	0
	女	6 (0.7%)	0	0	2	0	2	1	1
	計	11 (0.4%)	0	0	2	0	7	1	1
食道潰瘍	男	1 (0.1%)	0	0	0	0	0	1	0
	女	1 (0.1%)	0	0	0	0	0	1	0
	計	2 (0.1%)	0	0	0	0	0	2	0
胃潰瘍瘢痕	男	48 (3.0%)	0	0	1	5	20	19	3
	女	14 (1.6%)	0	0	1	3	6	4	0
	計	62 (2.5%)	0	0	2	8	26	23	3
胃ポリープ	男	235 (14.5%)	0	0	24	73	102	31	5
	女	265 (29.5%)	0	0	42	113	83	24	3
	計	500 (19.9%)	0	0	66	186	185	55	8
粘膜下腫瘍	男	38 (2.4%)	0	0	2	12	18	5	1
	女	15 (1.7%)	0	0	2	2	8	3	0
	計	53 (2.1%)	0	0	4	14	26	8	1
その他の所見 (十二指腸所見)	男	86 (5.3%)	0	0	4	33	33	16	0
	女	14 (1.6%)	0	1	3	6	4	0	0
	計	100 (4.0%)	0	1	7	39	37	16	0
その他の所見 (胃所見)	男	185 (11.4%)	0	0	18	67	68	26	6
	女	97 (10.8%)	0	1	12	38	31	13	2
	計	282 (11.2%)	0	1	30	105	99	39	8
その他の所見 (食道所見)	男	83 (5.1%)	0	0	5	25	38	12	3
	女	24 (2.7%)	0	1	3	9	8	2	1
	計	107 (4.3%)	0	1	8	34	46	14	4
検診者総数	男	1,616	0	7	205	540	590	245	29
	女	897	0	7	152	340	284	103	11
	計	2,513	0	14	357	880	874	348	40

胃部X線検査所見

所見	総数	年齢								所見	総数	年齢								
		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上	19歳以下			20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上			
胃 部	総数									狭窄	総数									
伸展不良	男	24 (0.2%)	0	1	1	10	6	5	1	狭窄	男	3 (0.0%)	0	0	0	0	0	2	1	
	女	10 (0.1%)	0	0	0	4	2	1	3		女	1 (0.0%)	0	0	0	1	0	0	0	
	計	34 (0.2%)	0	1	1	14	8	6	4		計	4 (0.0%)	0	0	0	1	0	2	1	
壁硬化	男	3 (0.0%)	0	0	0	0	2	1	0	食道裂孔ヘルニア	男	26 (0.2%)	0	0	2	6	10	6	2	
	女	3 (0.0%)	0	0	0	0	0	2	1		女	14 (0.2%)	0	0	0	0	2	7	5	
	計	6 (0.0%)	0	0	0	0	2	3	1		計	40 (0.2%)	0	0	2	6	12	13	7	
辺縁不整	男	5 (0.0%)	0	0	0	1	1	1	2	食道炎の疑い	男	1 (0.0%)	0	0	0	0	0	0	1	
	女	2 (0.0%)	0	0	0	1	1	0	0		女	0 (0.0%)	0	0	0	0	0	0	0	
	計	7 (0.0%)	0	0	0	2	2	1	2		計	1 (0.0%)	0	0	0	0	0	0	1	
変形	男	217 (1.7%)	0	0	19	60	67	59	12	その他の所見	男	28 (0.2%)	0	0	2	4	8	8	6	
	女	45 (0.5%)	0	0	2	18	12	10	3		女	20 (0.2%)	0	0	1	3	7	5	4	
	計	262 (1.2%)	0	0	21	78	79	69	15		計	48 (0.2%)	0	0	3	7	15	13	10	
粘膜不整	男	76 (0.6%)	0	0	12	20	19	17	8	十二指腸部										
	女	40 (0.5%)	0	0	3	8	10	14	5		十二指腸球部変形	男	238 (1.9%)	0	1	25	54	85	58	15
	計	116 (0.6%)	0	0	15	28	29	31	13			女	51 (0.6%)	0	0	2	16	18	14	1
ニッシュ	男	7 (0.1%)	0	0	1	2	3	1	0	十二指腸潰瘍の疑い		男	59 (0.5%)	0	0	10	17	13	17	2
	女	4 (0.0%)	0	1	1	0	0	1	1		女	18 (0.2%)	0	0	0	10	8	0	0	
	計	11 (0.1%)	0	1	2	2	3	2	1		計	77 (0.4%)	0	0	10	27	21	17	2	
レリーフ集中像	男	69 (0.5%)	0	0	7	17	16	23	6	十二指腸潰瘍瘢痕の疑い	男	17 (0.1%)	0	0	1	6	9	1	0	
	女	21 (0.2%)	0	0	4	3	7	7	0		女	1 (0.0%)	0	0	0	0	1	0	0	
	計	90 (0.4%)	0	0	11	20	23	30	6		計	18 (0.1%)	0	0	1	6	10	1	0	
レリーフ異常	男	0 (0.0%)	0	0	0	0	0	0	0	十二指腸炎の疑い	男	1 (0.0%)	0	0	1	0	0	0	0	
	女	2 (0.0%)	0	0	0	1	0	1	0		女	1 (0.0%)	0	0	0	0	1	0	0	
	計	2 (0.0%)	0	0	0	1	0	1	0		計	2 (0.0%)	0	0	1	0	1	0	0	
バリウム斑	男	5 (0.0%)	0	0	0	2	2	1	0	十二指腸憩室の疑い	男	7 (0.1%)	0	0	0	1	2	2	2	
	女	2 (0.0%)	0	0	0	1	0	1	0		女	5 (0.1%)	0	0	0	1	1	3	0	
	計	7 (0.0%)	0	0	0	3	2	2	0		計	12 (0.1%)	0	0	0	2	3	5	2	
陰影欠損	男	1 (0.0%)	0	0	0	0	0	1	0	造影不良	男	12 (0.1%)	0	0	3	4	2	3	0	
	女	1 (0.0%)	0	0	0	0	1	0	0		女	1 (0.0%)	0	0	0	1	0	0	0	
	計	2 (0.0%)	0	0	0	0	1	1	0		計	13 (0.1%)	0	0	3	5	2	3		