

動脈硬化検査判定結果

判定	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
	男	女							
異常なし	男	122 (17.9%)	0	0	11	73	30	8	0
	女	140 (23.8%)	0	0	17	71	39	13	0
	計	262 (20.6%)	0	0	28	144	69	21	0
放置可	男	77 (11.3%)	0	0	6	29	31	9	2
	女	94 (16.0%)	0	0	5	22	46	20	1
	計	171 (13.5%)	0	0	11	51	77	29	3
要経過観察	男	436 (64.1%)	0	0	9	88	162	142	35
	女	326 (55.3%)	0	0	2	36	132	129	27
	計	762 (60.0%)	0	0	11	124	294	271	62
要精密検査	男	40 (5.9%)	0	0	1	2	7	18	12
	女	29 (4.9%)	0	0	0	2	8	11	8
	計	69 (5.4%)	0	0	1	4	15	29	20
要治療	男	0 (0.0%)	0	0	0	0	0	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	0 (0.0%)	0	0	0	0	0	0	0
治療中	男	5 (0.7%)	0	0	0	0	0	4	1
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	5 (0.4%)	0	0	0	0	0	4	1
合計	男	680	0	0	27	192	230	181	50
	女	589	0	0	24	131	225	173	36
	計	1,269	0	0	51	323	455	354	86

頸部動脈超音波検査所見

所見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
	男	女							
頸動脈の内腔狭窄 (50-75%)	男	16 (2.4%)	0	0	0	1	4	6	5
	女	6 (1.0%)	0	0	0	0	2	2	2
	計	22 (1.7%)	0	0	0	1	6	8	7
頸動脈の高度狭窄 (75%以上)	男	1 (0.1%)	0	0	0	0	0	0	1
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	1 (0.1%)	0	0	0	0	0	0	1
総頸動脈内径拡大	男	2 (0.3%)	0	0	0	0	0	2	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	2 (0.2%)	0	0	0	0	0	2	0
頸動脈描出不良	男	32 (4.7%)	0	0	4	6	11	10	1
	女	16 (2.7%)	0	0	0	5	5	6	0
	計	48 (3.8%)	0	0	4	11	16	16	1
不安定プラーク (III b)	男	1 (0.1%)	0	0	0	1	0	0	0
	女	2 (0.3%)	0	0	0	1	1	0	0
	計	3 (0.2%)	0	0	0	2	1	0	0
総頸動脈壁肥厚	男	39 (5.7%)	0	0	0	2	7	19	11
	女	21 (3.6%)	0	0	0	0	5	10	6
	計	60 (4.7%)	0	0	0	2	12	29	17
その他の所見	男	7 (1.0%)	0	0	1	0	4	2	0
	女	2 (0.3%)	0	0	0	0	1	1	0
	計	9 (0.7%)	0	0	1	0	5	3	0
検診者総数	男	680	0	0	27	192	230	181	50
	女	589	0	0	24	131	225	173	36
	計	1,269	0	0	51	323	455	354	86

頸部動脈超音波検査の計測値の性年齢別平均値

検査項目	総数		29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
	男	女	平均	標準偏差	平均	標準偏差	平均	標準偏差	平均	標準偏差	平均	標準偏差	平均	標準偏差
平均IMT 右	男	680	0.730	0	27	0.580	192	0.620	230	0.700	181	0.830	50	0.950
	女	589	0.690	0	24	0.540	131	0.580	225	0.670	173	0.760	36	0.910
	計	1,269	0.710	0	51	0.560	323	0.610	455	0.690	354	0.790	86	0.930
平均IMT 左	男	680	0.780	0	27	0.590	192	0.660	230	0.760	181	0.880	50	1.000
	女	589	0.720	0	24	0.540	131	0.600	225	0.710	173	0.780	36	0.950
	計	1,269	0.750	0	51	0.570	323	0.630	455	0.740	354	0.830	86	0.980
最大IMT 右	男	680	1.370	0	27	1.050	192	1.110	230	1.310	181	1.640	50	1.880
	女	589	1.250	0	24	0.870	131	1.040	225	1.210	173	1.420	36	1.690
	計	1,269	1.320	0	51	0.960	323	1.080	455	1.260	354	1.530	86	1.800
最大IMT 左	男	680	1.390	0	27	1.000	192	1.150	230	1.360	181	1.630	50	1.840
	女	589	1.280	0	24	0.880	131	1.070	225	1.280	173	1.430	36	1.620
	計	1,269	1.340	0	51	0.940	323	1.120	455	1.320	354	1.530	86	1.750

血圧脈波検査の性年齢別平均値

検査項目	総数		29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
	男	女	平均	標準偏差	平均	標準偏差	平均	標準偏差	平均	標準偏差	平均	標準偏差	平均	標準偏差
ABI検査 右	男	674	1.150	0	27	1.130	192	1.140	230	1.160	176	1.160	49	1.150
	女	584	1.120	0	24	1.110	131	1.110	223	1.120	171	1.130	35	1.130
	計	1,258	1.140	0	51	1.120	323	1.130	453	1.140	347	1.140	84	1.140
ABI検査 左	男	674	1.140	0	27	1.110	192	1.130	230	1.150	176	1.150	49	1.130
	女	584	1.110	0	24	1.080	131	1.100	223	1.110	171	1.110	35	1.110
	計	1,258	1.130	0	51	1.100	323	1.120	453	1.130	347	1.130	84	1.120
PWV検査 右 cm/s	男	674	1414.9	0	27	1243.7	192	1273.3	230	1393.4	176	1557.3	49	1654.1
	女	584	1348.1	0	24	1125.2	131	1164.2	223	1331.7	171	1469.3	35	1702.4
	計	1,258	1383.9	0	51	1188.0	323	1229.1	453	1363.0	347	1513.9	84	1674.3
PWV検査 左 cm/s	男	674	1413.9	0	27	1263.6	192	1259.3	230	1393.9	176	1564.8	49	1654.4
	女	584	1357.4	0	24	1135.7	131	1175.9	223	1346.8	171	1477.8	35	1668.4
	計	1,258	1387.7	0	51	1203.4	323	1225.5	453	1370.7	347	1521.9	84	1660.3