

動脈硬化検査判定結果

判定	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
	男	女							
異常なし	男	113 (14.9%)	0	1	10	68	32	2	0
	女	117 (20.7%)	0	3	12	55	40	7	0
	計	230 (17.3%)	0	4	22	123	72	9	0
放置可	男	83 (10.9%)	0	0	4	31	40	7	1
	女	83 (14.7%)	0	0	0	26	44	11	2
	計	166 (12.5%)	0	0	4	57	84	18	3
要経過観察	男	499 (65.7%)	0	0	14	79	240	131	35
	女	329 (58.1%)	0	0	1	47	137	114	30
	計	828 (62.4%)	0	0	15	126	377	245	65
要精密検査	男	61 (8.0%)	0	0	1	2	24	20	14
	女	35 (6.2%)	0	0	0	1	7	19	8
	計	96 (7.2%)	0	0	1	3	31	39	22
要治療	男	0 (0.0%)	0	0	0	0	0	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	0 (0.0%)	0	0	0	0	0	0	0
治療中	男	4 (0.5%)	0	0	0	0	0	1	3
	女	2 (0.4%)	0	0	0	0	2	0	0
	計	6 (0.5%)	0	0	0	0	2	1	3
合計	男	760	0	1	29	180	336	161	53
	女	566	0	3	13	129	230	151	40
	計	1,326	0	4	42	309	566	312	93

頸部動脈超音波検査所見

所見	総数		19歳以下	20-29歳	30-39歳	40-49歳	50-59歳	60-69歳	70歳以上
	男	女							
プラーク個数	男	760 (100.0%)	0	1	29	180	336	161	53
	女	566 (100.0%)	0	3	13	129	230	151	40
	計	1,326 (100.0%)	0	4	42	309	566	312	93
頸動脈の内腔狭窄 (50-75%)	男	16 (2.1%)	0	0	0	1	5	6	4
	女	4 (0.7%)	0	0	0	0	3	1	0
	計	20 (1.5%)	0	0	0	1	8	7	4
総頸動脈内径拡大	男	3 (0.4%)	0	0	0	0	0	2	1
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	3 (0.2%)	0	0	0	0	0	2	1
頸動脈描出不良	男	21 (2.8%)	0	0	0	4	14	3	0
	女	6 (1.1%)	0	0	0	1	4	1	0
	計	27 (2.0%)	0	0	0	5	18	4	0
不安定プラーク (III b)	男	1 (0.1%)	0	0	0	0	1	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0
	計	1 (0.1%)	0	0	0	0	1	0	0
総頸動脈壁肥厚	男	34 (4.5%)	0	0	0	2	12	13	7
	女	6 (1.1%)	0	0	0	0	3	1	2
	計	40 (3.0%)	0	0	0	2	15	14	9
その他の所見	男	3 (0.4%)	0	0	0	1	0	2	0
	女	1 (0.2%)	0	0	0	0	1	0	0
	計	4 (0.3%)	0	0	0	1	1	2	0
検診者総数	男	760	0	1	29	180	336	161	53
	女	566	0	3	13	129	230	151	40
	計	1,326	0	4	42	309	566	312	93

頸部動脈超音波検査の計測値の性年齢別平均値

検査項目	総数		29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		
	男	女	男	女	男	女	男	女	男	女	男	女	男	女	
平均IMT 右・ mm	男	760	0.8	1	0.5	29	0.6	180	0.7	336	0.8	161	0.9	53	0.9
	女	566	0.7	3	0.5	13	0.5	129	0.6	230	0.7	151	0.8	40	0.8
	計	1,326	0.7	4	0.5	42	0.6	309	0.6	566	0.7	312	0.8	93	0.9
平均IMT 左・ mm	男	760	0.8	1	0.5	29	0.7	180	0.7	336	0.8	161	0.9	53	1.0
	女	566	0.7	3	0.5	13	0.5	129	0.6	230	0.7	151	0.8	40	0.8
	計	1,326	0.8	4	0.5	42	0.6	309	0.7	566	0.8	312	0.9	93	0.9
最大IMT 右・ mm	男	760	1.4	1	0.8	29	1.2	180	1.1	336	1.4	161	1.7	53	1.9
	女	566	1.3	3	0.8	13	0.8	129	1.1	230	1.3	151	1.4	40	1.5
	計	1,326	1.4	4	0.8	42	1.1	309	1.1	566	1.3	312	1.6	93	1.8
最大IMT 左・ mm	男	760	1.5	1	0.7	29	1.3	180	1.2	336	1.5	161	1.8	53	1.9
	女	566	1.3	3	0.8	13	0.9	129	1.1	230	1.3	151	1.5	40	1.6
	計	1,326	1.4	4	0.8	42	1.2	309	1.2	566	1.4	312	1.6	93	1.8

血圧脈波検査の性年齢別平均値

検査項目	総数		29歳以下		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		
	男	女	男	女	男	女	男	女	男	女	男	女	男	女	
ABI検査 右	男	738	1.2	1	1.1	29	1.1	179	1.1	321	1.2	159	1.2	49	1.2
	女	546	1.1	3	1.1	12	1.1	127	1.1	223	1.1	143	1.1	38	1.1
	計	1,284	1.1	4	1.1	41	1.1	306	1.1	544	1.1	302	1.1	87	1.2
ABI検査 左	男	738	1.2	1	1.1	29	1.1	179	1.1	321	1.2	159	1.2	49	1.2
	女	546	1.1	3	1.0	12	1.1	127	1.1	223	1.1	143	1.1	38	1.1
	計	1,284	1.1	4	1.1	41	1.1	306	1.1	544	1.1	302	1.1	87	1.1
PWV検査 右・ cm/s	男	738	1432.8	1	1131.0	29	1275.8	179	1291.9	321	1408.9	159	1569.7	49	1759.2
	女	546	1396.1	3	1099.7	12	1093.2	127	1201.8	223	1361.8	143	1562.4	38	1739.3
	計	1,284	1417.2	4	1107.5	41	1222.4	306	1254.5	544	1389.6	302	1566.2	87	1750.5
PWV検査 左・ cm/s	男	738	1443.9	1	1030.0	29	1289.2	179	1295.1	321	1421.2	159	1587.7	49	1769.2
	女	546	1410.1	3	1119.3	12	1125.9	127	1221.1	223	1365.7	143	1576.1	38	1790.7
	計	1,284	1429.5	4	1097.0	41	1241.4	306	1264.4	544	1398.4	302	1582.2	87	1778.6