

上部消化管検査判定結果

判定	総数		19歳以下	20-24歳	25-29歳	30-34歳	35-39歳	40-44歳	45-49歳	50-54歳	55-59歳	60-64歳	65-69歳	70-74歳	75-79歳	80歳以上
	男	女														
異常なし	男	12,960 (72.6%)	5	113	138	556	1,781	2,089	2,271	1,923	1,760	1,299	581	335	81	28
	女	7,420 (68.1%)	1	40	50	263	954	1,306	1,385	1,157	971	637	350	237	53	16
	計	20,380 (70.9%)	6	153	188	819	2,735	3,395	3,656	3,080	2,731	1,936	931	572	134	44
問題なし	男	2,095 (11.7%)	0	1	2	39	177	253	393	400	299	253	143	87	32	16
	女	1,543 (14.2%)	0	0	0	28	113	213	300	326	243	159	79	68	7	7
	計	3,638 (12.6%)	0	1	2	67	290	466	693	726	542	412	222	155	39	23
要経過観察	男	2,061 (11.5%)	0	1	2	46	150	200	294	341	363	274	207	130	34	19
	女	1,357 (12.5%)	0	0	0	24	93	178	243	228	209	181	100	75	20	6
	計	3,418 (11.9%)	0	1	2	70	243	378	537	569	572	455	307	205	54	25
要精密検査	男	682 (3.8%)	0	0	6	22	47	97	94	109	108	92	43	42	19	3
	女	557 (5.1%)	0	2	6	18	37	74	101	97	83	68	30	31	6	4
	計	1,239 (4.3%)	0	2	12	40	84	171	195	206	191	160	73	73	25	7
要治療	男	44 (0.2%)	0	0	0	0	3	10	9	7	1	6	5	2	0	1
	女	17 (0.2%)	0	0	0	0	1	2	7	3	2	0	2	0	0	0
	計	61 (0.2%)	0	0	0	0	4	12	16	10	3	6	7	2	0	1
治療中	男	19 (0.1%)	0	0	0	0	0	0	1	4	8	4	0	2	0	0
	女	5 (0.0%)	0	0	0	0	0	1	0	1	0	1	2	0	0	0
	計	24 (0.1%)	0	0	0	0	0	1	1	5	8	5	2	2	0	0
合計	男	17,861	5	115	148	663	2,158	2,649	3,062	2,784	2,539	1,928	979	598	166	67
	女	10,899	1	42	56	333	1,198	1,774	2,036	1,812	1,508	1,046	563	411	86	33
	計	28,760	6	157	204	996	3,356	4,423	5,098	4,596	4,047	2,974	1,542	1,009	252	100

胃内視鏡検査所見

所見	総数			19歳以下	20-24歳	25-29歳	30-34歳	35-39歳	40-44歳	45-49歳	50-54歳	55-59歳	60-64歳	65-69歳	70-74歳	75-79歳	80歳以上
	男	女	計														
食道裂孔ヘルニア	男	622	(26.5%)	0	0	0	11	48	95	111	127	114	74	27	13	1	1
	女	159	(11.1%)	0	0	0	5	9	29	31	28	28	17	7	4	1	0
	計	781	(20.7%)	0	0	0	16	57	124	142	155	142	91	34	17	2	1
逆流性食道炎	男	477	(20.3%)	0	0	0	6	28	63	88	112	80	65	28	4	2	1
	女	70	(4.9%)	0	0	0	0	3	14	11	16	9	10	3	3	1	0
	計	547	(14.5%)	0	0	0	6	31	77	99	128	89	75	31	7	3	1
バレット食道	男	34	(1.5%)	0	0	0	0	2	0	5	11	5	5	5	0	1	0
	女	11	(0.8%)	0	0	0	0	2	0	4	3	0	2	0	0	0	0
	計	45	(1.2%)	0	0	0	0	4	0	9	14	5	7	5	0	1	0
その他の所見 (食道所見)	男	96	(4.1%)	0	0	0	3	3	12	22	15	19	12	7	2	1	0
	女	40	(2.8%)	0	0	0	1	5	2	6	11	6	6	3	0	0	0
	計	136	(3.6%)	0	0	0	4	8	14	28	26	25	18	10	2	1	0
慢性胃炎	男	277	(11.8%)	0	0	0	3	15	15	39	70	61	46	16	12	0	0
	女	104	(7.3%)	0	0	0	1	5	9	15	29	22	17	4	2	0	0
	計	381	(10.1%)	0	0	0	4	20	24	54	99	83	63	20	14	0	0
萎縮性胃炎	男	143	(6.1%)	0	0	0	1	9	14	33	20	21	19	17	6	2	1
	女	85	(6.0%)	0	0	0	0	4	6	17	17	13	19	7	2	0	0
	計	228	(6.0%)	0	0	0	1	13	20	50	37	34	38	24	8	2	1
びらん性胃炎	男	122	(5.2%)	0	0	0	1	7	15	25	27	18	23	6	0	0	0
	女	42	(2.9%)	0	0	0	0	0	3	10	12	5	8	4	0	0	0
	計	164	(4.3%)	0	0	0	1	7	18	35	39	23	31	10	0	0	0
クオイボびらん	男	1	(0.0%)	0	0	0	0	0	0	0	0	1	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	計	1	(0.0%)	0	0	0	0	0	0	0	0	1	0	0	0	0	0
胃潰瘍	男	9	(0.4%)	0	0	0	0	0	2	1	2	1	2	1	0	0	0
	女	2	(0.1%)	0	0	0	0	0	1	0	0	0	0	1	0	0	0
	計	11	(0.3%)	0	0	0	0	0	3	1	2	1	2	2	0	0	0
胃潰瘍瘢痕	男	14	(0.6%)	0	0	0	1	1	0	4	4	0	2	0	2	0	0
	女	4	(0.3%)	0	0	0	0	0	1	1	0	0	2	0	0	0	0
	計	18	(0.5%)	0	0	0	1	1	1	5	4	0	4	0	2	0	0
胃底腺ポリープ	男	415	(17.7%)	0	0	0	2	22	52	85	93	68	64	17	10	1	1
	女	461	(32.3%)	0	0	0	5	44	73	100	102	72	40	21	3	1	0
	計	876	(23.2%)	0	0	0	7	66	125	185	195	140	104	38	13	2	1
除菌後胃炎	男	203	(8.7%)	0	0	0	0	8	22	24	43	41	34	24	7	0	0
	女	113	(7.9%)	0	0	0	2	3	10	22	22	18	20	9	7	0	0
	計	316	(8.4%)	0	0	0	2	11	32	46	65	59	54	33	14	0	0
表層性胃炎	男	15	(0.6%)	0	0	0	0	1	0	1	5	4	3	0	0	0	1
	女	14	(1.0%)	0	0	0	0	4	1	2	3	2	1	0	0	1	0
	計	29	(0.8%)	0	0	0	0	5	1	3	8	6	4	0	0	1	1
胃術後	男	6	(0.3%)	0	0	0	0	0	1	0	0	1	2	0	2	0	0
	女	2	(0.1%)	0	0	0	0	0	0	0	0	1	0	0	1	0	0
	計	8	(0.2%)	0	0	0	0	0	1	0	0	2	2	0	3	0	0
その他の所見 (胃所見)	男	388	(16.6%)	0	0	0	2	32	41	68	85	75	48	19	11	6	1
	女	142	(9.9%)	0	0	1	2	10	17	28	29	26	20	8	1	0	0
	計	530	(14.1%)	0	0	1	4	42	58	96	114	101	68	27	12	6	1
十二指腸潰瘍	男	3	(0.1%)	0	0	0	0	0	0	2	1	0	0	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	計	3	(0.1%)	0	0	0	0	0	0	2	1	0	0	0	0	0	0
十二指腸潰瘍瘢痕	男	59	(2.5%)	0	0	0	0	5	3	9	13	8	13	4	4	0	0
	女	18	(1.3%)	0	0	0	0	1	0	0	6	4	3	0	4	0	0
	計	77	(2.0%)	0	0	0	0	6	3	9	19	12	16	4	8	0	0
ポリープ	男	183	(7.8%)	0	0	0	3	15	19	36	37	33	29	5	4	1	1
	女	116	(8.1%)	0	0	0	4	8	24	25	27	18	10	0	0	0	0
	計	299	(7.9%)	0	0	0	7	23	43	61	64	51	39	5	4	1	1
憩室	男	5	(0.2%)	0	0	0	0	0	0	1	0	1	1	0	2	0	0
	女	6	(0.4%)	0	0	0	0	1	1	1	1	1	0	0	1	0	0
	計	11	(0.3%)	0	0	0	0	1	1	2	1	2	1	0	3	0	0
粘膜下腫瘍	男	61	(2.6%)	0	0	0	0	3	9	11	16	11	6	3	2	0	0
	女	22	(1.5%)	0	0	0	1	1	0	6	7	1	2	1	3	0	0
	計	83	(2.2%)	0	0	0	1	4	9	17	23	12	8	4	5	0	0
その他の所見 (十二指腸所見)	男	130	(5.5%)	0	0	1	1	7	14	20	29	23	16	15	4	0	0
	女	31	(2.2%)	0	0	0	0	2	1	6	14	3	5	0	0	0	0
	計	161	(4.3%)	0	0	1	1	9	15	26	43	26	21	15	4	0	0
検診者総数	男	2,344		0	0	1	39	202	318	456	493	387	287	107	45	6	3
	女	1,428		0	0	4	31	154	223	299	279	225	142	50	20	1	0
	計	3,772		0	0	5	70	356	541	755	772	612	429	157	65	7	3

胃部X線検査所見(1)

所見		総数		19歳以下	20-24歳	25-29歳	30-34歳	35-39歳	40-44歳	45-49歳	50-54歳	55-59歳	60-64歳	65-69歳	70-74歳	75-79歳	80歳以上
<b>胃 部</b>																	
伸展不良	男	7	(0.1%)	0	0	0	0	0	0	0	2	1	0	0	2	2	0
	女	2	(0.0%)	0	0	0	0	0	0	0	2	0	0	0	0	0	0
	計	9	(0.1%)	0	0	0	0	0	0	0	4	1	0	0	2	2	0
辺縁不整	男	18	(0.2%)	0	0	0	0	2	2	1	2	2	3	3	2	1	0
	女	3	(0.0%)	0	0	0	0	0	1	0	1	1	0	0	0	0	0
	計	21	(0.1%)	0	0	0	0	2	3	1	3	3	3	3	2	1	0
壁硬化	男	5	(0.0%)	0	0	0	0	1	0	0	0	3	0	1	0	0	0
	女	3	(0.0%)	0	0	0	0	0	0	1	0	1	0	0	1	0	0
	計	8	(0.0%)	0	0	0	0	1	0	1	0	4	0	1	1	0	0
変形	男	68	(0.6%)	0	0	0	1	1	4	11	7	10	18	7	4	5	0
	女	6	(0.1%)	0	0	0	0	2	0	1	1	0	0	1	1	0	0
	計	74	(0.4%)	0	0	0	1	3	4	12	8	10	18	8	5	5	0
粘膜不整	男	24	(0.2%)	0	0	0	0	1	3	3	3	6	3	1	3	1	0
	女	10	(0.1%)	0	0	0	1	1	0	4	0	0	0	1	3	0	0
	計	34	(0.2%)	0	0	0	1	2	3	7	3	6	3	2	6	1	0
レリーフ集中像	男	22	(0.2%)	0	0	0	0	2	1	2	4	3	3	6	0	1	0
	女	7	(0.1%)	0	0	0	1	1	2	0	0	1	1	0	1	0	0
	計	29	(0.2%)	0	0	0	1	3	3	2	4	4	4	6	1	1	0
レリーフ異常	男	2	(0.0%)	0	0	0	0	0	0	1	0	0	0	1	0	0	0
	女	2	(0.0%)	0	0	0	0	0	0	0	0	1	0	0	1	0	0
	計	4	(0.0%)	0	0	0	0	0	0	1	0	1	0	1	1	0	0
バリウム斑	男	22	(0.2%)	0	0	0	0	1	4	2	3	7	3	0	0	2	0
	女	9	(0.1%)	0	0	0	0	0	0	0	1	0	3	1	4	0	0
	計	31	(0.2%)	0	0	0	0	1	4	2	4	7	6	1	4	2	0
陰影欠損	男	2	(0.0%)	0	0	0	0	0	0	0	0	0	0	0	2	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	計	2	(0.0%)	0	0	0	0	0	0	0	0	0	0	0	2	0	0
胃炎の疑い	男	1,218	(11.3%)	0	1	1	29	87	112	142	179	164	142	165	122	51	23
	女	473	(6.9%)	0	0	0	3	20	34	49	63	58	88	62	69	20	7
	計	1,691	(9.5%)	0	1	1	32	107	146	191	242	222	230	227	191	71	30
胃潰瘍瘢痕の疑い	男	7	(0.1%)	0	0	0	0	0	0	2	1	1	1	0	0	2	0
	女	0	(0.0%)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	計	7	(0.0%)	0	0	0	0	0	0	2	1	1	1	0	0	2	0
胃潰瘍の疑い	男	22	(0.2%)	0	0	0	1	0	1	3	4	3	5	1	1	3	0
	女	3	(0.0%)	0	0	0	0	1	0	0	0	0	2	0	0	0	0
	計	25	(0.1%)	0	0	0	1	1	1	3	4	3	7	1	1	3	0
粗大皺壁の疑い	男	5	(0.0%)	0	0	0	0	0	0	1	1	1	0	0	2	0	0
	女	6	(0.1%)	0	0	0	1	0	0	1	1	0	1	1	1	0	0
	計	11	(0.1%)	0	0	0	1	0	0	2	2	1	1	1	3	0	0
陥凹性病変の疑い	男	3	(0.0%)	0	0	0	0	0	0	0	2	0	1	0	0	0	0
	女	0	(0.0%)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	計	3	(0.0%)	0	0	0	0	0	0	0	2	0	1	0	0	0	0
憩室の疑い	男	51	(0.5%)	0	0	0	2	4	5	9	3	6	7	7	6	2	0
	女	30	(0.4%)	0	0	0	1	2	2	5	3	4	2	3	6	2	0
	計	81	(0.5%)	0	0	0	3	6	7	14	6	10	9	10	12	4	0
術後胃	男	30	(0.3%)	0	0	0	0	0	0	1	3	2	9	3	7	1	4
	女	2	(0.0%)	0	0	0	0	0	0	0	0	1	1	0	0	0	0
	計	32	(0.2%)	0	0	0	0	0	0	1	3	3	10	3	7	1	4
ヒロリ菌感染疑い	男	47	(0.4%)	0	0	0	7	8	11	8	7	3	2	1	0	0	0
	女	26	(0.4%)	0	0	0	4	3	3	5	1	4	2	4	0	0	0
	計	73	(0.4%)	0	0	0	11	11	14	13	8	7	4	5	0	0	0
その他の所見	男	49	(0.5%)	0	0	0	3	1	4	10	6	10	1	7	4	1	2
	女	27	(0.4%)	0	1	0	0	1	2	2	2	3	6	5	3	1	1
	計	76	(0.4%)	0	1	0	3	2	6	12	8	13	7	12	7	2	3

胃部X線検査所見(2)

所見	総数	19歳以下	20-24歳	25-29歳	30-34歳	35-39歳	40-44歳	45-49歳	50-54歳	55-59歳	60-64歳	65-69歳	70-74歳	75-79歳	80歳以上	
<b>食道部</b>																
狭窄	男	2 (0.0%)	0	0	0	0	0	0	0	0	1	1	0	0	0	
	女	0 (0.0%)	0	0	0	0	0	0	0	0	0	0	0	0	0	
	計	2 (0.0%)	0	0	0	0	0	0	0	0	1	1	0	0	0	
食道裂孔ヘルニア	男	69 (0.6%)	0	0	0	0	0	2	6	9	8	6	12	13	8	5
	女	30 (0.4%)	0	0	0	0	1	0	1	3	6	6	9	4	0	
	計	99 (0.6%)	0	0	0	0	1	2	6	10	11	12	18	22	12	5
食道炎の疑い	男	60 (0.6%)	0	0	0	0	3	8	8	3	9	11	9	7	1	1
	女	15 (0.2%)	0	0	0	0	0	0	1	2	5	3	1	1	2	0
	計	75 (0.4%)	0	0	0	0	3	8	9	5	14	14	10	8	3	1
その他の所見	男	14 (0.1%)	0	0	0	1	0	1	1	3	2	2	2	1	1	0
	女	7 (0.1%)	0	0	0	0	0	2	0	1	0	1	2	1	0	0
	計	21 (0.1%)	0	0	0	1	0	3	1	4	2	3	4	2	1	0
<b>十二指腸部</b>																
十二指腸球部変形	男	213 (2.0%)	0	0	0	6	15	22	32	40	28	31	19	12	6	2
	女	23 (0.3%)	0	0	0	0	2	2	3	2	1	3	5	4	0	1
	計	236 (1.3%)	0	0	0	6	17	24	35	42	29	34	24	16	6	3
ニッシュェ	男	4 (0.0%)	0	0	0	0	1	0	0	1	1	1	0	0	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	計	4 (0.0%)	0	0	0	0	1	0	0	1	1	1	0	0	0	0
十二指腸潰瘍の疑い	男	10 (0.1%)	0	0	0	1	1	1	0	1	2	3	0	1	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	計	10 (0.1%)	0	0	0	1	1	1	0	1	2	3	0	1	0	0
十二指腸潰瘍瘢痕の疑い	男	18 (0.2%)	0	0	0	2	1	1	2	2	2	4	3	1	0	0
	女	2 (0.0%)	0	0	0	0	0	0	0	0	1	1	0	0	0	
	計	20 (0.1%)	0	0	0	2	1	1	2	2	2	5	4	1	0	0
十二指腸憩室の疑い	男	2 (0.0%)	0	0	0	0	0	0	1	0	0	0	1	0	0	0
	女	7 (0.1%)	0	0	0	0	0	1	0	1	0	0	1	3	1	0
	計	9 (0.1%)	0	0	0	0	0	1	1	1	0	0	2	3	1	0
隆起性病変の疑い	男	1,083 (10.0%)	0	1	2	25	113	136	183	167	169	121	85	58	15	8
	女	1,566 (22.7%)	0	0	0	37	105	241	323	292	248	166	79	60	9	6
	計	2,649 (15.0%)	0	1	2	62	218	377	506	459	417	287	164	118	24	14
造影不良	男	21 (0.2%)	0	0	0	1	2	0	8	6	1	3	0	0	0	0
	女	2 (0.0%)	0	0	0	0	0	0	1	0	0	0	1	0	0	0
	計	23 (0.1%)	0	0	0	1	2	0	9	6	1	3	1	0	0	0
その他の所見	男	2 (0.0%)	0	0	0	0	0	0	0	0	0	0	1	1	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	計	2 (0.0%)	0	0	0	0	0	0	0	0	0	0	1	1	0	0
検診者総数	男	10,824	3	33	44	507	1,303	1,586	1,796	1,551	1,362	1,110	799	513	155	62
	女	6,888	0	23	18	267	655	1,047	1,162	1,081	961	730	467	370	79	28
	計	17,712	3	56	62	774	1,958	2,633	2,958	2,632	2,323	1,840	1,266	883	234	90